

# A NOTE ON NEW CLASSES OF INFINITELY DIVISIBLE DISTRIBUTIONS ON $\mathbb{R}^d$

MAKOTO MAEJIMA

*Keio University*

email: maejima@math.keio.ac.jp

GENTA NAKAHARA

*Keio University*

AMS 2000 Subject classification: 60E07

Keywords: infinitely divisible distribution on  $\mathbb{R}^d$ , stochastic integral representation.

## *Abstract*

This paper introduces and studies a family of new classes of infinitely divisible distributions on  $\mathbb{R}^d$  with two parameters. Depending on parameters, these classes connect the Goldie–Steutel–Bondesson class and the class of generalized type  $G$  distributions, connect the Thorin class and the class  $M$ , connect the class  $M$  and the class of generalized type  $G$  distributions. These classes are characterized by stochastic integral representations with respect to Lévy processes.

## 1. INTRODUCTION

Let  $I(\mathbb{R}^d)$  be the class of all infinitely divisible distributions on  $\mathbb{R}^d$ .  $\widehat{\mu}(z)$ ,  $z \in \mathbb{R}^d$ , denotes the characteristic function of  $\mu \in I(\mathbb{R}^d)$  and  $|x|$  denotes the Euclidean norm of  $x \in \mathbb{R}^d$ . We use the Lévy–Khintchine triplet  $(A, \nu, \gamma)$  of  $\mu \in I(\mathbb{R}^d)$  in the sense that

$$\widehat{\mu}(z) = \exp \left\{ -2^{-1} \langle z, Az \rangle + i \langle \gamma, z \rangle + \int_{\mathbb{R}^d} (e^{i \langle z, x \rangle} - 1 - i \langle z, x \rangle (1 + |x|^2)^{-1}) \nu(dx) \right\},$$

$z \in \mathbb{R}^d$ , where  $A$  is a symmetric nonnegative-definite  $d \times d$  matrix,  $\gamma \in \mathbb{R}^d$  and  $\nu$  is a measure (called the Lévy measure) on  $\mathbb{R}^d$  satisfying

$$\nu(\{0\}) = 0 \quad \text{and} \quad \int_{\mathbb{R}^d} (|x|^2 \wedge 1) \nu(dx) < \infty.$$

The following polar decomposition is a basic result on the Lévy measure of  $\mu \in I(\mathbb{R}^d)$ . Let  $\nu$  be the Lévy measure of some  $\mu \in I(\mathbb{R}^d)$  with  $0 < \nu(\mathbb{R}^d) \leq \infty$ . Then there exist a measure  $\lambda$  on  $S = \{x \in \mathbb{R}^d : |x| = 1\}$  with  $0 < \lambda(S) \leq \infty$  and a family  $\{\nu_\xi : \xi \in S\}$  of measures on  $(0, \infty)$  such that  $\nu_\xi(B)$  is measurable in  $\xi$  for each

$B \in \mathcal{B}((0, \infty))$ ,  $0 < \nu_\xi((0, \infty)) \leq \infty$  for each  $\xi \in S$ , and

$$(1.1) \quad \nu(B) = \int_S \lambda(d\xi) \int_0^\infty 1_B(r\xi) \nu_\xi(dr), \quad B \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}).$$

Here  $\lambda$  and  $\{\nu_\xi\}$  are uniquely determined by  $\nu$  up to multiplication of measurable functions  $c(\xi)$  and  $\frac{1}{c(\xi)}$ , respectively, with  $0 < c(\xi) < \infty$ . We say that  $\nu$  has the polar decomposition  $(\lambda, \nu_\xi)$  and  $\nu_\xi$  is called the radial component of  $\nu$ . (See, e.g., Barndorff-Nielsen et al. (2006), Lemma 2.1.)

A real-valued function  $f$  defined on  $(0, \infty)$  is said to be completely monotone if it has derivatives  $f^{(n)}$  of all orders and for each  $n = 0, 1, 2, \dots$ ,  $(-1)^n f^{(n)}(r) \geq 0, r > 0$ . Bernstein's theorem says that  $f$  on  $(0, \infty)$  is completely monotone if and only if there exists a (not necessarily finite) measure  $Q$  on  $[0, \infty)$  such that  $f(r) = \int_{[0, \infty)} e^{-ru} Q(du)$ . (See, e.g., Feller (1966), p.439.)

In this paper, we introduce and study the following classes.

**Definition 1.1.** (The class  $J_{\alpha, \beta}(\mathbb{R}^d)$ .) Let  $\alpha < 2$  and  $\beta > 0$ . We say that  $\mu \in I(\mathbb{R}^d)$  belongs to the class  $J_{\alpha, \beta}(\mathbb{R}^d)$  if  $\nu = 0$  or  $\nu \neq 0$  and, in case  $\nu \neq 0$ ,  $\nu_\xi$  in (1.1) has expression

$$(1.2) \quad \nu_\xi(dr) = r^{-\alpha-1} g_\xi(r^\beta) dr, \quad r > 0,$$

where  $g_\xi(x)$  is measurable in  $\xi$ , is completely monotone in  $x$  on  $(0, \infty)$   $\lambda$ -a.e.  $\xi$ , not identically zero and  $\lim_{x \rightarrow \infty} g_\xi(x) = 0$   $\lambda$ -a.e.  $\xi$ .

**Remark 1.2.** If  $\alpha \leq 0$ , then automatically  $\lim_{x \rightarrow \infty} g_\xi(x) = 0$   $\lambda$ -a.e.  $\xi$ , because of the finiteness of  $\int_{|x| > 1} \nu(dx)$ . So, when we consider the classes  $B(\mathbb{R}^d), G(\mathbb{R}^d), T(\mathbb{R}^d)$  and  $M(\mathbb{R}^d)$  appearing later, we do not have to write this condition explicitly.

**Remark 1.3.** The integrability condition of the Lévy measure  $\int_{\mathbb{R}^d} (|x|^2 \wedge 1) \nu(dx) < \infty$  implies that

$$(1.3) \quad \int_0^\infty (r^2 \wedge 1) r^{-\alpha-1} g_\xi(r^\beta) dr < \infty, \quad \lambda\text{-a.e. } \xi,$$

so we do not have to assume (1.3) in the definition. It is automatically satisfied.

**Remark 1.4.** The classes  $J_{\alpha, 1}(\mathbb{R}^d), \alpha < 2$ , are studied in Sato (2006b).

Before mentioning our motivation of this study, we state a general result on the relations among the classes  $J_{\alpha, \beta}(\mathbb{R}^d), \alpha < 2, \beta > 0$ .

**Theorem 1.5.** (i) Fix  $\alpha < 2$  and let  $0 < \beta_1 < \beta_2$ . Then

$$J_{\alpha, \beta_1}(\mathbb{R}^d) \subset J_{\alpha, \beta_2}(\mathbb{R}^d).$$

(ii) Fix  $\beta > 0$  and let  $\alpha_1 < \alpha_2 < 2$ . Then

$$J_{\alpha_2, \beta}(\mathbb{R}^d) \subset J_{\alpha_1, \beta}(\mathbb{R}^d).$$

*Proof.* For the proof of (i), we need the following lemma.

**Lemma 1.6.** (See Feller (1966), p.441, Corollary 2.) *Let  $\phi$  be a completely monotone function on  $(0, \infty)$  and let  $\psi$  be a nonnegative function on  $(0, \infty)$  whose derivative is completely monotone. Then  $\phi(\psi)$  is completely monotone.*

Let  $h_\xi(x) = g_\xi(x^{\beta_1/\beta_2}), x > 0$ , where  $g_\xi$  is the one in (1.2), which is completely monotone on  $(0, \infty)$ . Since  $\psi(x) = x^{\beta_1/\beta_2}, x > 0$ , has a completely monotone derivative, it follows from Lemma 1.6 that  $h_\xi(x)$  is completely monotone. Suppose  $\mu \in J_{\alpha, \beta_1}(\mathbb{R}^d)$  and let  $g_\xi$  be the one in (1.2). Since  $g_\xi(r^{\beta_1}) = h_\xi(r^{\beta_2})$ , where  $h_\xi$  is completely monotone as has been just shown above, we have  $\mu \in J_{\alpha, \beta_2}(\mathbb{R}^d)$ . This proves (i).

To prove (ii), suppose that  $\mu \in J_{\alpha_2, \beta}(\mathbb{R}^d)$ . Then  $\nu_\xi(dr) = r^{-\alpha_2-1}g_\xi(r^\beta)dr, r > 0$ , as in (1.2), where  $g_\xi$  is completely monotone on  $(0, \infty)$   $\lambda$ -a.e.  $\xi$ . Note that

$$h_\xi(x) = x^{-(\alpha_2-\alpha_1)/\beta}g_\xi(x)$$

is completely monotone, because  $x^{-p}, p > 0$ , is completely monotone and the product of two completely monotone functions is also completely monotone. We now have

$$\nu_\xi(dr) = r^{-\alpha_2-1}g_\xi(r^\beta)dr = r^{-\alpha_1-1}h_\xi(r^\beta)dr,$$

and thus  $\mu$  also belongs to  $J_{\alpha_1, \beta}(\mathbb{R}^d)$ . This proves (ii).  $\square$

The motivations for studying the classes  $J_{\alpha, \beta}(\mathbb{R}^d)$  are the following.

**I.** The classes connecting the Goldie–Steutel–Bondesson class and the class of generalized type  $G$  distributions.

Let  $\alpha = -1$  and consider the classes  $J_{-1, \beta}(\mathbb{R}^d), \beta > 0$ .

A distribution  $\mu \in I(\mathbb{R}^d)$  is said to be of generalized type  $G$  if  $\nu_\xi$  in (1.2) has expression  $\nu_\xi(dr) = g_\xi(r^2)dr$  for some completely monotone function  $g_\xi$  on  $(0, \infty)$ , and denote by  $G(\mathbb{R}^d)$  the class of all generalized type  $G$  distributions on  $\mathbb{R}^d$ . Let  $I_{\text{sym}}(\mathbb{R}^d) = \{\mu \in I(\mathbb{R}^d) \mid \mu \text{ is symmetric in the sense that } \mu(B) = \mu(-B), B \in \mathcal{B}(\mathbb{R}^d)\}$ .

**Remark 1.7.** A distribution  $\mu \in G(\mathbb{R}^d) \cap I_{\text{sym}}(\mathbb{R}^d)$  is a so-called type  $G$  distribution, which is, in one dimension, a variance mixture of the standard normal distribution with a positive infinitely divisible mixing distribution.

**Remark 1.8.**  $G(\mathbb{R}^d) = J_{-1,2}(\mathbb{R}^d)$ .

**Remark 1.9.** The Goldie-Steutel-Bondesson class denoted by  $B(\mathbb{R}^d)$  is  $J_{-1,1}(\mathbb{R}^d)$ . (For details on  $B(\mathbb{R}^d)$ , see Barndorff-Nielsen et al. (2006).)

Therefore, by Theorem 1.5 (i) with  $\alpha = -1$ , for  $1 < \beta < 2$ ,

$$B(\mathbb{R}^d) \subset J_{-1,\beta}(\mathbb{R}^d) \subset G(\mathbb{R}^d),$$

and hence  $\{J_{-1,\beta}(\mathbb{R}^d), 1 \leq \beta \leq 2\}$  is a family of classes of infinitely divisible distributions on  $\mathbb{R}^d$  connecting  $B(\mathbb{R}^d)$  and  $G(\mathbb{R}^d)$  with continuous parameter  $\beta \in [1, 2]$ .

**II.** The classes connecting the Thorin class and the class  $M(\mathbb{R}^d)$ .

Let  $\alpha = 0$  and consider the classes  $J_{0,\beta}(\mathbb{R}^d)$ ,  $\beta > 0$ .

**Remark 1.10.** The Thorin class denoted by  $T(\mathbb{R}^d)$  is  $J_{0,1}(\mathbb{R}^d)$ . (For details on  $T(\mathbb{R}^d)$ , see also Barndorff-Nielsen et al. (2006).)

**Remark 1.11.** The class  $M(\mathbb{R}^d)$  is defined by  $J_{0,2}(\mathbb{R}^d)$ . (The class  $M(\mathbb{R}^d) \cap I_{\text{sym}}(\mathbb{R}^d)$  is studied in Aoyama et al. (2008).)

By Theorem 1.5 (i) with  $\alpha = 0$ , for  $1 < \beta < 2$ ,

$$T(\mathbb{R}^d) \subset J_{0,\beta}(\mathbb{R}^d) \subset M(\mathbb{R}^d),$$

and hence  $\{J_{0,\beta}(\mathbb{R}^d), 1 \leq \beta \leq 2\}$  is a family of classes of infinitely divisible distributions on  $\mathbb{R}^d$  connecting  $T(\mathbb{R}^d)$  and  $M(\mathbb{R}^d)$  with continuous parameter  $\beta \in [1, 2]$ .

**III.** The classes connecting the classes  $M(\mathbb{R}^d)$  and  $G(\mathbb{R}^d)$ .

Let  $\beta = 2$  and consider the classes  $J_{\alpha,2}(\mathbb{R}^d)$ ,  $\alpha < 2$ . Then, by Theorem 1.5 (ii) with  $\beta = 2$ , for  $-1 \leq \alpha \leq 0$

$$M(\mathbb{R}^d) \subset J_{\alpha,2}(\mathbb{R}^d) \subset G(\mathbb{R}^d),$$

and hence  $\{J_{\alpha,2}(\mathbb{R}^d), -1 \leq \alpha \leq 0\}$  is a family of classes of infinitely divisible distributions on  $\mathbb{R}^d$  connecting  $M(\mathbb{R}^d)$  and  $G(\mathbb{R}^d)$  with continuous parameter  $\alpha \in [-1, 0]$ .

**IV.** The classes connecting the classes  $T(\mathbb{R}^d)$  and  $B(\mathbb{R}^d)$ .

Let  $\beta = 1$  and consider the classes  $J_{\alpha,1}(\mathbb{R}^d)$ ,  $\alpha < 2$ . Then, by Theorem 1.5 (ii) with  $\beta = 1$ , for  $-1 \leq \alpha \leq 0$

$$T(\mathbb{R}^d) \subset J_{\alpha,1}(\mathbb{R}^d) \subset B(\mathbb{R}^d),$$

and hence  $\{J_{\alpha,1}(\mathbb{R}^d), -1 \leq \alpha \leq 0\}$  is a family of classes of infinitely divisible distributions on  $\mathbb{R}^d$  connecting  $T(\mathbb{R}^d)$  and  $B(\mathbb{R}^d)$  with continuous parameter  $\alpha \in [-1, 0]$ . (This fact is already mentioned in Sato (2006b).)

## 2. STOCHASTIC INTEGRAL CHARACTERIZATIONS FOR $J_{\alpha,\beta}(\mathbb{R}^d)$

The purpose of this paper is to characterize the classes  $J_{\alpha,\beta}(\mathbb{R}^d)$  by stochastic integral representations. For that, we first define mappings from  $I(\mathbb{R}^d)$  into  $I(\mathbb{R}^d)$  and investigate the domains of those mappings.

We introduce the following function  $G_{\alpha,\beta}(u)$ . For  $\alpha < 2$  and  $\beta > 0$ , let

$$G_{\alpha,\beta}(u) = \int_u^\infty x^{-\alpha-1} e^{-x^\beta} dx, \quad u \geq 0,$$

and let  $G_{\alpha,\beta}^*(t)$  be the inverse function of  $G_{\alpha,\beta}(u)$ , that is,  $t = G_{\alpha,\beta}(u)$  if and only if  $u = G_{\alpha,\beta}^*(t)$ . Let  $\{X_t^{(\mu)}\}$  be a Lévy process on  $\mathbb{R}^d$  with the law  $\mu \in I(\mathbb{R}^d)$  at  $t = 1$ .

We consider the stochastic integrals

$$\int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dX_t^{(\mu)}, \quad \text{where } G_{\alpha,\beta}(0) = \begin{cases} \beta^{-1} \Gamma(-\alpha\beta^{-1}), & \text{if } \alpha < 0, \\ \infty, & \text{if } \alpha \geq 0. \end{cases}$$

As to the definition of stochastic integrals of non-random measurable functions  $f$  which are  $\int_0^T f(t) dX_t^{(\mu)}$ ,  $T < \infty$ ,  $\mu \in I(\mathbb{R}^d)$ , we follow the definition in Sato (2004, 2006a), whose idea is to define a stochastic integral with respect to  $\mathbb{R}^d$ -valued independently scattered random measure induced by a Lévy process on  $\mathbb{R}^d$ . The improper stochastic integral  $\int_0^\infty f(t) dX_t^{(\mu)}$  is defined as the limit in probability of  $\int_0^T f(t) dX_t^{(\mu)}$  as  $T \rightarrow \infty$  whenever the limit exists. See also Sato (2006b). In the following,  $\mathcal{L}(X)$  stands for “the law of  $X$ ”. If we write

$$\Psi_{\alpha,\beta}(\mu) = \mathcal{L} \left( \int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dX_t^{(\mu)} \right),$$

then  $\Psi_{\alpha,\beta}$  can be considered as a mapping with domain  $\mathfrak{D}(\Psi_{\alpha,\beta})$  being the class of  $\mu \in I(\mathbb{R}^d)$  for which  $\int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dX_t^{(\mu)}$  is definable.

**Theorem 2.1.** *If  $\alpha < 0$ , then  $\mathfrak{D}(\Psi_{\alpha,\beta}) = I(\mathbb{R}^d)$ .*

*Proof.* By Proposition 3.4 in Sato (2006a), since  $G_{\alpha,\beta}(0) < \infty$  for  $\alpha < 0$ , if  $\int_0^{G_{\alpha,\beta}(0)} (G_{\alpha,\beta}^*(t))^2 dt < \infty$ , then  $\int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dX_t^{(\mu)}$  is well-defined. Actually,

$$\int_0^{G_{\alpha,\beta}(0)} (G_{\alpha,\beta}^*(t))^2 dt = - \int_0^\infty u^2 dG_{\alpha,\beta}(u) = \int_0^\infty u^{1-\alpha} e^{-u^\beta} du < \infty.$$

□

To determine the domain of  $\Psi_{\alpha,\beta}$ ,  $\alpha \geq 0$ , we need the following result by Sato (2006b). In the following,  $a(t) \sim b(t)$  means that  $\lim_{t \rightarrow \infty} a(t)/b(t) = 1$ ,  $a(t) \asymp b(t)$  means that  $0 < \liminf_{t \rightarrow \infty} a(t)/b(t) \leq \limsup_{t \rightarrow \infty} a(t)/b(t) < \infty$  and  $I_{\log}(\mathbb{R}^d) = \{\mu \in I(\mathbb{R}^d) \mid \int_{\mathbb{R}^d} \log^+ |x| \mu(dx) < \infty\}$ , where  $\log^+ |x| = (\log |x|) \vee 0$ .

**Proposition 2.2.** (Sato (2006b), Theorems 2.4 and 2.8.) *Let  $p \geq 0$ . Denote*

$$\Phi_{\varphi_p}(\mu) = \mathcal{L} \left( \int_0^\infty \varphi_p(t) dX_t^{(\mu)} \right).$$

*Suppose that  $\varphi_p$  is locally square-integrable with respect to Lebesgue measure on  $[0, \infty)$  and satisfies*

- (1)  $\varphi_0(t) \asymp e^{-ct}$  as  $t \rightarrow \infty$  with some  $c > 0$ ,
- (2)  $\varphi_p(t) \asymp t^{-1/p}$  as  $t \rightarrow \infty$  for  $p \in (0, 1) \cup (1, \infty)$ ,
- (3)  $\varphi_1(t) \asymp t^{-1}$  as  $t \rightarrow \infty$  and for some  $t_0 > 0, c > 0$  and  $\psi(t), \varphi_1(t) = t^{-1}\psi(t)$  for  $t > t_0$  with  $\int_{t_0}^\infty t^{-1} |\psi(t) - c| dt < \infty$ .

*Then*

- (i) *If  $p = 0$ , then  $\mathfrak{D}(\Phi_{\varphi_0}) = I_{\log}(\mathbb{R}^d)$ .*
- (ii) *If  $0 < p < 1$ , then  $\mathfrak{D}(\Phi_{\varphi_p}) = \{\mu \in I(\mathbb{R}^d) : \int_{\mathbb{R}^d} |x|^p \mu(dx) < \infty\} =: I_p(\mathbb{R}^d)$ .*
- (iii) *If  $p = 1$ , then  $\mathfrak{D}(\Phi_{\varphi_1}) = \{\mu \in I(\mathbb{R}^d) : \int_{\mathbb{R}^d} |x| \mu(dx) < \infty$   
 $\lim_{T \rightarrow \infty} \int_{t_0}^T t^{-1} dt \int_{|x|>t} x \nu(dx)$  exists in  $\mathbb{R}^d, \int_{\mathbb{R}^d} x \mu(dx) = 0\} =: I_1^*(\mathbb{R}^d)$ .*
- (iv) *If  $1 < p < 2$ , then  $\mathfrak{D}(\Phi_{\varphi_p}) = \{\mu \in I(\mathbb{R}^d) : \int_{\mathbb{R}^d} |x|^p \mu(dx) < \infty, \int_{\mathbb{R}^d} x \mu(dx) = 0\}$   
 $=: I_p^0(\mathbb{R}^d)$ .*
- (v) *If  $p \geq 2$ , then  $\mathfrak{D}(\Phi_{\varphi_p}) = \{\delta_0\}$ , where  $\delta_0$  is the distribution with the total mass at 0.*

We apply Proposition 2.2 to our problem. First we note that when  $\alpha < 2$ ,  $G_{\alpha,\beta}^*(t)$  is locally square-integrable with respect to Lebesgue measure on  $[0, \infty)$ .

**Theorem 2.3.** (Case  $\alpha = 0$ .)  $\mathfrak{D}(\Psi_{0,\beta}) = I_{\log}(\mathbb{R}^d)$ .

*Proof.* Note that  $t(= G_{\alpha,\beta}(u)) \uparrow \infty$  if and only if  $u(= G_{\alpha,\beta}^*(t)) \downarrow 0$ , when  $\alpha \geq 0$ . It is enough to show that for some  $C_1 \in (0, \infty)$ ,  $u \sim C_1 e^{-t}$  as  $t \rightarrow \infty$ . We have

$$\begin{aligned} \frac{u}{e^{-t}} &= \frac{u}{\exp\{-G_{0,\beta}(u)\}} = \exp\{G_{0,\beta}(u) + \log u\} = \exp\left\{\int_u^\infty x^{-1} e^{-x^\beta} dx + \log u\right\} \\ &= \exp\left\{\beta^{-1} \int_{u^\beta}^\infty y^{-1} e^{-y} dy - \beta^{-1} \int_{u^\beta}^1 y^{-1} dy\right\} \end{aligned}$$

$$= \exp \left\{ \beta^{-1} \int_{u^\beta}^1 y^{-1}(e^{-y} - 1)dy + \beta^{-1} \int_1^\infty y^{-1}e^{-y}dy \right\} \rightarrow C_1,$$

say, as  $u \downarrow 0$ . Hence  $u \sim C_1 e^{-t}$  as  $t \rightarrow \infty$ , and the condition (1) of Proposition 2.2 is satisfied. Thus Proposition 2.2 (i) gives us the assertion.  $\square$

**Theorem 2.4.** (Case  $\alpha \in (0, \infty)$ .)

- (i) If  $0 < \alpha < 1$ , then  $\mathfrak{D}(\Psi_{\alpha,\beta}) = I_\alpha(\mathbb{R}^d)$ .
- (ii) If  $\alpha = 1$ , then  $\mathfrak{D}(\Psi_{1,\beta}) = I_1^*(\mathbb{R}^d)$ .
- (iii) If  $1 < \alpha < 2$ , then  $\mathfrak{D}(\Psi_{\alpha,\beta}) = I_\alpha^0(\mathbb{R}^d)$ .
- (iv) If  $\alpha \geq 2$ , then  $\mathfrak{D}(\Psi_{\alpha,\beta}) = \{\delta_0\}$ .

*Proof.* (i) and (iii). It is enough to show that  $u \sim C_2 t^{-1/\alpha}$  as  $t \rightarrow \infty$  for some  $C_2 \in (0, \infty)$ . We have, as  $t \rightarrow \infty$  (equivalently  $u \downarrow 0$ ), for some  $C_3 \in (0, \infty)$ ,

$$\frac{u}{t^{-1/\alpha}} = \frac{u}{(G_{\alpha,\beta}(u))^{-1/\alpha}} = \frac{u}{(\beta^{-1} \int_{u^\beta}^\infty y^{-(\alpha/\beta)-1} e^{-y} dy)^{-1/\alpha}} \sim \frac{u}{(C_3 u^{-\alpha})^{-1/\alpha}} = C_3^{1/\alpha} =: C_2,$$

and the condition (2) of Proposition 2.3 is satisfied. Thus Proposition 2.3 (ii) and (iv) give us the assertions.

(ii). Suppose  $\beta \neq 1$ . (The case  $\beta = 1$  is proved in Sato (2006b).) We first have

$$\begin{aligned} G_{1,\beta}(u) &= \int_u^\infty x^{-2} e^{-x^\beta} dx = \int_u^\infty x^{-2} dx + \int_u^\infty x^{-2}(e^{-x^\beta} - 1)dx \\ &= \int_u^\infty x^{-2} dx + \int_u^1 x^{-2}(e^{-x^\beta} - 1 + x^\beta)du - \int_u^1 x^{-2+\beta} dx + \int_1^\infty x^{-2}(e^{-x^\beta} - 1)dx \\ &= u^{-1} + (\beta - 1)^{-1} u^{-1+\beta} + O(1), \quad u \downarrow 0. \end{aligned}$$

Thus

$$t = G_{1,\beta}^*(t)^{-1} + (\beta - 1)^{-1} G_{1,\beta}^*(t)^{-1+\beta} + O(1), \quad t \rightarrow \infty.$$

Therefore,

$$(2.1) \quad G_{1,\beta}^*(t) = t^{-1} + (\beta - 1)^{-1} t^{-1} G_{1,\beta}^*(t)^\beta + O(t^{-1} G_{1,\beta}^*(t)), \quad t \rightarrow \infty.$$

We have shown in (i) and (iii) that  $u \sim C_2 t^{-1/\alpha}$ , but this is also true for  $\alpha = 1$ . Hence

$$(2.2) \quad u = G_{1,\beta}^*(t) = C_2 t^{-1}(1 + o(1)), \quad t \rightarrow \infty.$$

By substituting (2.2) into (2.1), we have

$$\begin{aligned} G_{1,\beta}^*(t) &= t^{-1} + C_2^\beta (\beta - 1)^{-1} t^{-1-\beta} + t^{-1} a(t), \quad t \rightarrow \infty, \\ &= t^{-1} \left( 1 + C_2^\beta (\beta - 1)^{-1} t^{-\beta} + a(t) \right), \quad t \rightarrow \infty, \end{aligned}$$

where

$$a(t) = \begin{cases} o(t^{-\beta}), & t \rightarrow \infty, & \text{when } 0 < \beta < 1, \\ O(t^{-1}), & t \rightarrow \infty, & \text{when } \beta > 1. \end{cases}$$

Thus

$$G_{1,\beta}^*(t) = t^{-1}\psi(t),$$

where

$$\psi(t) := 1 + C_2^\beta(\beta - 1)^{-1}t^{-\beta} + a(t),$$

and

$$\int_1^\infty t^{-1}|\psi(t) - 1|dt = \int_1^\infty t^{-1}|C_2^\beta(\beta - 1)^{-1}t^{-\beta} + a(t)|dt < \infty.$$

Thus the condition (3) of Proposition 2.2 is satisfied with  $t_0 = 1$  and  $c = 1$ , and Proposition 2.2 (iii) gives us the assertion (iii).

(iv) The same as in Sato (2006b).  $\square$

We now calculate the Lévy measure of  $\tilde{\mu} = \Psi_{\alpha,\beta}(\mu)$ , and note that the mapping  $\Psi_{\alpha,\beta}$  is one-to-one.

**Lemma 2.5.** *Let  $\alpha < 2$  and  $\beta > 0$ . Let  $\mu \in \mathfrak{D}(\Psi_{\alpha,\beta})$  and  $\tilde{\mu} = \Psi_{\alpha,\beta}(\mu)$ , and let  $\nu$  and  $\tilde{\nu}$  be the Lévy measures of  $\mu$  and  $\tilde{\mu}$ , respectively.*

(1) *We have*

$$(2.3) \quad \tilde{\nu}(B) = \int_0^\infty \nu(s^{-1}B)s^{-\alpha-1}e^{-s^\beta} ds, \quad B \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}).$$

(2) *If  $\nu \neq 0$ , and  $\nu$  has polar decomposition  $(\lambda, \nu_\xi)$ , then a polar decomposition of  $\tilde{\nu} = (\tilde{\lambda}, \tilde{\nu}_\xi)$  is given by  $\tilde{\lambda} = \lambda$  and  $\tilde{\nu}_\xi(dr) = r^{-\alpha-1}\tilde{g}_\xi(r^\beta)dr$ , where*

$$(2.4) \quad \tilde{g}_\xi(u) = \int_0^\infty r^\alpha e^{-u/r^\beta} \nu_\xi(dr).$$

(3)  *$\tilde{g}_\xi$  in (2.4) satisfies the requirements of  $g_\xi$  in (1.2).*

*Proof.* Suppose  $\mu \in \mathfrak{D}(\Psi_{\alpha,\beta})$  and  $\tilde{\mu} = \Psi_{\alpha,\beta}(\mu)$ .

(1) We see that (by using Proposition 2.6 of Sato (2006b)),

$$\begin{aligned} \tilde{\nu}(B) &= \int_0^{G_{\alpha,\beta}^{(0)}} dt \int_{\mathbb{R}^d} 1_B(xG_{\alpha,\beta}^*(t))\nu(dx) = - \int_0^\infty dG_{\alpha,\beta}(s) \int_{\mathbb{R}^d} 1_B(xs)\nu(dx) \\ &= \int_0^\infty s^{-\alpha-1}e^{-s^\beta} ds \int_{\mathbb{R}^d} 1_{s^{-1}B}(x)\nu(dx) = \int_0^\infty \nu(s^{-1}B)s^{-\alpha-1}e^{-s^\beta} ds, \end{aligned}$$

which is (2.3).

(2) Next assume that  $\nu \neq 0$  and  $\nu$  has polar decomposition  $(\lambda, \nu_\xi)$ . Then, we have

$$\begin{aligned}\tilde{\nu}(B) &= \int_0^\infty s^{-\alpha-1} e^{-s^\beta} ds \int_S \lambda(d\xi) \int_0^\infty 1_{s^{-1}B}(r\xi) \nu_\xi(dr) \\ &= \int_S \lambda(d\xi) \int_0^\infty \nu_\xi(dr) r^{-1} \int_0^\infty (u/r)^{-\alpha-1} e^{-(u/r)^\beta} 1_B(u\xi) du \\ &= \int_S \lambda(d\xi) \int_0^\infty 1_B(u\xi) u^{-\alpha-1} \tilde{g}_\xi(u^\beta) du,\end{aligned}$$

where  $\tilde{\lambda} = \lambda$  and

$$(2.5) \quad \tilde{g}_\xi(u) = \int_0^\infty r^\alpha e^{-u/r^\beta} \nu_\xi(dr),$$

which is (2.4). The finiteness of  $\tilde{g}_\xi$  is trivial for  $\alpha \leq 0$ . For  $\alpha > 0$ , since  $\mu \in \mathfrak{D}(\Psi_{\alpha,\beta})$ , we have that  $\int_{\mathbb{R}^d} |x|^\alpha \mu(dx) < \infty$ . When  $\alpha > 0$ , note that  $\int_{\mathbb{R}^d} |x|^\alpha \mu(dx) < \infty$  implies  $\int_1^\infty r^\alpha \nu_\xi(dr) < \infty$ , (see, e.g. Sato (1999), Theorem 25.3). Hence the integral  $\tilde{g}_\xi$  exists.

(3) If we put

$$\tilde{Q}(B) = \int_0^\infty r^\alpha 1_B(r^{-\beta}) \nu_\xi(dr),$$

then it follows that  $\tilde{g}_\xi(u) = \int_0^\infty e^{-uy} \tilde{Q}(dy)$ , and thus  $\tilde{g}_\xi$  is completely monotone by Bernstein's theorem. If  $\alpha \leq 0$ , then automatically  $\lim_{u \rightarrow \infty} \tilde{g}_\xi(u) = 0$   $\lambda$ -a.e.  $\xi$ , since

$$\infty > \int_{|x|>1} \tilde{\nu}(dx) = \int_S \lambda(d\xi) \int_1^\infty u^{-\alpha-1} \tilde{g}_\xi(u^\beta) du.$$

When  $\alpha > 0$ , since  $\int_1^\infty r^\alpha \nu_\xi(dr) < \infty$ , the assertion that  $\lim_{u \rightarrow \infty} \tilde{g}_\xi(u) = 0$   $\lambda$ -a.e.  $\xi$  also follows from (2.5) by the dominated convergence theorem.

The proof of the lemma is thus concluded.  $\square$

**Remark 2.6.** (2.3) can be written as, by introducing a transformation  $\Upsilon_{\alpha,\beta}$  of Lévy measures as  $\tilde{\nu} = \Upsilon_{\alpha,\beta}(\nu)$ . Then this  $\Upsilon_{\alpha,\beta}$  is a generalized Upsilon transformation discussed in Barndorff-Nielsen et al. (2008) with the dilation measure  $\tau(ds) = s^{-\alpha-1} e^{-s^\beta} ds$ .

**Theorem 2.7.** For each  $\alpha < 2$  and  $\beta > 0$ , the mapping  $\Psi_{\alpha,\beta}$  is one-to-one.

The proof is carried out in the same way as for Proposition 4.1 of Sato (2006b).

We are now ready to discuss stochastic integral characterizations of the classes  $J_{\alpha,\beta}(\mathbb{R}^d)$ , by showing that  $J_{\alpha,\beta}(\mathbb{R}^d)$  is the range of the mapping  $\Psi_{\alpha,\beta}$ . However, in this paper, we restrict ourselves to the case  $\alpha < 1$ , because in the case  $1 \leq \alpha < 2$ ,  $J_{\alpha,\beta}(\mathbb{R}^d)$  is strictly bigger than the range  $\Psi_{\alpha,\beta}(\mathfrak{D}(\Psi_{\alpha,\beta}))$  and more deep calculations

would be needed. (See, e.g., Sato (2006b) and Maejima et al. (2009).) Also, the classes appearing in our motivation of introducing the classes  $J_{\alpha,\beta}(\mathbb{R}^d)$  are restricted to the case  $\alpha \leq 0$ .

**Theorem 2.8.** *Let  $\alpha < 1$  and  $\beta > 0$ . The range of the mapping  $\Psi_{\alpha,\beta}$  equals  $J_{\alpha,\beta}(\mathbb{R}^d)$ , that is,*

$$J_{\alpha,\beta}(\mathbb{R}^d) = \Psi_{\alpha,\beta}(\mathfrak{D}(\Psi_{\alpha,\beta})).$$

**Remark 2.9.** This theorem is already known for  $\alpha = -1, 0$  and  $\beta = 1$  in Theorems A and C of Barndorff-Nielsen et al. (2006) and for  $\alpha < 1$  and  $\beta = 1$  in Theorem 4.2 of Sato (2006b).

*Proof of Theorem 2.8.* We first show that  $\Psi_{\alpha,\beta}(\mathfrak{D}(\Psi_{\alpha,\beta})) \subset J_{\alpha,\beta}(\mathbb{R}^d)$ . Suppose  $\mu \in \mathfrak{D}(\Psi_{\alpha,\beta})$  and  $\tilde{\mu} = \Psi_{\alpha,\beta}(\mu)$ , and let  $\nu$  and  $\tilde{\nu}$  be the Lévy measures of  $\mu$  and  $\tilde{\mu}$ , respectively. Thus, if  $\nu = 0$ , then  $\tilde{\nu} = 0$  and  $\tilde{\mu} \in J_{\alpha,\beta}(\mathbb{R}^d)$ . When  $\nu \neq 0$ , it follows from Lemma 2.5 that  $\tilde{\mu} \in J_{\alpha,\beta}(\mathbb{R}^d)$ .

Next we show that  $J_{\alpha,\beta}(\mathbb{R}^d) \subset \Psi_{\alpha,\beta}(\mathfrak{D}(\Psi_{\alpha,\beta}))$ . Suppose  $\tilde{\mu} \in J_{\alpha,\beta}(\mathbb{R}^d)$  with the Lévy-Khintchine triplet  $(\tilde{A}, \tilde{\nu}, \tilde{\gamma})$ . If  $\tilde{\nu} = 0$ , then  $\tilde{\mu} = \Psi_{\alpha,\beta}(\mu)$  for some  $\mu \in \mathfrak{D}(\Psi_{\alpha,\beta})$ . Thus, suppose that  $\tilde{\nu} \neq 0$ . Then, in a polar decomposition  $(\tilde{\lambda}, \tilde{\nu}_\xi)$  of  $\tilde{\nu}$ , we have  $\tilde{\nu}_\xi(dr) = r^{-\alpha-1} \tilde{g}_\xi(r^\beta) dr$ , where  $\tilde{g}_\xi(v)$  is completely monotone in  $v > 0$   $\tilde{\lambda}$ -a.e.  $\xi$ , and is measurable in  $\xi$ . Thus by Bernstein's theorem, there are measures  $\tilde{Q}_\xi$  on  $[0, \infty)$  such that

$$\tilde{g}_\xi(v) = \int_{[0,\infty)} e^{-vu} \tilde{Q}_\xi(du).$$

In general,  $\tilde{Q}_\xi$  is a measure on  $[0, \infty)$ , but since  $\lim_{v \rightarrow \infty} \tilde{g}_\xi(v) = 0$   $\tilde{\lambda}$ -a.e.  $\xi$ ,  $\tilde{Q}_\xi$  does not have a point mass at 0, and hence  $\tilde{Q}_\xi$  is a measure on  $(0, \infty)$ . We see that

$$\begin{aligned} (2.6) \quad \tilde{\nu}(B) &= \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_B(r\xi) r^{-\alpha-1} \tilde{g}_\xi(r^\beta) dr \\ &= \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_B(r\xi) r^{-\alpha-1} dr \int_0^\infty e^{-r^\beta u} \tilde{Q}_\xi(du). \end{aligned}$$

Since  $\int_{\mathbb{R}^d} (|x|^2 \wedge 1) \tilde{\nu}(dx) < \infty$ , we have

$$\int_S \tilde{\lambda}(d\xi) \int_0^1 r^{1-\alpha} dr \int_1^\infty e^{-r^\beta u} \tilde{Q}_\xi(du) + \int_S \tilde{\lambda}(d\xi) \int_1^\infty r^{-\alpha-1} dr \int_0^1 e^{-r^\beta u} \tilde{Q}_\xi(du) < \infty.$$

Hence, we have, by the change of variables  $r \rightarrow v$  by  $r^\beta u = v$ ,

$$\int_0^1 r^{1-\alpha} dr \int_1^\infty e^{-r^\beta u} \tilde{Q}_\xi(du) = \int_1^\infty \tilde{Q}_\xi(du) \int_0^1 r^{1-\alpha} e^{-r^\beta u} dr$$

$$= \beta^{-1} \int_1^\infty u^{(\alpha-2)/\beta} \tilde{Q}_\xi(du) \int_0^u v^{-1+(2-\alpha)/\beta} e^{-v} dv \geq C_4 \int_1^\infty u^{(\alpha-2)/\beta} \tilde{Q}_\xi(du),$$

where

$$C_4 = \beta^{-1} \int_0^1 v^{-1+(2-\alpha)/\beta} e^{-v} dv \in (0, \infty).$$

Thus

$$(2.7) \quad \int_S \tilde{\lambda}(d\xi) \int_1^\infty u^{(\alpha-2)/\beta} \tilde{Q}_\xi(du) < \infty.$$

We also have for any  $\alpha < 1$ ,

$$(2.8) \quad \begin{aligned} \int_1^\infty r^{-\alpha-1} dr \int_0^1 e^{-r^\beta u} \tilde{Q}_\xi(du) &= \int_0^1 \tilde{Q}_\xi(du) \int_1^\infty r^{-\alpha-1} e^{-r^\beta u} dr \\ &= \beta^{-1} \int_0^1 u^{\alpha/\beta} \tilde{Q}_\xi(du) \int_u^\infty v^{-1-(\alpha/\beta)} e^{-v} dv \geq C_5 \int_0^1 u^{\alpha/\beta} \tilde{Q}_\xi(du), \end{aligned}$$

where

$$C_5 = \beta^{-1} \int_1^\infty v^{-1-(\alpha/\beta)} e^{-v} dv \in (0, \infty).$$

Thus

$$(2.9) \quad \int_S \tilde{\lambda}(d\xi) \int_0^1 u^{\alpha/\beta} \tilde{Q}_\xi(du) < \infty.$$

In addition, if  $\alpha = 0$ , (2.8) is turned out to be

$$\begin{aligned} \int_1^\infty r^{-1} dr \int_0^1 e^{-r^\beta u} \tilde{Q}_\xi(du) &= \beta^{-1} \int_0^1 \tilde{Q}_\xi(du) \int_u^\infty v^{-1} e^{-v} dv \\ &\geq (\beta e)^{-1} \int_0^1 \tilde{Q}_\xi(du) \int_u^1 v^{-1} dv = (\beta e)^{-1} \int_0^1 (-\log u) \tilde{Q}_\xi(du). \end{aligned}$$

Thus, when  $\alpha = 0$ ,

$$(2.10) \quad \int_S \tilde{\lambda}(d\xi) \int_0^1 (-\log u) \tilde{Q}_\xi(du) < \infty.$$

Furthermore,

$$\int_1^\infty r^{-\alpha-1} dr \int_0^1 e^{-r^\beta u} \tilde{Q}_\xi(du) \geq \int_1^\infty r^{-\alpha-1} e^{-r^\beta} dr \int_0^1 \tilde{Q}_\xi(du) = C_6 \int_0^1 \tilde{Q}_\xi(du),$$

where

$$C_6 := \int_1^\infty r^{-\alpha-1} e^{-r^\beta} dr \in (0, \infty).$$

Thus we have

$$(2.11) \quad \int_S \tilde{\lambda}(d\xi) \int_0^1 \tilde{Q}_\xi(dr) < \infty.$$

Define

$$(2.12) \quad \nu_\xi(B) = \int_0^\infty u^{\alpha/\beta} 1_B(u^{-1/\beta}) \tilde{Q}_\xi(du), \quad B \in \mathcal{B}((0, \infty)).$$

Then, it follows from (2.7) and (2.9) that

$$(2.13) \quad \begin{aligned} \int_S \tilde{\lambda}(d\xi) \int_0^\infty (r^2 \wedge 1) \nu_\xi(dr) &= \int_S \tilde{\lambda}(d\xi) \int_0^\infty u^{\alpha/\beta} (u^{-2/\beta} \wedge 1) \tilde{Q}_\xi(du) \\ &= \int_S \tilde{\lambda}(d\xi) \left( \int_0^1 u^{\alpha/\beta} \tilde{Q}_\xi(du) + \int_1^\infty u^{(\alpha-2)/\beta} \tilde{Q}_\xi(du) \right) < \infty. \end{aligned}$$

Define  $\nu$  by

$$(2.14) \quad \nu(B) = \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_B(r\xi) \nu_\xi(dr).$$

Then, by (2.13),  $\nu$  is the Lévy measure of some infinitely divisible distribution  $\mu$ , and  $\mu$  belongs to  $\mathfrak{D}(\Psi_{\alpha,\beta})$  and satisfies

$$(2.15) \quad \tilde{\nu}(B) = \int_0^{G_{\alpha,\beta}(0)} \nu((G_{\alpha,\beta}^*(t))^{-1}B) dt.$$

To show (2.15), by (2.6), (2.12) and (2.14), we have

$$\begin{aligned} \tilde{\nu}(B) &= \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_B(r\xi) r^{-\alpha-1} dr \int_0^\infty e^{-r^\beta u} \tilde{Q}_\xi(du) \\ &= \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_B(u^{-1/\beta} s\xi) s^{-\alpha-1} e^{-s^\beta} ds \int_0^\infty u^{\alpha/\beta} \tilde{Q}_\xi(du) \\ &= \int_0^\infty s^{-\alpha-1} e^{-s^\beta} ds \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_B(u^{-1/\beta} s\xi) u^{\alpha/\beta} \tilde{Q}_\xi(du) \\ &= \int_0^\infty s^{-\alpha-1} e^{-s^\beta} ds \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_B(rs\xi) \nu_\xi(dr) \\ &= \int_0^\infty s^{-\alpha-1} e^{-s^\beta} ds \int_S \tilde{\lambda}(d\xi) \int_0^\infty 1_{s^{-1}B}(r\xi) \nu_\xi(dr) \\ &= \int_0^\infty \nu(s^{-1}B) s^{-\alpha-1} e^{-s^\beta} ds = - \int_0^\infty \nu(s^{-1}B) dG_{\alpha,\beta}(s) \\ &= \int_0^{G_{\alpha,\beta}(0)} \nu((G_{\alpha,\beta}^*(t))^{-1}B) dt. \end{aligned}$$

To show that  $\mu \in \mathfrak{D}(\Psi_{\alpha,\beta})$ , it is enough to show that  $\int_{|x|>1} |x|^\alpha \nu(dx) < \infty$ , which is if and only if  $\mu \in I_\alpha(\mathbb{R}^d)$ , when  $0 < \alpha < 1$ , and  $\int_{|x|>1} \log |x| \nu(dx) < \infty$ , which is if and only if  $\mu \in I_{\log}(\mathbb{R}^d)$ , when  $\alpha = 0$ , (see Sato (1999), Theorem 25.3). Note that by (2.12) we see, for any nonnegative measurable function  $f$  on  $(0, \infty)$ ,

$$\int_0^\infty f(r) \nu_\xi(dr) = \int_0^\infty u^{\alpha/\beta} f(u^{-1/\beta}) \tilde{Q}_\xi(du).$$

Thus if we choose  $f(r) = I[r > 1]r^\alpha$ , where  $I[A]$  is the indicator function of the set  $A$ , then  $\nu$  in (2.14) satisfies that for  $\alpha > 0$

$$(2.16) \quad \int_{|x|>1} |x|^\alpha \nu(dx) = \int_S \tilde{\lambda}(d\xi) \int_1^\infty r^\alpha \nu_\xi(dr) = \int_S \tilde{\lambda}(d\xi) \int_0^1 \tilde{Q}_\xi(du) < \infty$$

due to (2.11). When  $\alpha = 0$ ,

$$(2.17) \quad \begin{aligned} \int_{|x|>1} \log |x| \nu(dx) &= \int_S \tilde{\lambda}(d\xi) \int_1^\infty \log r \nu_\xi(dr) \\ &= \int_S \tilde{\lambda}(d\xi) \int_0^1 \log u^{-1/\beta} \tilde{Q}_\xi(du) = \beta^{-1} \int_S \tilde{\lambda}(d\xi) \int_0^1 (-\log u) \tilde{Q}_\xi(du) < \infty \end{aligned}$$

due to (2.10).

Notice again that

$$\int_0^{G_{\alpha,\beta}(0)} (G_{\alpha,\beta}^*(t))^2 dt = - \int_0^\infty u^2 dG_{\alpha,\beta}(u) = \int_0^\infty u^{1-\alpha} e^{-u^\beta} du < \infty.$$

Define  $A$  and  $\gamma$  by

$$(2.18) \quad \tilde{A} = \left( \int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t)^2 dt \right) A$$

and

$$(2.19) \quad \tilde{\gamma} = \int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dt \left( \gamma + \int_{\mathbb{R}^d} x \left( \frac{1}{1 + |G_{\alpha,\beta}^*(t)x|^2} - \frac{1}{1 + |x|^2} \right) \nu(dx) \right).$$

Here we have to check the finiteness of this integral. We first have

$$\int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dt = - \int_0^\infty u dG_{\alpha,\beta}(u) = \int_0^\infty u^{-\alpha} e^{-u^\beta} du < \infty,$$

since  $\alpha < 1$ . Below,  $C_7, C_8 \in (0, \infty)$  are suitable constants. Recall  $\alpha < 1$ . When  $\alpha \neq 0$ , we have

$$\begin{aligned} & \int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dt \int_{\mathbb{R}^d} |x| \left| \frac{1}{1 + |G_{\alpha,\beta}^*(t)x|^2} - \frac{1}{1 + |x|^2} \right| \nu(dx) \\ &= \int_0^\infty u^{-\alpha} e^{-u^\beta} du \int_{\mathbb{R}^d} |x| \left| \frac{1}{1 + |ux|^2} - \frac{1}{1 + |x|^2} \right| \nu(dx) \\ &\leq \int_0^\infty u^{-\alpha} (1 + u^2) e^{-u^\beta} du \int_{\mathbb{R}^d} \frac{|x|^3}{(1 + |ux|^2)(1 + |x|^2)} \nu(dx) \\ &\leq \int_0^\infty u^{-\alpha} (1 + u^2) e^{-u^\beta} du \\ &\quad \times \left( \int_{|x|\leq 1} |x|^2 \nu(dx) + \int_{|x|>1, |ux|\leq 1} |x| \nu(dx) + \int_{|x|>1, |ux|>1} \frac{|x|}{|ux|^2} \nu(dx) \right) \end{aligned}$$

$$\begin{aligned}
&= C_7 + \int_{|x|>1} |x| \nu(dx) \int_0^{1/|x|} u^{-\alpha} (1+u^2) e^{-u^\beta} du \\
&\quad + \int_{|x|>1} \nu(dx) \int_{1/|x|}^\infty u^{-\alpha-1} (1+u^2) e^{-u^\beta} du \\
&\leq C_7 + \int_{|x|>1} |x| \nu(dx) \int_0^{1/|x|} 2u^{-\alpha} du \\
&\quad + \int_{|x|>1} \nu(dx) \left\{ \left( \int_{1/|x|}^1 + \int_1^\infty \right) u^{-\alpha-1} (1+u^2) e^{-u^\beta} du \right\} \\
(2.20) \quad &\leq C_7 + 2(1-\alpha)^{-1} \int_{|x|>1} |x|^\alpha \nu(dx) \\
&\quad + \int_{|x|>1} \nu(dx) \left\{ \int_{1/|x|}^1 2u^{-\alpha-1} du + \int_1^\infty u^{-\alpha-1} (1+u^2) e^{-u^\beta} du \right\} \\
&= C_7 + 2(1-\alpha)^{-1} \int_{|x|>1} |x|^\alpha \nu(dx) \\
&\quad + \int_{|x|>1} \nu(dx) \{ -2\alpha^{-1} (1-|x|^\alpha) + C_8 \} \\
(2.21) \quad &= C_7 + 2(1-\alpha)^{-1} \int_{|x|>1} |x|^\alpha \nu(dx) \\
&\quad + 2\alpha^{-1} \int_{|x|>1} |x|^\alpha \nu(dx) + (C_8 - 2\alpha^{-1}) \int_{|x|>1} \nu(dx) < \infty,
\end{aligned}$$

by (2.16). When  $\alpha = 0$ , since

$$\int_{1/|x|}^1 u^{-\alpha-1} du = \int_{1/|x|}^1 u^{-1} du = \log |x|,$$

in (2.20), we have

$$(2.22) \quad \int_{|x|>1} \log |x| \nu(dx)$$

instead of  $\int_{|x|>1} |x|^\alpha \nu(dx)$  in (2.21) in the calculation above. The finiteness of (2.22) is assured by (2.17).

Thus  $\gamma$  can be defined. Hence, if we denote by  $\mu$  an infinitely divisible distribution having the Lévy-Khintchine triplet  $(A, \nu, \gamma)$  above, then by (2.15), (2.18) and (2.19), we see that

$$\tilde{\mu} = \mathcal{L} \left( \int_0^{G_{\alpha,\beta}(0)} G_{\alpha,\beta}^*(t) dX_t^{(\mu)} \right),$$

concluding that  $\tilde{\mu} \in \Psi_{\alpha,\beta}(\mathfrak{D}(\Psi_{\alpha,\beta}))$ . This completes the proof.  $\square$

**Acknowledgment** The authors greatly appreciate the referee's detailed comments, which improved the paper very much.

#### REFERENCES

- [1] T. Aoyama, M. Maejima and J. Rosiński (2008). A subclass of type  $G$  selfdecomposable distributions, *J. Theoret. Probab.*, **21**, 14–34.
- [2] O.E. Barndorff-Nielsen, M. Maejima and K. Sato (2006). Some classes of multivariate infinitely divisible distributions admitting stochastic integral representations, *Bernoulli*, **12**, 1–33.
- [3] Barndorff-Nielsen, O.E., Rosiński, J. and Thorbjørnsen, S. (2008): General  $\Upsilon$  transformations. *ALEA Lat. Am. J. Probab. Math. Statist.* **4**, 131-165.
- [4] W. Feller (1966). *An Introduction to Probability Theory and Its Applications, Vol. II, 2nd ed.*, John Wiley & Sons..
- [5] M. Maejima, M. Matsui and M. Suzuki (2009). Classes of infinitely divisible distributions on  $\mathbb{R}^d$  related to the class of selfdecomposable distributions, to appear in *Tokyo J. Math.*
- [6] K. Sato (1999). *Lévy Processes and Infinitely Divisible Distributions*. Cambridge University Press.
- [7] Sato, K. (2004) Stochastic integrals in additive processes and application to semi-Lévy processes, *Osaka J. Math.* **41**, 211–236.
- [8] K. Sato (2006a). Additive processes and stochastic integrals, *Illinois J. Math.* **50**, 825–851.
- [9] M. Sato (2006b). Two families of improper stochastic integrals with respect to Lévy processes, *ALEA Lat. Am. J. Probab. Math. Statist.* **1** 47–87.