

Research Report

KSTS/RR-04/001

Jan. 13, 2004

**Poisson point processes attached
to symmetric diffusions**

by

**Masatoshi Fukushima
Hiroshi Tanaka**

<p>Masatoshi Fukushima Kansai University Hiroshi Tanaka</p>

Department of Mathematics
Faculty of Science and Technology
Keio University

©2004 KSTS

3-14-1 Hiyoshi, Kohoku-ku, Yokohama, 223-8522 Japan

Poisson point processes attached to symmetric diffusions

by

Masatoshi Fukushima

Department of Mathematics, Faculty of Engineering, Kansai University,
Suita 564-8680, Japan
e-mail: fuku@ipcku.kansai-u.ac.jp

and

Hiroshi Tanaka

1-4-17-104 Miyamaedaira, Miyamae-ku, Kawasaki 216-0006 Japan

ABSTRACT. -Let a be a non-isolated point of a topological space S and $X^0 = (X_t^0, 0 \leq t < \zeta^0, P_x^0)$ be a symmetric diffusion on $S_0 = S \setminus \{a\}$ such that $P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 = a) > 0, x \in S_0$. By making use of Poisson point processes taking values in the spaces of excursions around a whose characteristic measures are uniquely determined by X^0 , we construct a symmetric diffusion \tilde{X} on S with no killing inside S which extends X^0 on S_0 . We also prove that such a process \tilde{X} is unique in law and its resolvent and Dirichlet form admit explicit expressions in terms of X^0 .

Keywords: symmetric diffusion, Poisson point process, excursions, entrance law, energy functional, Dirichlet form

1 Introduction

Let S be a locally compact separable metric space and a be a non-isolated point of S . We put $S_0 = S \setminus \{a\}$. The one point compactification of S is denoted by S_Δ . When S is compact already, Δ is added as an isolated point. Let m be a positive Radon measure on S_0 with $\text{Supp}[m] = S_0$. m is extended to S by setting $m(\{a\}) = 0$.

We assume that we are given an m -symmetric diffusion $X^0 = (X_t^0, P_x^0)$ on S_0 with life time ζ^0 satisfying the following four conditions:

A.1 $P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 \in \{a\} \cup \{\Delta\}) = P_x^0(\zeta^0 < \infty), \quad \forall x \in S_0.$

We define the functions $\varphi(x)$, $u_\alpha(x)$, $\alpha > 0$, of $x \in S_0$ by

$$\varphi(x) = P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 = a), \quad u_\alpha(x) = E_x^0(e^{-\alpha\zeta^0}; X_{\zeta^0-}^0 = a).$$

A.2 $\varphi(x) > 0, \quad \forall x \in S_0,$

A.3 $u_\alpha \in L^1(S_0; m), \quad \forall \alpha > 0.$

A.4 $u_\alpha \in C_b(S_0), \quad G_\alpha^0(C_b(S_0)) \subset C_b(S_0), \quad \alpha > 0,$

where G_α^0 is the resolvent of X^0 and $C_b(S_0)$ is the space of all bounded continuous functions on S_0 .

By making use of excursion-valued Poisson point processes whose characteristic measures are uniquely determined by X^0 , or to be a little more precise, by piecing together

those excursions which start from a and return to a and then possibly by adding the last one that never returns to a , we shall construct in §4 of the present paper a process \tilde{X} on S satisfying

- (1) \tilde{X} is an m -symmetric diffusion process on S with no killing inside S ,
- (2) \tilde{X} is an extension of X^0 : the process on S_0 obtained from \tilde{X} by killing upon the hitting time of a is identical in law with X^0 .

We call a process \tilde{X} on S satisfying (1),(2) a *symmetric extension of X^0* .

We shall also prove in §5 that, under conditions **A.1**, **A.2** for the given m -symmetric diffusion X^0 on S_0 , its symmetric extension is unique in law, satisfies condition **A.3** automatically and admits the resolvent expressible as

$$G_\alpha f(x) = G_\alpha^0 f(x) + u_\alpha(x) \cdot G_\alpha f(a), \quad x \in S_0, \quad G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)},$$

where (\cdot, \cdot) denotes the inner product in $L^2(S_0; m)$ and $L(m_0, \psi)$ is the energy functional in Meyer's sense [21] of the X^0 -excessive measure $m_0 = \varphi \cdot m$ and X^0 -excessive function $\psi = 1 - \varphi$.

Furthermore the associated Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(S; m)$ will be seen in §5 to have the following simple expression; if we denote by \mathcal{F}_e its extended Dirichlet space, then

$$\mathcal{F}_e = \{w = u_0 + c\varphi : u_0 \in \mathcal{F}_{0,e}, \quad c \text{ constant}\}, \quad \mathcal{F} = \mathcal{F}_e \cap L^2(S; m),$$

$$\mathcal{E}(w, w) = \mathcal{E}(u_0, u_0) + c^2 \mathcal{E}(\varphi, \varphi), \quad \mathcal{E}(\varphi, \varphi) = L(m_0, \psi),$$

where $(\mathcal{F}_{0,e}, \mathcal{E})$ is the extended Dirichlet space for the given diffusion X^0 .

In §6, we shall present four examples. Example 6.1 concerns the uniqueness of the symmetric extension of the one dimensional absorbing Brownian motion.

Example 6.2 treats the case where S_0 is a bounded open subset of \mathbb{R}^d , ($d \geq 1$), $S = S_0 \cup \{a\}$ is the one point compactification of S_0 and X^0 is the absorbing Brownian motion on S_0 . In this case, $\varphi(x) = 1$, $x \in S_0$. The resulting Dirichlet form on $L^2(S; m)$ (m is the Lebesgue measure on S_0 extended to S by $m(\{a\}) = 0$) is given by

$$\mathcal{F} = \{w = u_0 + c : u_0 \in H_0^1(S_0), \quad c \text{ constant}\},$$

$$\mathcal{E}(w, w) = \frac{1}{2} \int_{S_0} |\nabla u_0|^2(x) dx,$$

which is easily seen to be regular, strongly local and irreducible recurrent. A more general Dirichlet form of this type will be presented in §3.2. This type of Dirichlet form first appeared in the paper [8] by the first author and it is recently utilized in a study of the asymptotics of the spectral gap for one parameter family of energy forms([17]). Our study is motivated by a wish to conceive a clearer picture of the sample path of the diffusion on S associated with such a Dirichlet form.

Example 6.3 is essentially one-dimensional, where we shall see that the conditions **A.2** and **A.3** are satisfied if and only if the boundary is regular in Feller's sense. This example is reminiscent of an example by N. Ikeda and S. Watanabe[14].

Example 6.4 is higher dimensional, where the Dirichlet form associated with the constructed process \tilde{X} may not be regular.

In order to identify right quantities to describe the excursion-valued Poisson point processes to be constructed in §4, we shall study in §2 and §3 a strongly local regular Dirichlet form on $L^2(S; m)$ for which the point $\{a\}$ has a positive capacity. In particular, we shall find that the Dirichlet form and the associated resolvent admit exactly the above mentioned expressions. Furthermore, we shall see that the entrance law $\{\mu_t\}$ governing the excursion law ought to be determined by

$$m_0 = \int_0^\infty \mu_t dt,$$

an equation investigated by E.B.Dynkin, R.K.Gettoor, P.J.Fitzsimmons and others ([11]).

In a seminal work [15], K.Itô considered a standard process X on S for which a point a is regular for itself. A Poisson point process \mathbf{Y} taking value in the space of excursions around a was then associated, and it was shown that the stopped process X^0 obtained from X by the hitting time at a and the characteristic measure of \mathbf{Y} together determine the law of X uniquely. It was implicitly assumed in [15] that the point a is recurrent in the sense that

$$\varphi(x) = P_x(\sigma_a < \infty) = 1, \quad x \in S, \quad \sigma_a = \inf\{t > 0 : X_t = a\}.$$

But, as was shown in P.A. Meyer [20], an *absorbed* Poisson point process can be still associated with X when $\{a\}$ is non-recurrent. See Remark 4.2 in this regard.

Since our present assumption on X^0 requires φ only to be positive, we must handle not only returning excursions from the point a but also non-returning excursions. By restricting ourselves to the case that both X^0 and \tilde{X} are symmetric diffusions however, we shall see that the characteristic measures on these different type of excursion spaces are uniquely determined by X^0 so that, starting with X^0 , we can give an explicit construction of \tilde{X} .

The Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(S; m)$ associated with a symmetric extension \tilde{X} of X^0 may not be regular but it is quasi-regular in the sense of [19]. Accordingly we can make use of the quasi-homeomorphism in [3] to connect \tilde{X} with the regular Dirichlet form studied in §2, yielding the uniqueness of \tilde{X} and the explicit expression of $(\mathcal{E}, \mathcal{F})$.

There are quite a few works [1], [24], [25], [26] dealing with generalizations of Itô's one [15]. See Remark 2.2 and Remark 4.1 in these regards. But construction and uniqueness of a symmetric extension X of a symmetric X^0 as are formulated in the present paper have never been considered.

2 Strongly local Dirichlet form with a point of positive capacity

2.1 Description of the form and resolvent by absorbed process

Let S be a locally compact separable metric space and a be a non-isolated point of S . We denote the complementary set $S \setminus \{a\}$ by S_0 . Let m be a positive Radon measure on S with $\text{Supp}[m] = S$ and with $m(\{a\}) = 0$. The inner product in each of the spaces $L^2(S; m)$, $L^2(S_0, m)$ will be designated by (\cdot, \cdot) .

A Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(S; m)$ is called *regular* if $\mathcal{F} \cap C_0(S)$ is \mathcal{E}_1 -dense in \mathcal{F} and uniformly dense in $C_0(S)$, where $C_0(S)$ denotes the space of continuous functions on

S with compact support. It is called *strongly local* if $\mathcal{E}(u, v)$ vanishes whenever $u, v \in \mathcal{F}$, $\text{Supp}[u]$, $\text{Supp}[v]$ are compact and v is constant on a neighbourhood of $\text{Supp}[u]$, where $\text{Supp}[u]$ denotes the topological support of the measure $u \cdot m$. For the sake of a use in §3.2, we make here a remark:

Remark 2.1. If a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(S; m)$ is regular and strongly local, then the strong locality stated above holds without assuming that $\text{Supp}[v]$ is compact. Indeed, assuming the boundedness of v , take a function $w \in \mathcal{F} \cap C_0(S)$ with $w = 1$ on a neighbourhood of $K = \text{Supp}[u]$ and put $v_1 = v \cdot w$, $v_0 = v - v_1$. Then $\mathcal{E}(u, v_1) = 0$. Since v_0 belongs to the part \mathcal{F}_G of $(\mathcal{E}, \mathcal{F})$ on the open set $G = S \setminus K$ and $(\mathcal{E}, \mathcal{F}_G)$ is a regular Dirichlet form on $L^2(G; m)$ (cf. [9, Th.4.4.3]), we can find $v_n \in \mathcal{F} \cap C_0(G)$ which are \mathcal{E}_1 -convergent to v_0 . Hence $\mathcal{E}(u, v_0) = \lim_{n \rightarrow \infty} \mathcal{E}(u, v_n) = 0$ and $\mathcal{E}(u, v) = 0$.

We consider a strongly local regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(S; m)$ and an associated m -symmetric Hunt process $X = (X_t, P_x)$ on S . In view of [9, The.4.5.3], X can then be taken to be a diffusion on S_Δ in the sense that all sample paths are continuous functions from $[0, \infty)$ to S_Δ , where S_Δ is the one-point compactification of S when S is non-compact and Δ is an extra point isolated from S when S is compact. In either case Δ will be the cemetery of the sample paths. Furthermore, X can be taken to be of no killing inside S in the sense that

$$P_x(X_{\zeta-} = \Delta, \zeta < \infty) = P_x(\zeta < \infty), \quad x \in S,$$

where $\zeta(\omega)$ denotes the life time, namely, the hitting time of the cemetery Δ of the sample path ω . In particular, when S is compact, $P_x(\zeta = \infty) = 1$ for all $x \in S$.

We make the assumption that

B.1 $\text{Cap}(\{a\}) > 0$.

Here $\text{Cap}(A)$ for $A \subset S$ is its 1-capacity relative to $(\mathcal{E}, \mathcal{F})$. In what follows, the quasi-continuity of functions on S will be understood with respect to this capacity. Each function $u \in \mathcal{F}$ admits its quasi-continuous version denoted by \tilde{u} . ‘q.e.’ will mean ‘except for a set of zero capacity’.

The hitting probability and the α -order hitting probability of $\{a\}$ are denoted by φ and u_α respectively:

$$\varphi(x) = P_x(\sigma < \infty), \quad u_\alpha(x) = E_x(e^{-\alpha\sigma}), \quad x \in S, \quad (2.1)$$

where σ is the hitting time of a by the process X defined by

$$\sigma = \inf\{t > 0 : X_t = a\}. \quad (2.2)$$

The assumption **B.1** implies that u_α is a non-trivial element of \mathcal{F} and it is the α -potential $U_\alpha \nu_\alpha$ of a positive measure ν_α concentrated on $\{a\}$ (cf. [9, §2.2]):

$$\mathcal{E}_\alpha(u_\alpha, v) = \tilde{v}(a) \nu_\alpha(\{a\}) \quad v \in \mathcal{F}. \quad (2.3)$$

Put

$$\mathcal{F}_0 = \{u \in \mathcal{F} : \tilde{u}(a) = 0\}. \quad (2.4)$$

Then $(\mathcal{E}, \mathcal{F}_0)$ is a regular strongly local Dirichlet form on $L^2(S_0; m)$, which is associated with the part $X^0 = (X_t^0, P_x^0)$ of X on the set S_0 , namely, the diffusion process X^0

obtained from X by killing upon the hitting time σ (cf. [9, §4.4]). X^0 is of no killing inside S_0 and, if we denote the life time of X^0 by ζ^0 , then φ , u_α admit the expressions

$$\varphi(x) = P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 = a), \quad u_\alpha(x) = E_x^0(e^{-\alpha\zeta^0}; X_{\zeta^0-}^0 = a), \quad x \in S_0, \quad (2.5)$$

in terms of the absorbed process X^0 . We further consider the functions

$$\psi^{(1)}(x) = P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 = \Delta), \quad \psi^{(2)}(x) = P_x^0(\zeta^0 = \infty), \quad x \in S_0, \quad (2.6)$$

and put $\psi = \psi^{(1)} + \psi^{(2)}$ so that $\psi = 1 - \varphi$.

Denote by p_t and G_α the transition function and the resolvent of X respectively. The same notions for the absorbed process X^0 will be denoted by p_t^0 and G_α^0 . The functions φ , $\psi^{(1)}$, $\psi^{(2)}$ on S_0 are X^0 -excessive. In particular, $\psi^{(2)}$ is X^0 -invariant in the sense that $\psi^{(2)} = p_t^0 \psi^{(2)}$, $t > 0$. Because of the m -symmetry of X^0 , the measure

$$m_0 = \varphi \cdot m \quad (2.7)$$

is an X^0 -excessive measure with $m_0 p_t^0 = p_t^0 \varphi \cdot m$.

Our first aim in this section is to show under the present setting that the form \mathcal{E} as well as the resolvent G_α are uniquely and explicitly determined by quantities depending only on the absorbed process X^0 .

We prepare a lemma.

Lemma 2.1. *For an X^0 -excessive function v on S_0 ,*

$$L(m_0, v) = \lim_{t \downarrow 0} \frac{1}{t} \langle m_0 - m_0 p_t^0, v \rangle = \lim_{t \downarrow 0} \frac{1}{t} (\varphi - p_t^0 \varphi, v) (\leq \infty). \quad (2.8)$$

is well defined as an increasing limit and it holds that

$$L(m_0, v) = \lim_{\alpha \rightarrow \infty} \alpha(u_\alpha, v). \quad (2.9)$$

If v is p_t^0 -invariant, then for each $t > 0$ and $\alpha > 0$,

$$L(m_0, v) = \frac{1}{t} (\varphi - p_t^0 \varphi, v) = \alpha(u_\alpha, v).$$

Proof. If we set $e(t) = (\varphi - p_t^0 \varphi, v)$, then

$$e(t+s) = e(t) + (p_t^0 \varphi - p_{t+s}^0 \varphi, v) = e(t) + (\varphi - p_s^0 \varphi, p_t^0 v) \leq e(t) + e(s),$$

and hence $e(t)/t$ is increasing as t decreases and constant if v is p_t^0 -invariant.. We also see that

$$\alpha(u_\alpha, v) = \alpha(\varphi - \alpha G_\alpha^0 \varphi, v) = \int_0^\infty e^{-t} (t/\alpha)^{-1} (\varphi - p_{t/\alpha}^0 \varphi, v) t dt$$

increases to $L(v)$ as $\alpha \uparrow \infty$. □

We note that $L(m_0, v)$ is nothing but the *energy functional* of the X^0 -excessive measure m_0 and the X^0 -excessive function v in the sense of P.A. Meyer [21] when X^0 is transient (cf. [4, §39], [11, p16]). In [4, §39], it is called the mass of v relative to m_0 .

Let \mathcal{F}_e (resp. $\mathcal{F}_{0,e}$) be the extended Dirichlet space of $(\mathcal{F}, \mathcal{E})$ (resp. $(\mathcal{F}_0, \mathcal{E})$). Each element $u \in \mathcal{F}_e$ admits its quasi continuous version denoted by \tilde{u} again. In view of [9, §4.6], it holds then that

$$\begin{aligned} \mathcal{F}_{0,e} &= \mathcal{F}_{e,0} = \{u \in \mathcal{F}_e : \tilde{u}(a) = 0\}, \\ \varphi \in \mathcal{F}_e, \quad \mathcal{E}(\varphi, u) &= 0 \quad \forall u \in \mathcal{F}_{e,0}, \end{aligned} \tag{2.10}$$

$$\mathcal{F} = \mathcal{F}_e \cap L^2(S; m) \quad \mathcal{F}_0 = \mathcal{F}_{0,e} \cap L^2(S_0, m). \tag{2.11}$$

Furthermore any $w \in \mathcal{F}_e$ can be decomposed as

$$w = u_0 + c \varphi, \quad u_0 \in \mathcal{F}_{e,0}, \quad c \text{ constant} \tag{2.12}$$

and

$$\mathcal{E}(w, w) = \mathcal{E}(u_0, u_0) + c^2 \mathcal{E}(\varphi, \varphi). \tag{2.13}$$

Theorem 2.1. (i) *It holds that*

$$\mathcal{E}(\varphi, \varphi) = L(m_0, \psi) (= L(m_0, \psi^{(1)}) + L(m_0, \psi^{(2)})). \tag{2.14}$$

(ii) u_α is a non-trivial element of $\mathcal{F} \cap L^1(S_0; m)$.

(iii) For any $f \in L^2(S, m)$ and $x \in S$,

$$G_\alpha f(x) = G_\alpha^0 f(x) + \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)} u_\alpha(x), \quad G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)}. \tag{2.15}$$

(iv) Let δ_a be a unit mass concentrated at $\{a\}$. Then it is of finite energy integral and its α -potential $U_\alpha \delta_a$ is related to u_α by

$$\widetilde{U_\alpha \delta_a} = \frac{1}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)} u_\alpha. \tag{2.16}$$

(v) The point a is regular for itself and also an instantaneous state with respect to X :

$$P_a(\sigma = 0, \tau_a = 0) = 1, \quad \tau_a = \inf\{t > 0 : X_t \in S_0\}. \tag{2.17}$$

Proof. We first give a proof of (ii). According to a general theorem ([9, Chap 4]), the formula obtained by the strong Markov property

$$G_\alpha f(x) = G_\alpha^0 f(x) + u_\alpha(x) G_\alpha f(a) \quad x \in S, \quad f \in L^2(S; m), \tag{2.18}$$

represents the orthogonal decomposition of $G_\alpha f \in \mathcal{F}$ into the space \mathcal{F}_0 and its orthogonal complement $\mathcal{H}_\alpha = \{c \cdot u_\alpha : c \text{ constant}\}$ in the Hilbert space $(\mathcal{F}, \mathcal{E}_\alpha)$. We see that $G_\alpha f(a) > 0$ for some $f \in C_0^+(S)$, because otherwise $\mathcal{F} = \mathcal{F}_0$ from (2.18) contradicting to $u_\alpha \in \mathcal{F}$. By (2.18),

$$(u_\alpha, 1) G_\alpha f(a) \leq (G_\alpha f, 1) = (f, G_\alpha 1) \leq \frac{1}{\alpha} (f, 1) < \infty.$$

Next we prove (i) and (iii). For $f \in C_0(S)$, the function $w = G_\alpha f$ has two expressions:

$$w = G_\alpha^0 f + c u_\alpha = u_0 + c \varphi, \quad c = G_\alpha f(a), \quad u_0 \in \mathcal{F}_{e,0}.$$

By [9, Cor.1.6.3, Th.2.1.7], We can find a sequence $\{g_n\}$ of uniformly bounded functions in \mathcal{F} such that

$$\lim_{n \rightarrow \infty} g_n = \varphi \text{ } m\text{-a.e.}, \quad \lim_{n \rightarrow \infty} \mathcal{E}(g_n - \varphi, g_n - \varphi) = 0.$$

Letting $n \rightarrow \infty$ in the equation

$$\mathcal{E}(w, g_n) + \alpha(w, g_n) = (f, g_n),$$

we get

$$c\mathcal{E}(\varphi, \varphi) + c\alpha(u_\alpha, \varphi) = (f, \varphi) - (\alpha G_\alpha^0 f, \varphi).$$

Since the right hand side equals

$$(f, \varphi - \alpha G_\alpha^0 \varphi) = (f, u_\alpha),$$

we arrive at

$$G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + \mathcal{E}(\varphi, \varphi)}, \quad f \in C_0(S). \quad (2.19)$$

(2.19) holds for any bounded Borel f . In particular, we have for any $\alpha > 0$,

$$G_\alpha 1(a) = \frac{(u_\alpha, 1)}{\alpha(u_\alpha, \varphi) + \mathcal{E}(\varphi, \varphi)} \leq \frac{1}{\alpha},$$

and hence

$$\mathcal{E}(\varphi, \varphi) \geq \alpha(u_\alpha, \psi).$$

By letting $\alpha \rightarrow \infty$, we get from Lemma 2.1

$$\mathcal{E}(\varphi, \varphi) \geq L(m_0, \psi).$$

In order to prove (2.14), notice that the assumption of the strong locality of \mathcal{E} implies that the killing measure k in the Beurling-Deny representation of \mathcal{E} vanishes (cf. [9, Th.4.5.3]). On account of [9, Lemma 4.5.2],

$$\int_S f^2 dk = \lim_{\alpha \rightarrow \infty} \alpha \int_S f(x)^2 (1 - \alpha G_\alpha 1(x)) m(dx), \quad f \in \mathcal{F} \cap C_0(S).$$

From (2.18) and (2.19), we have

$$\begin{aligned} 1 - \alpha G_\alpha 1(x) &= 1 - \alpha G_\alpha^0 1(x) - \frac{\alpha(u_\alpha, 1)}{\alpha(u_\alpha, \varphi) + \mathcal{E}(\varphi, \varphi)} u_\alpha(x) \\ &\geq u_\alpha(x) - \frac{\alpha(u_\alpha, 1)}{\alpha(u_\alpha, \varphi) + \mathcal{E}(\varphi, \varphi)} u_\alpha(x) \\ &= \frac{\mathcal{E}(\varphi, \varphi) - \alpha(u_\alpha, \psi)}{\alpha(u_\alpha, \varphi) + \mathcal{E}(\varphi, \varphi)} u_\alpha(x). \end{aligned}$$

Take $f \in \mathcal{F} \cap C_0(S)$ such that $f(a) \neq 0$. We have from (2.19) and the above inequality

$$\alpha \int_S f^2 (1 - \alpha G_\alpha 1) dm \geq (\mathcal{E}(\varphi, \varphi) - \alpha(u_\alpha, \psi)) (\alpha G_\alpha f^2)(a).$$

By letting $\alpha \rightarrow \infty$, we get

$$0 \geq (\mathcal{E}(\varphi, \varphi) - L(m_0, \psi))f(a)^2,$$

proving the desired identity (2.14).

Proof of (iv). By (2.3),

$$(u_\alpha, f) = \mathcal{E}_\alpha(u_\alpha, G_\alpha f) = G_\alpha f(a) \nu_\alpha(\{a\}),$$

which combined with (2.15) gives

$$\nu_\alpha = (\alpha(u_\alpha, \varphi) + L(m_0, \psi))\delta_a.$$

Proof of (v). The regularity $P_a(\sigma = 0) = 1$ of the point a for itself follows from **A.1** and a general fact that, for any Borel set B , the set of irregular points $x \in B$ for B is of zero capacity ([9, Chap. 4]). If $P_a(0 < \tau_a < \infty) > 0$, then $P_a(X_{\tau_a} \in S_0 \cup \Delta) = 1$ contradicting the sample continuity and absence of the killing inside S for X . If a were a trap with respect to X , then $G_\alpha f(a) = f(a)/\alpha$ for any $f \in L^2(S; m)$ contradicting (2.15). Accordingly, a is an instantaneous state. □

Remark 2.2. (i) The present assumptions can be relaxed as follows:

- (a) The measure m on S is replaced by $\bar{m} = m + \gamma\delta_a$ for a non-negative constant γ .
- (b) $(\mathcal{E}, \mathcal{F})$ is assumed to be a (not necessarily strongly) local regular Dirichlet form on $L^2(S; \bar{m})$, while its part $(\mathcal{E}, \mathcal{F}_0)$ on S_0 is assumed to be a strongly local Dirichlet form on $L^2(S_0; m)$.

Then, in view of the above proof of Theorem 2.1, we readily see that (2.14) and (2.15) remain true under the following modifications:

$$\mathcal{E}(\varphi, \varphi) = L(m_0, \psi) + \delta,$$

$$G_\alpha f(x) = G_\alpha^0 f(x) + \frac{(u_\alpha, f) + \gamma f(a)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi) + \delta + \alpha\gamma} u_\alpha(x),$$

for a non-negative constant δ .

Example 6.1 will indicate stochastic interpretations of the parameters γ and δ .

(ii) The parameters γ, δ have appeared in Rogers' description [24] of the most general extension of a general resolvent G_α^0 under a setting corresponding to $\psi^{(1)} = 0$. Another parameter appearing in [24] is a family of measures $n_\alpha, \alpha > 0$, on S_0 , which is reduced to $u_\alpha \cdot m$ under the present symmetry assumption.

(iii) In the setting (i) in the above, G_α is conservative if and only if $\psi^{(1)} = 0$ and $\delta = 0$, and in this case the above expression is reduced to

$$G_\alpha f(x) = G_\alpha^0 f(x) + \frac{(1 - \alpha G_\alpha^0 1, f) + \gamma f(a)}{\alpha(1 - \alpha G_\alpha^0 1, 1) + \alpha\gamma} (1 - \alpha G_\alpha^0 1(x)).$$

Such a formula was found by Y. Le Jan [18] (see also [4, §78]) in a general setting to produce conservative resolvents out of a (not necessarily symmetric) submarkovian resolvent and its dual preserving the duality.

2.2 Description of the inverse local time

In §4, we shall construct a diffusion on S with resolvent (2.15) by means of Poisson point processes of excursions, namely, by piecing together the excursions. In this subsection, let us study more about the roles of the measure m_0 and the energy functional $L(m_0, \psi)$ played in the present diffusion X on S .

Let $L(t)$ be the positive continuous additive functional (admitting exceptional set) associated with the smooth measure δ_a (cf.[9, §5.1]):

$$\widetilde{U}_\alpha \delta_a(x) = E_x \left(\int_0^\infty e^{-\alpha t} dL(t) \right) \quad \text{for q.e. } x \in S. \quad (2.20)$$

In particular, (2.20) holds for $x = a$. $L(t)$ is a local time at $\{a\}$ in the sense that it increases only when $X_t = a$:

$$L(t) = \int_0^t I_a(X_s) dL(s).$$

We consider the right continuous inverse $S(t) = \inf\{s : L(s) > t\}$ of $L(t)$.

It is well known that the increasing process $(S(t), P_a)$ is a subordinator killed upon an exponential holding time (cf.[2]). Theorem 2.1 enables us to identify the Lévy measure of the subordinator and the killing rate. Indeed, according to [2, v (3.17)], (2.20) implies the identity

$$E_a(e^{-\alpha S(t)}) = \exp(-t/\widetilde{U}_\alpha \delta_a(a)),$$

which combined with (2.16) leads us to

$$E_a \left(e^{-\alpha S(t)} \right) = e^{-tL(m_0, \psi)} \exp[-t\alpha(u_\alpha, \varphi)]. \quad (2.21)$$

We need a lemma which will play a basic role in §4 again. A family $\{\nu_t\}_{t>0}$ of σ -finite measures on S_0 is called an X^0 -entrance law if $\nu_t p_s^0 = \nu_{s+t}$, $s, t > 0$. Then $\nu_t(f)$, $f \in \mathcal{B}^+(S_0)$, is measurable in t and we may let

$$\hat{\nu}_\alpha(f) = \int_0^\infty e^{-\alpha t} \nu_t(f) dt, \quad \alpha > 0, \quad f \in \mathcal{B}^+(S_0).$$

Lemma 2.2. (i) *There exists a unique X^0 -entrance law $\{\mu_t\}$ such that*

$$m_0 = \int_0^\infty \mu_t dt. \quad (2.22)$$

(ii) $\hat{\mu}_\alpha(f) = (u_\alpha, f)$, $\alpha > 0$, $f \in \mathcal{B}^+(S_0)$.

Consequently,

$$\int_0^t \mu_s(f) ds = \int_{S_0} P_x^0(\zeta^0 \leq t, X_{\zeta^0-} = a) f(x) m(dx), \quad t > 0, \quad f \in \mathcal{B}(S_0). \quad (2.23)$$

(iii) $\mu_t(S_0) < \infty$, $t > 0$.

(iv) *For any bounded X^0 -excessive function v on S_0 , $\mu_t(v)$ is right continuous in $t > 0$.*

(v) *For any X^0 -excessive function v on S_0 , the energy functional $L(m_0, v)$ introduced in Lemma 2.1 admits an expression*

$$L(m_0, v) = \lim_{t \downarrow 0} \mu_t(v).$$

When v is p_t^0 -invariant, it holds for any $t > 0$ that

$$L(m_0, v) = \mu_t(v).$$

(vi) $L(m_0, \varphi) = \infty$.

Proof. (i) Since

$$p_t^0 \varphi(x) = P_x^0(t < \zeta^0 < \infty, X_{\zeta^-}^0 = a) \downarrow 0, \quad t \rightarrow \infty,$$

$\lim_{t \downarrow 0} m_0 p_t^0(f) = (p_t^0 \varphi, f) = 0$ for $f \in L^1(S_0, m)$, namely, m_0 is purely excessive. Hence the desired assertion follows from a well known representation theorem provided that X^0 is transient ([11, Th. 5.25]). But the present situation can be reduced to this case by observing that

$$S_1 = \{x \in S_0 : \varphi(x) > 0\}$$

is a non-trivial X^0 -invariant set q.e. and the restriction of X^0 to S_1 is transient (cf. [9, §4.6]).

(ii) For $f \in C_0^+(S_0)$, we have

$$\int_t^\infty \mu_t(f) dt = \int_0^\infty \mu_{t+s}(f) dt = \int_0^\infty \mu_s(p_t^0 f) ds = (\varphi, p_t^0 f),$$

and

$$\mu_t(f) = -\frac{d}{dt}(\varphi, p_t^0 f), \quad \text{a.e. } t.$$

Hence

$$\begin{aligned} \hat{\mu}_\alpha(f) &= -\int_0^\infty e^{-\alpha t} \frac{d}{dt}(\varphi, p_t^0 f) dt \\ &= [-e^{-\alpha t}(\varphi, p_t^0 f)]_0^\infty - \alpha \int_0^\infty e^{-\alpha t}(\varphi, p_t^0 f) dt \\ &= (\varphi, f) - \alpha(\varphi, G_\alpha^0 f) = (\varphi - \alpha G_\alpha^0 \varphi, f) = (u_\alpha, f). \end{aligned}$$

(iii) By (ii) and Theorem 2.1 (ii), $\hat{\mu}_\alpha(1) = (u_\alpha, 1) < \infty$, from which the desired finiteness follows.

(iv) On account of (iii), we have $\mu_{t+s}(v) = \mu_t(p_s^0 v) \rightarrow \mu_t(v)$, $s \downarrow 0$.

(v) Since $\langle \mu_t, v \rangle$ is increasing as $t \downarrow 0$ (independent of t when v is p_t^0 -invariant), the assertions follow from

$$\langle m_0 - m_0 p_t^0, v \rangle = \int_0^t \langle \mu_s, v \rangle ds.$$

(vi) Since $S(t)$ is the right continuous inverse of an increasing continuous process $L(t)$, $P_a(S(t) > 0) = 1$ and consequently we have

$$L(m_0, \varphi) = \lim_{\alpha \rightarrow \infty} \alpha(u_\alpha, \varphi) = \infty$$

by letting $\alpha \rightarrow \infty$ in (2.21). □

We see by the above lemma that $\mu_t(\varphi)$ is decreasing and right continuous in $t > 0$ and so we can define a measure Θ on $(0, \infty)$ by

$$\Theta((s, t]) = \mu_s(\varphi) - \mu_t(\varphi), \quad 0 < s < t. \quad (2.24)$$

It then holds that

$$\Theta((s, t]) = \mu_s(\varphi - p_{t-s}^0 \varphi) = \langle \mu_s, P.(\sigma \leq t - s) \rangle,$$

and we get by letting $t \rightarrow \infty$,

$$\Theta((s, \infty)) = \mu_s(\varphi). \quad (2.25)$$

We note that

$$\Theta([\delta, \infty)) < \infty$$

for each $\delta > 0$ by virtue of Lemma 2.2 (iii).

Lemma 2.3. *It holds that*

$$\alpha(u_\alpha, \varphi) = \int_0^\infty (1 - e^{-\alpha u}) \Theta(du).$$

Proof. we have from Lemma 2.2 (ii) and (2.25)

$$\begin{aligned} \alpha(u_\alpha, \varphi) &= \alpha \hat{\mu}_\alpha(\varphi) = \alpha \int_0^\infty e^{-\alpha t} \Theta((t, \infty)) dt \\ &= \int_0^\infty \int_0^s \alpha e^{-\alpha t} dt \Theta(ds) = \int_0^\infty (1 - e^{-\alpha s}) \Theta(ds). \end{aligned}$$

□

On account of the formula (2.21), Lemma 2.3 and by noting that $\lim_{\alpha \downarrow 0} \alpha(u_\alpha, \varphi) = 0$, we can get the next theorem from [2, Theorem 3.21].

Theorem 2.2. *Define a measure Θ on $(0, \infty)$ by (2.24). On a certain probability space (Ω, \mathcal{B}, P) , construct a subordinator $\{Y_t\}_{t \geq 0}$ with Lévy measure Θ and zero drift and a random variable Z , independent of $\{Y_t\}$, with*

$$P(Z \geq t) = e^{-L(m_0, \psi)t}, \quad t \geq 0.$$

If we let

$$S^*(t) = \begin{cases} Y(t) & t < Z, \\ \infty & t \geq Z, \end{cases}$$

then the process $(\{S^*(t)\}_{t \geq 0}, P)$ is equivalent in law to $(\{S(t)\}_{t \geq 0}, P_a)$.

3 Strongly local Dirichlet form with a recurrent point

Let S and m be as in §2. In this section, we consider a special case of the Dirichlet form of §2 for which the point a is recurrent.

3.1 Description of associated Poisson point process and entrance law

Let $(\mathcal{E}, \mathcal{F})$ be a strongly local regular Dirichlet form on $L^2(S; m)$ and $X = (X_t, P_x)$ be an associated diffusion on S . In place of the assumption **B.1** of §2, let us assume that

B.2 $\varphi(x) > 0$ m -a.e. $x \in S_0$

B.3 $1 \in \mathcal{F}_e$ and $\mathcal{E}(1, 1) = 0$.

In the next subsection, we shall construct a typical example of a Dirichlet form $(\mathcal{E}, \mathcal{F})$ satisfying these conditions by a method of the one point compactification.

The assumption **B.2** implies that $u_1 > 0$, m -a.e. and $\text{Cap}(\{a\}) = \mathcal{E}_1(u_1, u_1) \geq (u_1, u_1) > 0$, namely, the assumption **B.1** of §1 (cf. [9, Lemma 4.2.1]). Further, the Dirichlet form $(\mathcal{E}, \mathcal{F})$ becomes irreducible because, from (2.15), we have for any Borel sets $B_1, B_2 \subset S$ of positive m -measures

$$(I_E, G_\alpha I_F) \geq (u_\alpha, I_E)(u_\alpha, I_F)/\alpha(u_\alpha, \varphi) > 0.$$

Since $(\mathcal{E}, \mathcal{F})$ is recurrent by **B.3**, we have actually the property

$$\varphi(x) = 1, \quad \text{q.e. } x \in S, \quad (3.1)$$

stronger than the assumption **B.2** in view of [9, Th.4.6.6].

Thus the point a is not only regular for itself, instantaneous, but also recurrent. (2.15) is now reduced to

$$G_\alpha f(x) = G_\alpha^0 f(x) + \frac{(u_\alpha, f)}{\alpha(u_\alpha, 1)} u_\alpha(x), \quad x \in S, \quad G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, 1)}. \quad (3.2)$$

The positive continuous additive functional $L(t)$ of X associated with the unit mass δ_a has the property that $L(\infty) = \infty$ and its right continuous inverse $S(t)$ is a subordinator satisfying

$$E_a \left(\int_0^\infty e^{-\alpha S(s)} ds \right) = \frac{1}{\alpha(u_\alpha, 1)} \quad (3.3)$$

on account of (2.16) and (2.20).

Therefore we can follow directly the argument of [15, §6 case 2(b)] to conclude that

$$D_{\mathbf{p}} = \{s : S(s) - S(s-) > 0\}, \quad (3.4)$$

$$\mathbf{p}_s(t) = X_{S(s-)+t}, \quad s \in D_{\mathbf{p}}, \quad 0 \leq t < S(s) - S(s-), \quad (3.5)$$

defines, under the law P_a , a W_a -valued Poisson point process \mathbf{p} , where W_a is the space of continuous excursions in S_0 from a to a :

$$W_a = \{w : [0, \zeta(w)) \rightarrow S_0, \text{ continuous}, 0 < \zeta(w) < \infty, w(0) = a, w(\zeta-) = a\}. \quad (3.6)$$

Let \mathbf{n} be the characteristic measure of the Poisson point process \mathbf{p} . Then \mathbf{n} is a σ -finite measure on the space W_a and $\{w(t), \mathbf{n}\}$ is Markovian with respect to the transition function p_t^0 of X^0 . The entrance law $\{\nu_t\}$ associated with the characteristic measure \mathbf{n} is defined by

$$\nu_t(B) = \mathbf{n}\{w : \zeta(w) > t, w(t) \in B\}, \quad B \in \mathcal{B}(S), \quad t > 0. \quad (3.7)$$

Recall that we have already considered an X^0 -entrance law $\{\mu_t\}$ specified by (2.22) which is now reduced to

$$m = \int_0^\infty \mu_t dt. \quad (3.8)$$

The description (2.23) of $\{\mu_t\}$ now reads

$$\int_0^t \mu_s(f) ds = \int_{S_0} P_x^0(\zeta^0 \leq t) f(x) m(dx), \quad t > 0, \quad f \in \mathcal{B}(S_0). \quad (3.9)$$

Theorem 3.1. $\nu_t = \mu_t, \quad t > 0.$

Proof. By virtue of Lemma 2.2, it suffices to show that

$$\hat{\nu}_\alpha(f) = (u_\alpha, f), \quad f \in \mathcal{B}_b(S_0). \quad (3.10)$$

We make use of the next general formula

$$E_a \left(\sum_{s \leq t} a(s, \mathbf{p}_s, \omega) \right) = E_a \left(\int_{W_a \times (0, t]} a(s, w, \omega) \mathbf{n}(dw) ds \right) \quad (3.11)$$

holding for any non-negative predictable function $a(s, w, \omega)$ on $[0, \infty) \times W_a \times \Omega$, Ω being a filtered sample space on which the diffusion process X is defined (cf. [14, p62].)

Since $m(\{a\})$ is assumed to be zero, $\int_0^\infty I_a(X_t) dt = 0$, P_a -almost surely. By (3.4) and (3.5), we have for $f \in B_b(S)$,

$$\begin{aligned} G_\alpha f(a) &= E_a \left(\int_0^\infty e^{-\alpha t} f(X_t) dt \right) = E_a \left(\sum_{s > 0} \int_{S(s-)}^{S(s)} e^{-\alpha t} f(X_t) dt \right) \\ &= E_a \left(\sum_{s > 0} e^{-\alpha S(s-)} \int_0^{\zeta(\mathbf{p}_s)} e^{-\alpha t} f(\mathbf{p}_s(t)) dt \right). \end{aligned}$$

We let

$$\Gamma(w) = \int_0^{\zeta(w)} e^{-\alpha t} f(w(t)) dt.$$

$a(s, w, \omega) = \Gamma(w) \cdot e^{-\alpha S(s-, \omega)}$ is then predictable and we get by (3.11)

$$\begin{aligned} G_\alpha f(a) &= E_a \left(\sum_{s > 0} e^{-\alpha S(s-)} \Gamma(\mathbf{p}_s) \right) \\ &= \int_{W_a} \Gamma(w) \mathbf{n}(dw) \cdot \int_0^\infty E_a \left(e^{-\alpha S(s)} \right) ds. \end{aligned}$$

Since

$$\int_{W_a} \Gamma(w) \mathbf{n}(dw) = \hat{\nu}_\alpha(f),$$

(3.2) and (3.3) lead us to the desired identity (3.10). \square

By Theorem 3.1 and [15, Th. 6.3], the finite dimensional distribution of $\{W_a, \mathbf{n}\}$ can be described as follows:

$$\int_{W_a} f_1(w(t_1)) f_2(w(t_2)) \cdots f_n(w(t_n)) \mathbf{n}(dw) = \mu_{t_1} f_1 p_{t_2-t_1}^0 f_2 \cdots p_{t_{n-1}-t_{n-2}}^0 f_{n-1} p_{t_n-t_{n-1}}^0 f_n, \quad (3.12)$$

for any $0 < t_1 < t_2 < \cdots < t_{n-1}, t_n$, $f_1, f_2, \cdots, f_n \in B_b(S_0)$. Here we use the convention that $w \in W$ satisfies $w(t) = \Delta, \forall t \geq \zeta(w)$, and any function f on S_0 is extended to $S_0 \cup \Delta$ by setting $f(\Delta) = 0$.

In §4, we shall start with an m -symmetric diffusion X^0 on S_0 and an expression like the above with μ_t being specified by (2.22). See §4 for the abbreviated notation appearing on the right hand side of (3.12).

Actually Theorem 3.1 can be extended to a general case where condition **B.3** of the recurrence is not assumed as we shall see in Remark 4.2 at the end of §4.

We note that the excursion law around a regular point of a general Markov process can be also formulated in terms of Maisonneuve's exit system[5]. Some property of the integral in t of the associated entrance law was investigated by R.K. Gettoor [10].

3.2 Construction of form by one-point compactification

In this subsection, we start with a Dirichlet form with underlying space S_0 and extend it by the one-point compactification to a Dirichlet form with underlying space $S = S_0 \cup a$ satisfying **B.2** and **B.3** (and consequently **B.1**).

Let S_0 be a locally compact separable metric space and m be a bounded positive measure on S_0 with $\text{Supp}[m] = S_0$. We consider a regular strongly local Dirichlet form $(\mathcal{E}, \mathcal{F}_0)$ on $L^2(S_0; m)$ satisfying the *Poincaré inequality*:

$$(u, u) \leq A \cdot \mathcal{E}(u, u) \quad u \in \mathcal{F}_0 \quad \exists A > 0. \quad (3.13)$$

Denote by $S = S_0 \cup a$ the one-point compactification of S_0 and by $L^2(S; m)(= L^2(S_0; m))$ the space of square integrable functions on S with respect to $I_{S_0} \cdot m$. Let us introduce a space $(\mathcal{E}, \mathcal{F})$ by

$$\mathcal{F} = \mathcal{F}_0 + \text{constant functions on } S, \quad (3.14)$$

$$\mathcal{E}(w_1, w_2) = \mathcal{E}(f_1, f_2), \quad w_1 = f_1 + c_1, \quad w_2 = f_2 + c_2, \quad f_i \in \mathcal{F}_0, \quad c_i \text{ constant.} \quad (3.15)$$

Theorem 3.2. (i) $(\mathcal{E}, \mathcal{F})$ is a regular strongly local Dirichlet form on $L^2(S; m)$ possessing as its core the space

$$\mathcal{C} = \mathcal{C}_0 + \text{constant functions on } S_0,$$

where $\mathcal{C}_0 = \mathcal{F}_0 \cap C_0(S_0)$.

(ii) $(\mathcal{E}, \mathcal{F})$ and the associated diffusion on S satisfy **B.2**, **B.3**.

Proof. (i) Suppose $f \in \mathcal{F}_0$ is a constant. By the regularity of $(\mathcal{E}, \mathcal{F}_0)$, there exist $f_n \in \mathcal{F}_0 \cap C_0(S_0)$ which are \mathcal{E}_1 -convergent to f . We have then $\mathcal{E}(f, f) = \lim_{n \rightarrow \infty} \mathcal{E}(f, f_n) = 0$ on account of the strong locality of $(\mathcal{E}, \mathcal{F}_0)$ and Remark 2.1 stated in the beginning of §2.1. (3.13) then implies $f = 0$ and the definition (3.14) and (3.15) makes sense.

If $w_n = f_n + c_n \in \mathcal{F}$ is an \mathcal{E}_1 -Cauchy sequence, then f_n is \mathcal{E}_1 -convergent to some $f \in \mathcal{F}_0$ by (3.13) and hence w_n is \mathcal{E}_1 -convergent to $f + c$ for some constant c .

Clearly \mathcal{C} is dense both in \mathcal{F} and $C(S)$, namely, $(\mathcal{E}, \mathcal{F})$ is regular.

Suppose, for $w_i = f_i + c_i \in \mathcal{C}$, that w_1 is constant on a neighbourhood of $\text{Supp}(w_2)$. When $c_2 = 0$, $\mathcal{E}(w_1, w_2) = 0$ by the strong locality of $(\mathcal{E}, \mathcal{F}_0)$. When $c_2 \neq 0$, the set $U = S \setminus \text{Supp}(w_2)$ is either empty or a non-empty relatively compact open subset of S_0 . In the former case, $f_1 = 0$ and $\mathcal{E}(w_1, w_2) = 0$. In the latter case, $f_2 = -c_2$ on U , while $\text{Supp}(f_1) \subset U$ and $\mathcal{E}(w_1, w_2) = \mathcal{E}(f_1, f_2) = 0$ again. Hence $(\mathcal{E}, \mathcal{F})$ is strongly local on account of [9, Th.3.1.2].

The Markov property

$$w \in \mathcal{F} \Rightarrow v = (0 \vee w) \wedge 1 \in \mathcal{F}, \mathcal{E}(v, v) \leq \mathcal{E}(w, w)$$

is evident, because, for $w = f + c$, $w \in \mathcal{F}_0$, c constant, we have $v = [(-c) \vee f] \wedge (1 - c) + c$.

(ii) **B.2** follows from the Poincaré inequality (3.13). Denote by X and $X^0 = (X_t^0, P_x^0, \zeta^0)$ the diffusions associated with $(\mathcal{E}, \mathcal{F})$ and $(\mathcal{E}, \mathcal{F}_0)$ respectively. Then X^0 is the part of X on S_0 and hence

$$\varphi(x) = P_x^0(\zeta^0 < \infty), \quad x \in S_0,$$

Denote by G^0 the 0-order resolvent operator of X^0 . Since $m(S_0) < \infty$, (3.13) implies that $G^0 1 \in \mathcal{F}_0$ and

$$E_x^0(\zeta^0) = G^0 1(x) < \infty \quad \text{q.e.}$$

proving (3.1). It is obvious from (3.14),(3.15) that $1 \in \mathcal{F}$ and $\mathcal{E}(1, 1) = 0$. \square

$(\mathcal{E}, \mathcal{F}_0)$ is not necessarily irreducible on S_0 , but $(\mathcal{E}, \mathcal{F})$ defined by (3.14),(3.15) is irreducible recurrent on S in view of the observation made in the preceding subsection. See Example 6.2.

4 Construction of a symmetric extension via excursion valued Poisson point processes

In this section, we start with an m -symmetric diffusion X^0 on S_0 and construct first an excursion law with which Poisson point processes of two different kinds of excursions around the point a are associated. We then construct an m -symmetric diffusion \tilde{X} on $S = S_0 \cup a$ by piecing together those excursions. The resolvent of the resulting diffusion \tilde{X} turns out to be identical with (2.15).

4.1 An excursion law and its basic properties

Let S be a locally compact separable metric space and a be a non-isolated point of S . We put $S_0 = S \setminus \{a\}$. The one point compactification of S is denoted by S_Δ . When S is compact already, Δ is added as an isolated point. Let m be a positive Radon measure on S_0 with $\text{Supp}[m] = S_0$. m is extended to S by setting $m(\{a\}) = 0$.

We assume that we are given an m -symmetric diffusion $X^0 = (X_t^0, P_x^0)$ on S_0 with life time ζ^0 satisfying the following:

$$\mathbf{A.1} \quad P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 \in \{a\} \cup \{\Delta\}) = P_x^0(\zeta^0 < \infty), \quad \forall x \in S_0.$$

We define the functions $\varphi, u_\alpha, \psi^{(1)}, \psi^{(2)}, \psi$ by (2.5) and (2.6), namely, for $x \in S_0$,

$$\varphi(x) = P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 = a), \quad u_\alpha(x) = E_x^0(e^{-\alpha\zeta^0}; X_{\zeta^0-} = a),$$

$$\psi = 1 - \varphi = \psi^{(1)} + \psi^{(2)}, \quad \psi^{(1)}(x) = P_x^0(\zeta^0 < \infty, X_{\zeta^0-} = \Delta), \quad \psi^{(2)}(x) = P_x^0(\zeta^0 = \infty).$$

Let us assume that

$$\mathbf{A.2} \quad \varphi(x) > 0, \quad \forall x \in S_0,$$

and

$$\mathbf{A.3} \quad u_\alpha \in L^1(S_0; m), \quad \forall \alpha > 0.$$

Denote by p_t^0, G_α^0 the transition function and the resolvent of X^0 respectively. Our last assumption concerns the regularity:

$$\mathbf{A.4} \quad u_\alpha \in C_b(S_0), \quad G_\alpha^0(C_b(S_0)) \subset C_b(S_0), \quad \alpha > 0,$$

where $C_b(S_0)$ is the space of all bounded continuous functions on S_0 .

The measure m could be infinite on a compact neighbourhood of a in S , but it is finite on each level set of u_α due to the condition **A.3**. We also note here the next relation which will be utilized in the sequel:

$$u_\alpha(x) = \varphi(x) - \alpha G_\alpha^0 \varphi(x) \leq 1 - \alpha G_\alpha^0 1(x), \quad x \in S_0.$$

Define m_0 by

$$m_0 = \varphi \cdot m,$$

which is an X^0 -excessive measure with $m_0 p_t^0 = p_t^0 \varphi \cdot m$. In view of Lemma 2.2, there exists a unique X^0 -entrance law $\{\mu_t\}$ related to the measure m_0 by (2.22), namely,

$$m_0 = \int_0^\infty \mu_t dt.$$

and it satisfies that

$$\hat{\mu}_\alpha(f) = (u_\alpha, f), \quad f \in \mathcal{B}^+(S_0). \quad (4.1)$$

On account of the assumption **(A.3)**, we then have that

$$\mu_t(S_0) < \infty, \quad t > 0, \quad \int_0^1 \mu_t(S_0) dt < \infty. \quad (4.2)$$

We now introduce the spaces W' , W of excursions by

$$\begin{aligned} W' &= \{w : \exists \zeta(w) \in (0, \infty], \text{ } w \text{ is a continuous function from } (0, \zeta(w)) \text{ to } S_0\}, \\ W &= \{w \in W' : \text{if } \zeta(w) < \infty, \text{ then } \exists w(\zeta(w)-) \in \{a\} \cup \{\Delta\}\}. \end{aligned} \quad (4.3)$$

$\zeta(w)$ will be called the *terminal time* of the excursion w .

We are concerned with a measure \mathbf{n} on the space W specified in terms of the entrance law $\{\mu_t\}$ and the transition function p_t^0 by

$$\int_W f_1(w(t_1)) f_2(w(t_2)) \cdots f_n(w(t_n)) \mathbf{n}(dw) = \mu_{t_1} f_1 p_{t_2-t_1}^0 f_2 \cdots p_{t_{n-1}-t_{n-2}}^0 f_{n-1} p_{t_n-t_{n-1}}^0 f_n, \quad (4.4)$$

for any $0 < t_1 < t_2 < \cdots < t_n$, $f_1, f_2, \cdots, f_n \in B_b(S_0)$. Here, we use the convention that $w \in W$ satisfies $w(t) = \Delta, \forall t \geq \zeta(w)$, and any function f on S_0 is extended to $S_0 \cup \Delta$ by setting $f(\Delta) = 0$. Further, on the right hand side of (4.4), we employ an abbreviated notation for the repeated operations

$$\mu_{t_1} [f_1 p_{t_2-t_1}^0 \{f_2 \cdots p_{t_{n-1}-t_{n-2}}^0 (f_{n-1} p_{t_n-t_{n-1}}^0 f_n)\}].$$

Proposition 4.1. *There exists a unique measure \mathbf{n} on the space W satisfying (4.4).*

Proof. Let \mathbf{n} be the Kuznetsov measure on W' uniquely associated with the transition semigroup $\{p_t^0\}$ and the entrance rule $\{\eta_u\}$ defined by

$$\eta_u = 0 \quad \text{for } u \leq 0, \quad \eta_u = \mu_u \quad \text{for } u > 0$$

as is constructed in [5, Chap XIX, 9] for a right semigroup. Because of the present choice of the entrance rule, it holds that $\alpha = 0$ where α is the birth time which is random in general (cf. [11, p54].)

On account of the assumption **A.1** for the diffusion X^0 on S_0 , the same method of the construction of the Kuznetsov measure as in [5, Chap.XIX, 9] works in proving that \mathbf{n} is supported by the space W and satisfies (4.4). \square

We call \mathbf{n} the *excursion law* associated with the entrance law $\{\mu_t\}$. We split the space W of excursions into two parts:

$$W^+ = \{w \in W : \zeta(w) < \infty, w(\zeta-) = a\}, \quad W^- = W \setminus W^+. \quad (4.5)$$

Note that $W^- = W_1^- \cup W_2^-$ with

$$W_1^- = \{w \in W : \zeta(w) < \infty, w(\zeta-) = \Delta\}, \quad W_2^- = \{w \in W : \zeta(w) = \infty\}.$$

For $w \in W^+$, we define $\hat{w} \in W$ by

$$\hat{w}(t) = w(\zeta - t), \quad 0 < t < \zeta. \quad (4.6)$$

The next lemma says that the restriction of the excursion law to W^+ is invariant under time reversion. This is a present variant of the time reversal arguments that have been formulated in general contexts ([23], [12], [6], [7]).

Lemma 4.1. *For any $t_k > 0$ and $f_k \in \mathcal{B}_b(S_0)$, ($1 \leq k \leq n$),*

$$\mathbf{n} \left\{ \prod_{k=1}^n f_k(w(t_1 + \cdots + t_k)); W^+ \right\} = \mu_{t_1} f_1 p_{t_2}^0 f_2 \cdots p_{t_{n-1}}^0 f_{n-1} p_{t_n}^0 f_n \varphi, \quad (4.7)$$

$$\mathbf{n} \left\{ \prod_{k=1}^n f_k(w(t_1 + \cdots + t_k)); W^+ \right\} = \mathbf{n} \left\{ \prod_{k=1}^n f_k(\hat{w}(t_1 + \cdots + t_k)); W^+ \right\}. \quad (4.8)$$

Proof. (4.7) readily follows from (4.4) and the Markov property of \mathbf{n} . As for (4.8) we observe that, for $\alpha_1, \dots, \alpha_n > 0$,

$$\int_0^\infty \cdots \int_0^\infty e^{-\alpha_1 t_1 - \cdots - \alpha_n t_n} \mathbf{n} \left\{ \prod_{k=1}^n f_k(w(t_1 + \cdots + t_k)); W^+ \right\} dt_1 \cdots dt_n \quad (4.9)$$

equals

$$\mathbf{n}\{F(w); \zeta < \infty, w(\zeta-) = a\}$$

with

$$F(w) = \int \cdots \int_{0 < t_1 < \cdots < t_n < \zeta} \prod_{k=1}^n \left\{ e^{-\alpha_k(t_k - t_{k-1})} f_k(w(t_k)) \right\} dt_1 \cdots dt_n, \quad (t_0 = 0).$$

Hence, for (4.8), it suffices to prove

$$\mathbf{n}\{F(w); \zeta < \infty, w(\zeta-) = a\} = \mathbf{n}\{F(\hat{w}); \zeta < \infty, w(\zeta-) = a\}. \quad (4.10)$$

Performing the change of variables

$$\zeta - t_k = s_k, \quad 1 \leq k \leq n,$$

in the expression of $F(\hat{w})$ and by noting that

$$\begin{aligned} t_k = \zeta - s_k, \quad t_k - t_{k-1} = s_{k-1} - s_k, \quad 1 \leq k \leq n, \quad s_0 = \zeta, \\ 0 < t_1 < \cdots < t_n < \zeta \iff 0 < s_n < \cdots < s_1 < \zeta, \end{aligned}$$

we obtain

$$\begin{aligned} F(\hat{w}) &= \int \cdots \int_{0 < s_n < \cdots < s_1 < \zeta} \prod_{k=1}^n \left\{ e^{-\alpha_k(s_{k-1}-s_k)} f_k(w(s_k)) \right\} ds_1 \cdots ds_n \\ &= \int \cdots \int_{0 < s_n < \cdots < s_1 < \infty} \Gamma_{s_1 \cdots s_n}(w) ds_1 \cdots ds_n \end{aligned}$$

with

$$\Gamma_{s_1 \cdots s_n}(w) = \prod_{k=2}^n \left\{ e^{-\alpha_k(s_{k-1}-s_k)} f_k(w(s_k)) \right\} \cdot e^{-\alpha_1(\zeta-s_1)} f_1(w(s_1)) I_{(0,\zeta)}(s_1).$$

On the other hand, we get from (4.4) and the Markov property of \mathbf{n} that

$$\begin{aligned} &\mathbf{n}\{\Gamma_{s_1 s_2 \cdots s_n}(w); \zeta < \infty, w(\zeta-) = a\} \\ &= \mathbf{n}\left\{ f_n(w(s_n)) e^{-\alpha_n(s_{n-1}-s_n)} \cdots \right. \\ &\quad \left. f_2((w(s_2)) e^{-\alpha_3(s_1-s_2)} f_1(w(s_1)) u_{\alpha_1}(w(s_1)); s_1 < \zeta) \right\} \\ &= e^{-\alpha_n(s_{n-1}-s_n) - \alpha_{n-1}(s_{n-2}-s_{n-1}) - \cdots - \alpha_2(s_1-s_2)} \cdot \\ &\quad \mu_{s_n} f_n p_{s_{n-1}-s_n}^0 f_{n-1} p_{s_{n-2}-s_{n-1}}^0 f_{n-1} \cdots p_{s_2-s_3}^0 f_2 p_{s_1-s_2}^0 f_1 u_{\alpha_1}. \end{aligned}$$

Therefore,

$$\mathbf{n}\{F(\hat{w}); \zeta < \infty, w(\zeta-) = a\} = \int_0^\infty ds_n \mu_{s_n} f_n G_{\alpha_n}^0 f_{n-1} G_{\alpha_{n-1}}^0 \cdots f_3 G_{\alpha_3}^0 f_2 G_{\alpha_2}^0 f_1 u_{\alpha_1}.$$

In view of (2.7), the symmetry of G_α^0 , (4.7) and (4.9), we arrive at

$$\begin{aligned} &\mathbf{n}\{F(\hat{w}); \zeta < \infty, w(\zeta-) = a\} = \langle m_0, f_n G_{\alpha_n}^0 f_{n-1} G_{\alpha_{n-1}}^0 \cdots f_3 G_{\alpha_3}^0 f_2 G_{\alpha_2}^0 f_1 u_{\alpha_1} \rangle \\ &= (f_n \varphi, G_{\alpha_n}^0 f_{n-1} G_{\alpha_{n-1}}^0 \cdots f_3 G_{\alpha_3}^0 f_2 G_{\alpha_2}^0 f_1 u_{\alpha_1}) = (f_1 G_{\alpha_2}^0 f_2 G_{\alpha_3}^0 f_3 \cdots G_{\alpha_n} f_n \varphi, u_{\alpha_1}) \\ &= \int_0^\infty e^{-\alpha_1 t_1} \mu_{t_1} f_1 G_{\alpha_2}^0 f_2 G_{\alpha_3}^0 f_3 \cdots G_{\alpha_n}^0 f_n \varphi dt_1 = \mathbf{n}\{F(w); \zeta < \infty, w(\zeta-) = a\} \end{aligned}$$

the desired identity (4.10). \square

Next we put

$$W_a = \{w \in W : \lim_{t \downarrow 0} w(t) = a\}. \quad (4.11)$$

Lemma 4.2. $\mathbf{n}\{W \setminus W_a\} = 0$.

Proof. The preceding lemma implies that

$$\begin{aligned} \mathbf{n}\{W^+ \setminus W_a\} &= \mathbf{n}\{W^+ \cap (w(0+) = a)^c\} \\ &= \mathbf{n}\{W^+ \cap (\hat{w}(0+) = a)^c\} = \mathbf{n}\{W^+ \cap (w(\zeta-) = a)^c\} = 0. \end{aligned}$$

We then have for each $t > 0$

$$\mathbf{n}\{\varphi(w(t)); (\zeta > t) \cap (w(0+) = a)^c\} = \mathbf{n}\{(W^+ \setminus W_a) \cap (\zeta > t)\} = 0,$$

which combined with the assumption **A.2** leads us to

$$\mathbf{n}\{(W \setminus W_a) \cap (\zeta > t)\} = 0.$$

It then suffices to let $t \downarrow 0$. □

Lemma 4.3. *For any neighbourhood U of a in S , we let*

$$\tau_{U^c} = \inf\{t > 0 : w(t) \in U^c\}, \quad w \in W.$$

It holds then that

$$\mathbf{n}\{\tau_{U^c} < \zeta\} < \infty.$$

Proof. We may assume that the closure \bar{U} in S is compact. Let $f(x) = \varphi(x) - u_1(x)$, $x \in S_0$. Then

$$f(x) = E_x^0 \left\{ 1 - e^{-\zeta^0}; \zeta^0 < \infty, X_{\zeta^0-} = a \right\} > 0, \quad \forall x \in S_0.$$

Since $u_\alpha(x) - u_1(x) \uparrow f(x)$, $\alpha \downarrow 0$, the assumption **A.3** implies that f is lower semicontinuous on S_0 and hence

$$c = \inf_{x \in \partial U} f(x)$$

is positive. We then have, for each $\delta > 0$ and $x \in \partial U$,

$$\begin{aligned} P_x^0(\delta < \zeta^0 < \infty, X_{\zeta^0-} = a) &\geq E_x^0 \left\{ 1 - e^{-\zeta^0}; \delta < \zeta^0 < \infty, X_{\zeta^0-} = a \right\} \\ &\geq c - E_x^0 \left\{ 1 - e^{-\zeta^0}; \zeta^0 \leq \delta, X_{\zeta^0-} = a \right\} \geq c - (1 - e^{-\delta}). \end{aligned}$$

Choose $\delta > 0$ so small that

$$r = c - (1 - e^{-\delta})$$

is positive. For such δ ,

$$P_x^0(\delta < \zeta^0 < \infty, X_{\zeta^0-} = a) \geq r, \quad \forall x \in \partial U. \quad (4.12)$$

We shall use the notation τ_{U^c} not only for $w \in W$ but also for the sample path of the Markov process X^0 . Using the preceding lemma, (4.12) and (4.2), we are led to

$$\begin{aligned} \mathbf{n}\{\tau_{U^c} < \zeta\} &= \lim_{\epsilon \downarrow 0} \mathbf{n}\{\epsilon < \tau_{U^c} < \zeta\} = \lim_{\epsilon \downarrow 0} \int_U \mu_\epsilon(dx) P_x^0 \{\tau_{U^c} < \zeta^0\} \\ &\leq \overline{\lim}_{\epsilon \downarrow 0} \int_U \mu_\epsilon(dx) E_x^0 \left\{ r^{-1} P_{X_{\tau_{U^c}}^0}^0(\delta < \zeta^0 < \infty, X_{\zeta^0-} = a); \tau_{U^c} < \zeta^0 \right\} \\ &\leq r^{-1} \lim_{\epsilon \downarrow 0} \int_{S_0} \mu_\epsilon(dx) P_x^0(\delta < \zeta^0 < \infty, X_{\zeta^0-} = a) \leq r^{-1} \lim_{\epsilon \downarrow 0} \int_{S_0} \mu_\epsilon(dx) P_x^0(\delta < \zeta^0) \\ &= r^{-1} \lim_{\epsilon \downarrow 0} \mu_{\epsilon+\delta}(S_0) \leq r^{-1} \mu_\delta(S_0) < \infty. \end{aligned}$$

□

The next lemma states a relation of the excursion law \mathbf{n} to energy functionals $L(m_0, v)$ introduced in Lemma 2.1.

Lemma 4.4.

- (i) $\mathbf{n}(W^+) = L(m_0, \varphi)$, $\mathbf{n}(W^-) = L(m_0, \psi)$, $\mathbf{n}(W_i^-) = L(m_0, \psi^{(i)})$, $i = 1, 2$.
- (ii) $\mathbf{n}(W_1^-) < \infty$, $\mathbf{n}(W_2^-) = \mu_t(\psi^{(2)}) = \alpha \hat{\mu}_\alpha(\psi^{(2)}) = \alpha(u_\alpha, \psi^{(2)}) < \infty$, $t > 0, \alpha > 0$.

Proof. (i) Since $\mathbf{n}(\zeta > t; W^+) = \langle \mu_t, \varphi \rangle$, the first identity follows from Lemma 2.2 (v) by letting $t \downarrow 0$. The proof of the other identities is the same.

(ii) Take a neighbourhood U of a in S with compact \bar{U} . We have then by the preceding lemma

$$\mathbf{n}(W_1^-) = \mathbf{n}(\zeta < \infty, w(\zeta-) = \Delta) \leq \mathbf{n}\{\tau_{U^c} < \zeta\} < \infty.$$

Since $\psi^{(2)}$ is p_t^0 -invariant, the second assertion follows from (i), Lemma 2.1, Lemma 2.2 and assumption **A.3**. □

In particular, $\mathbf{n}(W^-) = \mathbf{n}(W_1^-) + \mathbf{n}(W_2^-)$ is finite. We shall see that $\mathbf{n}(W^+) = \infty$.

4.2 Poisson point processes on W_a and a new process X

By Lemma 4.2, the excursion law \mathbf{n} is concentrated on the space W_a defined by (4.11). Accordingly, we consider the spaces

$$W_a^+ = \{w \in W^+ : \lim_{t \downarrow 0} w(t) = a\}, \quad W_a^- = \{w \in W^- : \lim_{t \downarrow 0} w(t) = a\},$$

so that $W_a = W_a^+ + W_a^-$. In the sequel however, we shall employ slightly modified but equivalent definitions of those spaces by extending each w from an S_0 -valued excursion to S -valued continuous one as follows:

$$W_a = \{w : \exists \zeta(w) \in (0, \infty], w \text{ is a continuous function from } [0, \zeta(w)) \text{ to } S, w(0) = a, \\ w(t) \in S_0, t \in (0, \zeta(w)), w(\zeta(w)-) \in \{a\} \cup \{\Delta\} \text{ if } \zeta(w) < \infty\}, \quad (4.13)$$

Any $w \in W_a$ for which $\zeta(w) < \infty, w(\zeta(w)-) = a$ will be regarded to be a continuous function from $[0, \zeta(w)]$ to S by setting $w(\zeta(w)) = a$. We further let

$$W_a^+ = \{w : \exists \zeta(w) \in (0, \infty), w \text{ is a continuous function from } [0, \zeta(w)] \text{ to } S, \\ w(t) \in S_0, t \in (0, \zeta(w)), w(0) = w(\zeta(w)) = a\}, \quad (4.14)$$

$$W_a^- = W_a \setminus W_a^+. \quad (4.15)$$

The excursion law \mathbf{n} will be considered to be a measure on W_a defined by (4.13) and we denote by \mathbf{n}^+ , \mathbf{n}^- , the restrictions of \mathbf{n} to W_a^+ , W_a^- defined by (4.14) and (4.15) respectively.

Let $\{\mathbf{p}_s, s > 0\}$ be a Poisson point process on W_a with characteristic measure \mathbf{n} defined on an appropriate probability space (Ω, P) . We then let

$$\mathbf{p}_s^+ = \begin{cases} \mathbf{p}_s & \text{if } \mathbf{p}_s \in W_a^+, \\ \partial & \text{otherwise,} \end{cases} \quad (4.16)$$

$$\mathbf{p}_s^- = \begin{cases} \mathbf{p}_s & \text{if } \mathbf{p}_s \in W_a^-, \\ \partial & \text{otherwise,} \end{cases} \quad (4.17)$$

where ∂ is an extra point disjoint of W_a . Then $\{\mathbf{p}_s^+, s > 0\}$, $\{\mathbf{p}_s^-, s > 0\}$ are mutually independent Poisson point processes on W_a^+ , W_a^- with characteristic measures \mathbf{n}^+ , \mathbf{n}^- respectively. Furthermore

$$\mathbf{p}_s = \mathbf{p}_s^+ + \mathbf{p}_s^-. \quad (4.18)$$

By means of the terminal time $\zeta(\mathbf{p}_r^+)$ of the excursion \mathbf{p}_r^+ , we let

$$J(s) = \sum_{r \leq s} \zeta(\mathbf{p}_r^+), \quad s > 0. \quad (4.19)$$

We put $J(0) = 0$.

Lemma 4.5. (i) $J(s) < \infty$ a.s. for $s > 0$.
(ii) $\{J(s)\}_{s \geq 0}$ is a subordinator with

$$E \left\{ e^{-\alpha J(s)} \right\} = \exp \left\{ -\alpha(u_\alpha, \varphi)s \right\}. \quad (4.20)$$

Proof. (i) We write $J(s)$ as $J(s) = I + II$ with

$$I = \sum_{r \leq s, \zeta(\mathbf{p}_r^+) \leq 1} \zeta(\mathbf{p}_r^+), \quad II = \sum_{r \leq s, \zeta(\mathbf{p}_r^+) > 1} \zeta(\mathbf{p}_r^+).$$

Since $\mathbf{n}^+(\zeta > 1) \leq \mu_1(S_0) < \infty$ by (4.2), r in the sum II is finite a.s. and hence $II < \infty$ a.s. On the other hand,

$$\begin{aligned} E(I) &= \mathbf{sn}^+(\zeta; \zeta \leq 1) \leq \mathbf{sn}^+(\zeta \wedge 1) \\ &= \mathbf{sn}^+ \left\{ \int_0^1 I_{(0, \zeta)}(t) dt \right\} = s \int_0^1 \mathbf{n}^+(\zeta > t) dt \leq s \int_0^1 \mu_t(S_0) dt, \end{aligned}$$

which is finite by (4.2). Hence $I < \infty$ a.s.

(ii) Clearly $\{J(s)\}_{s \geq 0}$ is increasing and of stationary independent increment. Since

$$e^{-\alpha J(s)} - 1 = \sum_{r \leq s} \left\{ e^{-\alpha J(r)} - e^{-\alpha J(r-)} \right\} = \sum_{r \leq s} e^{-\alpha J(r-)} \left\{ e^{-\alpha \zeta(\mathbf{p}_r^+)} - 1 \right\},$$

we have

$$E \left\{ e^{-\alpha J(s)} \right\} - 1 = -c \int_0^s E \left\{ e^{-\alpha J(r)} \right\} dr,$$

with

$$\begin{aligned} c &= \mathbf{n}^+(1 - e^{-\alpha \zeta}) = \mathbf{n}(1 - e^{-\alpha \zeta}; \zeta < \infty, w(\zeta) = a) \\ &= \mathbf{n} \left\{ \alpha \int_0^\zeta e^{-\alpha t} dt; \zeta < \infty, w(\zeta) = a \right\} \\ &= \alpha \int_0^\infty e^{-\alpha t} \mathbf{n}(t < \zeta < \infty, w(\zeta) = a) dt \\ &= \alpha \int_0^\infty e^{-\alpha t} \mu_t(\varphi) dt = \alpha \hat{\mu}_\alpha(\varphi) = \alpha(u_\alpha, \varphi) < \infty. \end{aligned}$$

□

In virtue of Lemma 4.3 and Lemma 4.5, we may assume that the next three properties hold for any $\omega \in \Omega$ by subtracting a P -negligible set from Ω if necessary:

$$J(s) < \infty \quad \forall s > 0, \quad (4.21)$$

$$\lim_{s \rightarrow \infty} J(s) = \infty, \quad (4.22)$$

and, for any finite interval $I \subset (0, \infty)$ and any neighbourhood U of a in S ,

$$\{s \in I : \tau_{U^c}(\mathbf{p}_s^+) < \zeta(\mathbf{p}_s^+)\} \text{ is a finite set.} \quad (4.23)$$

Let T be the time of occurrence of the first excursion of the point process $\{\mathbf{p}_s^-, s > 0\}$, namely,

$$T = \min\{s > 0 : \mathbf{p}_s^- \neq \partial\}. \quad (4.24)$$

Since $\mathbf{n}(W_a^-) = L(m_0, \psi) < \infty$ by Lemma 4.4, we can see that T and \mathbf{p}_T^- are independent and

$$P(T > t) = e^{-L(m_0, \psi)t}, \quad \text{the distribution of } \mathbf{p}_T^- = L(m_0, \psi)^{-1} \mathbf{n}^-. \quad (4.25)$$

We are now in a position to produce a new process $X = \{X_t\}_{t \geq 0}$ out of the point processes of excursions \mathbf{p}^\pm .

(i) For $0 \leq t < J(T-)$, we determine s by

$$J(s-) \leq t \leq J(s), \quad (4.26)$$

and let

$$X_t = \begin{cases} \mathbf{p}_s^+(t - J(s-)) & \text{if } J(s) - J(s-) > 0, \\ a & \text{if } J(s) - J(s-) = 0. \end{cases} \quad (4.27)$$

(ii) For $J(T-) \leq t < \zeta_\omega \equiv J(T-) + \zeta(\mathbf{p}_T^-)$, we let

$$X_t = \mathbf{p}_T^-(t - J(T-)). \quad (4.28)$$

In this way, the S -valued continuous path

$$X_t, \quad 0 \leq t < \zeta_\omega,$$

is defined and

$$X_{\zeta_\omega-} = \Delta \quad \text{if } \zeta_\omega < \infty.$$

Continuity of the path is a consequence of (4.23).

For this process $\{X_t, 0 \leq t < \zeta_\omega, P\}$, let us put

$$G_\alpha f(a) = E \left(\int_0^{\zeta_\omega} e^{-\alpha t} f(X_t) dt \right), \quad \alpha > 0, \quad f \in \mathcal{B}(S). \quad (4.29)$$

Proposition 4.2. *It holds that*

$$G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)}. \quad (4.30)$$

Proof. We use the notation

$$\hat{f}_\alpha(w) = \int_0^{\zeta(w)} e^{-\alpha t} f(w(t)) dt, \quad w \in W_a.$$

We have then

$$\begin{aligned} \int_0^{\zeta_\omega} e^{-\alpha t} f(X_t) dt &= \sum_{s < T} \int_{J(s^-)}^{J(s)} e^{-\alpha t} f(X_t) dt + \int_{J(T^-)}^{J(T^-) + \zeta(\mathbf{p}_T^-)} e^{-\alpha t} f(X_t) dt \\ &= \sum_{s < T} e^{-\alpha J(s^-)} \hat{f}_\alpha(\mathbf{p}_s^+) + e^{-\alpha J(T^-)} \hat{f}_\alpha(\mathbf{p}_T^-), \end{aligned}$$

and consequently

$$\begin{aligned} G_\alpha f(a) &= E \left(\sum_{s < T} e^{-\alpha J(s^-)} \hat{f}_\alpha(\mathbf{p}_s^+) + e^{-\alpha J(T^-)} \hat{f}_\alpha(\mathbf{p}_T^-) \right) \\ &= E \left(\int_0^T e^{-\alpha \hat{\mu}_\alpha(\varphi) s} ds \right) \mathbf{n}^+(\hat{f}_\alpha) + E \left(e^{-\alpha \hat{\mu}_\alpha(\varphi) T} \right) L(m_0, \psi)^{-1} \mathbf{n}^-(\hat{f}_\alpha) \\ &= \frac{\mathbf{n}^+(\hat{f}_\alpha)}{\alpha \hat{\mu}_\alpha(\varphi) + L(m_0, \psi)} + \frac{\mathbf{n}^-(\hat{f}_\alpha)}{\alpha \hat{\mu}_\alpha(\varphi) + L(m_0, \psi)} \\ &= \frac{\mathbf{n}(\hat{f}_\alpha)}{\alpha \hat{\mu}_\alpha(\varphi) + L(m_0, \psi)} = \frac{\hat{\mu}_\alpha(f)}{\alpha \hat{\mu}_\alpha(\varphi) + L(m_0, \psi)}. \end{aligned}$$

It then suffices to substitute (4.1) in the last expression. \square

4.3 Continuity of resolvent along X

Lemma 4.6. *For $\alpha > 0$ and $f \in \mathcal{B}(S)$, define $G_\alpha f(a)$ by the right hand side of (4.30) and extend it to a function on S by setting*

$$G_\alpha f(x) = G_\alpha^0 f(x) + G_\alpha f(a) u_\alpha(x), \quad x \in S_0. \quad (4.31)$$

Then $\{G_\alpha\}_{\alpha > 0}$ is an m -symmetric (sub)Markovian resolvent on S .

Proof. By making use of the resolvent equation for G_α^0 , the m -symmetry of G_α^0 and the equation

$$u_\alpha(x) - u_\beta(x) + (\alpha - \beta) G_\alpha^0 u_\beta(x) = 0, \quad \alpha, \beta > 0, \quad x \in S_0,$$

we can easily check the resolvent equation

$$G_\alpha f(x) - G_\beta f(x) + (\alpha - \beta) G_\alpha G_\beta f(x) = 0, \quad x \in S.$$

The m -symmetry of G_α

$$\int_S G_\alpha f(x) g(x) m(dx) = \int_S f(x) G_\alpha g(x) m(dx)$$

holding for any non-negative Borel functions f, g is clear. Moreover we get by Lemma 2.1 that

$$\begin{aligned}\alpha G_\alpha 1(x) &= \alpha G_\alpha^0 1(x) + u_\alpha(x) \frac{\alpha(u_\alpha, \varphi + \psi)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)} \\ &\leq 1 - u_\alpha(x) + u_\alpha(x) = 1, \quad x \in S_0,\end{aligned}$$

and similarly, $\alpha G_\alpha 1(a) \leq 1$. \square

Let $\{U_n\}$ be a decreasing sequence of open neighbourhoods of the point a in S such that $U_n \supset \overline{U_{n+1}}$ and $\bigcap_{n=1}^\infty U_n = \{a\}$. Let

$$A = A_{\alpha, \rho} = \{x \in S_0 : u_\alpha(x) < \rho\} \text{ for } \alpha > 0, 0 < \rho < 1.$$

We then set

$$\sigma_n = \inf\{t > 0 : X_t^0 \in U_n \cap S_0\}, \quad \sigma_a = \lim_{n \rightarrow \infty} \sigma_n, \quad \tau_n = \inf\{t > 0 : X_t^0 \in U_n \cap A\},$$

with the convention that $\inf \emptyset = \infty$.

Lemma 4.7. *For any $\alpha > 0$, $\rho \in (0, 1)$ and $x \in S_0$,*

$$\lim_{n \rightarrow \infty} P_x^0 \{\tau_n < \sigma_a < \infty\} = 0. \quad (4.32)$$

Proof. Since

$$\{\sigma_a < \infty\} = \{\zeta^0 < \infty, X_{\zeta^0-}^0 = a\}$$

and $\sigma_a = \zeta^0$ on the set $\{\sigma_a < \infty\}$, we have for $x \in S_0$ and $m < n$

$$\begin{aligned}u_\alpha(x) &= E_x^0 \{e^{-\alpha\sigma_a}; \tau_n < \sigma_a\} + E_x^0 \{e^{-\alpha\sigma_a}; \tau_n \geq \sigma_a\} \\ &= E_x^0 \{e^{-\alpha\tau_n} u_\alpha(X_{\tau_n}^0); \tau_n < \sigma_a\} + E_x^0 \{e^{-\alpha\sigma_a}; \tau_n \geq \sigma_a\} \\ &\leq \rho E_x^0 \{e^{-\alpha\tau_n}; \tau_n < \sigma_a\} + E_x^0 \{e^{-\alpha\sigma_a}; \tau_n \geq \sigma_a\} \\ &\leq \rho E_x^0 \{e^{-\alpha(\tau_n \wedge \sigma_a)}; \tau_m < \sigma_a\} + E_x^0 \{e^{-\alpha\sigma_a}; \tau_n \geq \sigma_a\}.\end{aligned}$$

By letting first $n \rightarrow \infty$ and then $m \rightarrow \infty$, we obtain

$$\begin{aligned}u_\alpha(x) &\leq \rho \lim_{m \rightarrow \infty} E_x^0 \{e^{-\alpha\sigma_a}; \tau_m < \sigma_a\} + \lim_{n \rightarrow \infty} E_x^0 \{e^{-\alpha\sigma_a}; \tau_n \geq \sigma_a\} \\ &= E_x^0 \{e^{-\alpha\sigma_a}\} - (1 - \rho) \lim_{n \rightarrow \infty} E_x^0 \{e^{-\alpha\sigma_a}; \tau_n < \sigma_a\} \\ &= u_\alpha(x) - (1 - \rho) \lim_{n \rightarrow \infty} E_x^0 \{e^{-\alpha\sigma_a}; \tau_n < \sigma_a\},\end{aligned}$$

which implies

$$\lim_{n \rightarrow \infty} E_x^0 \{e^{-\alpha\sigma_a}; \tau_n < \sigma_a\} = 0$$

and so (4.32) must hold. \square

Lemma 4.8. *Let $\alpha > 0$.*

(i) *For any $x \in S_0$,*

$$\lim_{t \uparrow \sigma_a} u_\alpha(X_t^0) = 1 \quad P_x^0\text{-a.s. on } \{\sigma_a < \infty\}. \quad (4.33)$$

(ii) $\mathbf{n}(\Lambda) = 0$ *where*

$$\Lambda = \left\{ w \in W_a^+ : \exists \alpha > 0, \lim_{t \uparrow \zeta} u_\alpha(w(t)) \neq 1 \right\}.$$

Proof. If $\sigma_a < \infty$ and if $\underline{\lim}_{t \uparrow \sigma_a} u_\alpha(X_t^0) < \rho$, then for any small $\epsilon > 0$ there exists $t \in (\sigma_a - \epsilon, \sigma_a)$ such that $u_\alpha(X_t^0) < \rho$, and so $\tau_n < \sigma_a$ for all n . Therefore by the preceding lemma

$$P_x^0 \{ \underline{\lim}_{t \uparrow \sigma_a} u_\alpha(X_t^0) < \rho, \sigma_a < \infty \} = 0.$$

Since u_α is decreasing in α and ρ can be taken arbitrarily close to 1, we obtain (4.33).

(ii) follows from (i) as

$$\begin{aligned} \mathbf{n}(\Lambda) &= \lim_{\epsilon \downarrow 0} \mathbf{n}(\Lambda \cap \{\epsilon < \zeta\}) \\ &= \lim_{\epsilon \downarrow 0} \int_{S_0} \mu_\epsilon(dx) P_x^0(\lim_{t \uparrow \sigma_a} u_\alpha(X_t^0) \neq 1) = 0. \end{aligned}$$

□

We extend u_α to a function on S by setting $u_\alpha(a) = 1$. By Lemma 4.8 (ii) combined with Lemma 4.1 and a similar reasoning as in the proof of Lemma 4.2, we may assume, subtracting a suitable \mathbf{n} -negligible set from W_a^+ (resp. W_a^-), that $u_1(w(t))$ is continuous in $t \in [0, \zeta]$ (resp. $t \in [0, \zeta)$.)

Lemma 4.9. *Let $0 < \rho < 1$ and set*

$$\tilde{W}_\rho = \left\{ w \in W_a^+ : \max_{0 \leq t \leq \zeta} \{1 - u_1(w(t))\} > \rho \right\}.$$

Then $\mathbf{n}^+(\tilde{W}_\rho) < \infty$.

Proof. The proof is similar to that of Lemma 4.3. For any x such that $1 - u_1(x) = \rho$ and for $\delta = -\log(1 - \frac{\rho}{2}) > 0$, we have

$$\begin{aligned} P_x^0(\sigma_a > \delta) &\geq E_x^0 \{1 - e^{-\sigma_a}; \sigma_a > \delta\} \\ &= E_x^0 \{1 - e^{-\sigma_a}\} - E_x^0 \{1 - e^{-\sigma_a}; \sigma_a \leq \delta\} \\ &\geq 1 - u_1(x) - (1 - e^{-\delta}) = \rho - (1 - e^{-\delta}) = \frac{\rho}{2}. \end{aligned}$$

Therefore if we set

$$A = \{x \in S_0 : 1 - u_1(x) \leq \rho\}, \quad \tau = \inf\{t > 0 : w(t) \in S_0 \setminus A\},$$

then

$$\begin{aligned} \mathbf{n}^+(\tilde{W}_\rho) &= \mathbf{n}^+(\tau < \zeta) = \lim_{\epsilon \downarrow 0} \mathbf{n}^+(\epsilon < \tau < \zeta^0) = \lim_{\epsilon \downarrow 0} \int_A \mu_\epsilon(dx) P_x^0(\tau < \zeta^0) \\ &\leq \overline{\lim}_{\epsilon \downarrow 0} \int_A \mu_\epsilon(dx) E_x^0 \left\{ \left(\frac{2}{\rho} \right) P_{X_\tau^0}^0(\sigma_a > \delta); \tau < \zeta^0 \right\} \\ &\leq \frac{2}{\rho} \overline{\lim}_{\epsilon \downarrow 0} \int_{S_0} \mu_\epsilon(dx) P_x^0(\sigma_a > \delta) \\ &\leq \frac{2}{\rho} \lim_{\epsilon \downarrow 0} \int_{S_0} \mu_\epsilon(dx) P_x^0(\zeta^0 > \delta) + \frac{2}{\rho} \lim_{\epsilon \downarrow 0} \int_{S_0} \mu_\epsilon(dx) P_x^0(\zeta^0 < \sigma_a = \infty) \\ &= \frac{2}{\rho} \lim_{\epsilon \downarrow 0} \mu_{\epsilon+\delta}(1) + \frac{2}{\rho} \lim_{\epsilon \downarrow 0} \mu_\epsilon(\psi^{(1)}), \end{aligned}$$

which is finite in view of (4.2) and Lemma 4.4. \square

For $\alpha > 0$, $f \in \mathcal{B}(S)$, we defined the resolvent $G_\alpha f$ by

$$G_\alpha f(x) = G_\alpha^0 f(x) + G_\alpha f(a)u_\alpha(x), \quad x \in S_0$$

with $G_\alpha f(a)$ of Proposition 4.2. We now extend $G_\alpha^0 f(x)$ to S by setting

$$G_\alpha^0 f(a) = 0.$$

In the last subsection, we have constructed a process $\{X_t\}_{t \in [0, \zeta_\omega]}$ out of the Poisson point processes \mathbf{p}^+ , \mathbf{p}^- on W_a^+ , W_a^- defined on a probability space (Ω, P) .

Proposition 4.3. *Let $u = G_\alpha f$ with $f \in C_b(S)$. Then $u(X_t)$ is continuous in $t \in [0, \zeta_\omega)$, P -a.s.*

Proof. As was remarked immediately after the proof of Lemma 4.8, u_1 is continuous along any sample point functions of $\mathbf{p}^+ = \{\mathbf{p}_s^+, s > 0\}$ and $\mathbf{p}^- = \{\mathbf{p}_s^-, s > 0\}$. Moreover, by Lemma 4.9, we can subtract a suitable P -negligible set from Ω so that, in addition to the properties (4.21),(4.22) and (4.23), \mathbf{p}^+ satisfies the following property for every sample point $\omega \in \Omega$: for any finite interval $I \subset (0, \infty)$ and for any $\rho \in (0, 1)$,

$$\{s \in I : \max_{0 \leq t \leq \zeta(\mathbf{p}_s^+)} (1 - u_1(\mathbf{p}_s^+(t))) > \rho\} \text{ is a finite set.} \quad (4.34)$$

Then it is not hard to see that not only X_t but also $u_1(X_t)$ are continuous in $t \in [0, \zeta_\omega)$. From the inequality $G_1^0 1(x) \leq 1 - u_1(x)$, $x \in S$, we see that

$$\lim_{t \rightarrow t_0} G_1^0 1(X_t) = 0 \quad \text{if } X_{t_0} = a.$$

Hence $G_1^0 f(X_t)$ has the same property as the above for $f \in C_b(S)$. Since $G_1^0 f(X_t)$ is clearly continuous on $\{t \in [0, \zeta_\omega) : X_t \neq a\}$ by the assumption **A.4**, it is continuous on $[0, \zeta_\omega)$. We have thus proved the continuity of $G_1 f(X_t)$. The continuity of $G_\alpha f(X_t)$ follows from the resolvent equation proved in Lemma 4.6. \square

4.4 Markov property of X

Let us define $p_t f(x)$ for $t > 0, x \in S, f \in \mathcal{B}(S)$, as follows:

$$p_t f(a) = E(f(X_t); \zeta_\omega > t), \quad (4.35)$$

$$p_t f(x) = p_t^0 f(x) + E_x \{p_{t-\sigma_a} f(a); \sigma_a \leq t\}, \quad x \in S_0. \quad (4.36)$$

Evidently

$$\int_0^\infty e^{-\alpha t} p_t f dt = G_\alpha f, \quad \alpha > 0. \quad (4.37)$$

Lemma 4.10. $p_{t+s} = p_t p_s, \quad t, s > 0.$

Proof. Take any $f \in C_b(S)$. By (4.36) and the resolvent equation in Lemma 4.6, we have for any $x \in S$

$$\int_0^\infty e^{-\alpha t} \left\{ \int_0^\infty e^{-\beta s} p_{t+s} f(x) ds \right\} dt = \int_0^\infty e^{-\alpha t} \{p_t(G_\beta f)(x)\} dt, \quad (4.38)$$

because the left hand side equals $\frac{1}{\alpha - \beta}(G_\beta f(x) - G_\alpha f(x)) = G_\alpha G_\beta f(x)$.

We first consider the case where $x = a$. Then the functions inside $\{\cdot\}$ of the both hand sides of (4.38) are continuous in $t > 0$ in virtue of the continuity of X and Proposition 4.3. Hence we have for any $t > 0$

$$\int_0^\infty e^{-\beta s} p_{t+s} f(a) ds = p_s(G_\beta f)(a) = \int_0^\infty e^{-\beta s} p_t(p_s f)(a) ds.$$

Since both $p_{t+s} f(a)$, $p_t(p_s f)(a)$ are right continuous in $s > 0$, we get

$$p_{t+s} f(a) = p_t(p_s f)(a), \quad t > 0, s > 0. \quad (4.39)$$

We next consider the case where $x \in S_0$. Using (4.37), we obtain

$$\begin{aligned} p_{t+s} f(x) &= p_{t+s}^0 f(x) + E_x^0 \{p_{t+s-\sigma_a} f(a); \sigma_a \leq t+s\} \\ &= p_{t+s}^0 f(x) + E_x^0 \{p_{t-\sigma_a}(p_s f)(a); \sigma_a \leq t\} \\ &\quad + E_x^0 \{p_{t+s-\sigma_a} f(a) : t < \sigma_a \leq t+s\}. \end{aligned}$$

On the other hand,

$$p_t(p_s f)(x) = p_t^0(p_s f)(x) + E_x^0 \{p_{t-\sigma_a}(p_s f)(a); \sigma_a \leq t\}.$$

Hence it suffices to prove that

$$p_{t+s}^0 f(x) + E_x^0 \{p_{t+s-\sigma_a} f(a); t < \sigma_a \leq t+s\} = p_t^0(p_s f)(x). \quad (4.40)$$

Put

$$g(x) = E_x^0 \{p_{s-\sigma_a} f(a); \sigma_a \leq s\},$$

then, we are led from $p_s f(x) = p_s^0 f(x) + g(x)$ to

$$p_t^0(p_s f)(x) = p_{t+s}^0 f(x) + p_t^0 g(x),$$

and consequently, (4.40) is reduced to

$$E_x^0 \{p_{t+s-\sigma_a} f(a); t < \sigma_a \leq t+s\} = E_x^0(g(X_t^0); \zeta^0 > t). \quad (4.41)$$

With the notation θ_t to denote the usual shift, the left hand side of (4.41) equals

$$\begin{aligned} &E_x^0 \{p_{t+s-\sigma_a} f(a); \zeta^0 > t, \sigma_a > t, \sigma_a \circ \theta_t \leq s\} \\ &= E_x^0 \{p_{s-\sigma_a \circ \theta_t} f(a); \zeta^0 > t, \sigma_a \circ \theta_t \leq s\} \\ &= E_x^0 \left[E_{X_t^0}^0 \{p_{s-\sigma_a} f(a); \sigma_a \leq s\}; \zeta^0 > t \right], \end{aligned}$$

which coincides with the right hand side of (4.41) as was to be proved. \square

Lemma 4.11. *Suppose $g \in \mathcal{B}(S)$ and $\lim_{\epsilon \downarrow 0} p_\epsilon g(x) = g(x)$, $x \in S$. Then, for any $f \in C_b(S)$, $t > 0$,*

$$\lim_{\epsilon \downarrow 0} p_\epsilon(f p_t g)(x) = f(x) p_t g(x), \quad x \in S. \quad (4.42)$$

Proof. Fix $x \in S$. Clearly, for any neighbourhood U of x ,

$$\lim_{\epsilon \downarrow 0} p_\epsilon I_U(x) = 1,$$

and hence

$$p_\epsilon |f p_t g|(x) = p_\epsilon |f I_U p_t g|(x) + o(\epsilon).$$

For any $\delta > 0$, take a neighbourhood U of x such that

$$|f(y) - f(x)| < \delta, \quad y \in U.$$

Then

$$\begin{aligned} & |p_\epsilon (f p_t g)(x) - f(x) p_\epsilon (p_t g)(x)| \\ & \leq p_\epsilon (|f - f(x)| |p_t g|)(x) \\ & \leq p_\epsilon (|f - f(x)| I_U |p_t g|)(x) + o(\epsilon) \leq \delta \|g\|_\infty + o(\epsilon). \end{aligned}$$

On the other hand, we have from the preceding lemma that

$$\lim_{\epsilon \downarrow 0} f(x) p_\epsilon (p_t g)(x) = \lim_{\epsilon \downarrow 0} f(x) p_t (p_\epsilon g)(x) = f(x) p_t g(x).$$

Consequently

$$\overline{\lim}_{\epsilon \downarrow 0} |p_\epsilon (f p_t g)(x) - f(x) p_t g(x)| \leq \delta \|g\|_\infty,$$

which means (4.42) because $\delta > 0$ can be taken arbitrarily small. \square

Proposition 4.4. (i) For $\alpha_1, \dots, \alpha_n > 0$,

$$E \left\{ \int_{0 < t_1 < \dots < t_n < \zeta_\omega} \prod_{k=1}^n \left(e^{-\alpha_k (t_k - t_{k-1})} f_k(X_{t_k}) \right) dt_1 \dots dt_n \right\} = G_{\alpha_1} f_1 G_{\alpha_2} f_2 \dots G_{\alpha_n} f_n(a), \quad (4.43)$$

where we set $t_0 = 0$ by convention.

(ii). $X = \{X_t, 0 \leq t < \zeta_\omega, P\}$ is a Markov process on S with transition function p_t and initial distribution concentrated at $\{a\}$.

Proof. We shall employ the following notations:

$$F(X; t; \alpha_1, f_1, \dots, \alpha_n, f_n) = \int_{t < t_1 < \dots < t_n < \zeta_\omega} \prod_{k=1}^n \left\{ e^{-\alpha_k (t_k - t_{k-1})} f_k(X_{t_k}) \right\} dt_1 \dots dt_n,$$

and, for $w \in W_a$,

$$F(w; t; \alpha_1, f_1, \dots, \alpha_n, f_n) = \int_{t < t_1 < \dots < t_n < \zeta(w)} \prod_{k=1}^n \left\{ e^{-\alpha_k (t_k - t_{k-1})} f_k(w(t_k)) \right\} dt_1 \dots dt_n.$$

(i). The left hand side of (4.43) will be denoted by $G(\alpha_1, f_1, \dots, \alpha_n, f_n)$, namely,

$$E \{ F(X; 0; \alpha_1, f_1, \dots, \alpha_n, f_n) \} = G(\alpha_1, f_1, \dots, \alpha_n, f_n). \quad (4.44)$$

For $0 < s < T$, we denote by $I(s)$ the expression

$$\int_{J(s^-) < t_1 < J(s)} e^{-\alpha_1 t_1} f_1(X_{t_1}) \left\{ \int \cdots \int_{t_1 < t_2 < \cdots < t_n < \zeta_\omega} \prod_{k=2}^n \left(e^{-\alpha_k (t_k - t_{k-1})} f_k(X_{t_k}) \right) dt_2 \cdots dt_n \right\} dt_1.$$

Then

$$F(X; 0; \alpha_1, f_1, \cdots, \alpha_n, f_n) = \sum_{0 < s < T} I(s) + F(X; J(T^-); \alpha_1, f_1, \cdots, \alpha_n, f_n).$$

Further, if we put for $1 \leq m \leq n$

$$\begin{aligned} I_m(s) &= \int_{J(s^-) < t_1 < \cdots < t_m < J(s)} \prod_{k=1}^m \left\{ e^{-\alpha_k (t_k - t_{k-1})} f_k(X_{t_k}) \right\} dt_1 \cdots dt_m \\ &\cdot \int_{J(s) < t_{m+1} < \cdots < t_n < \zeta_\omega} \prod_{\ell=m+1}^n \left\{ e^{-\alpha_\ell (t_\ell - t_{\ell-1})} f_\ell(X_{t_\ell}) \right\} dt_{m+1} \cdots dt_n, \end{aligned}$$

then

$$I(s) = \sum_{m=1}^n I_m(s).$$

Moreover, each $I_m(s)$ can be written as

$$I_m(s) = F_m(s) G_m(s)$$

with

$$\begin{aligned} F_m(s) &= \int_{J(s^-) < t_1 < \cdots < t_m < J(s)} \prod_{k=1}^m \left\{ e^{-\alpha_k (t_k - t_{k-1})} f_k(X_{t_k}) \right\} e^{-\alpha_{m+1} (J(s) - t_m)} dt_1 \cdots dt_m, \\ G_m(s) &= \int_{J(s) < t_{m+1} < \cdots < t_n < \zeta_\omega} e^{-\alpha_{m+1} (t_{m+1} - J(s))} \prod_{\ell=m+2}^n \left\{ e^{-\alpha_\ell (t_\ell - t_{\ell-1})} f_\ell(X_{t_\ell}) \right\} dt_{m+1} \cdots dt_n. \end{aligned}$$

Therefore

$$F(X; 0; \alpha_1, f_1, \cdots, \alpha_n, f_n) = \sum_{0 < s < T} \sum_{m=1}^n F_m(s) G_m(s) + F(X; J(T^-); \alpha_1, f_1, \cdots, \alpha_n, f_n). \quad (4.45)$$

Next, let us put (with the convention that $\alpha_{n+1} = 0$)

$$\begin{aligned} &F(w; \alpha_1, f_1, \cdots, \alpha_m, f_m; \alpha_{m+1}) \\ &= \int_{0 < t_1 < \cdots < t_m < \zeta(w)} \prod_{k=1}^m \left\{ e^{-\alpha_k (t_k - t_{k-1})} f_k(w(t_k)) \right\} e^{-\alpha_{m+1} (\zeta(w) - t_m)} dt_1 \cdots dt_m, \end{aligned} \quad (4.46)$$

so that

$$F_m(s) = e^{-\alpha_1 J(s^-)} F(\mathbf{p}_s^+; \alpha_1, f_1, \cdots, \alpha_m, f_m; \alpha_{m+1}). \quad (4.47)$$

We furthermore put $Y_t = X_{J(s)+t}$ so that

$$G_m(s) = \int_{0 < t_{m+1} < \dots < t_n < \zeta_{\omega - J(s)}} \dots \int \prod_{\ell=m+1}^n \left\{ e^{-\alpha_\ell(t_\ell - t_{\ell-1})} f_\ell(X_{t_\ell}) \right\} dt_{m+1} \dots dt_n, \quad (4.48)$$

where we set $t_m = 0$.

For $\mathbf{p} = \{\mathbf{p}_t, t > 0\}$, we may use the following notations:

$$\begin{aligned} & G(\mathbf{p}; \alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n) \\ &= \int_{0 < t_{m+1} < \dots < t_n < \zeta_\omega} \dots \int \prod_{\ell=m+1}^n \left\{ e^{-\alpha_\ell(t_\ell - t_{\ell-1})} f_\ell(X_{t_\ell}) \right\} dt_{m+1} \dots dt_n, \end{aligned} \quad (4.49)$$

(with the convention that $t_m = 0$), and

$$\theta_s \mathbf{p} = \{\mathbf{p}_{s+t}, t > 0\}. \quad (4.50)$$

$\theta_s \mathbf{p}$ then has the same distribution as \mathbf{p} and independent of $\{\mathbf{p}_t, 0 < t < s\}$. Since Y_t is constructed from $\theta_s \mathbf{p}$ in the same way as X_t is from \mathbf{p} , (4.48) can be rewritten as

$$G_m(s) = G(\theta_s \mathbf{p}; \alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n), \quad (4.51)$$

which is identical in law to

$$G(\mathbf{p}; \alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n)$$

for each fixed $s > 0$. Further

$$F(X; J(T-); \alpha_1, f_1, \dots, \alpha_n, f_n) = e^{-\alpha_1 J(T-)} F(\mathbf{p}_T^-; 0; \alpha_1, f_1, \dots, \alpha_n, f_n). \quad (4.52)$$

Combining (4.45),(4.47),(4.51) and (4.52), we arrive at

$$\begin{aligned} & F(X; 0; \alpha_1, f_1, \dots, \alpha_n, f_n) \\ &= \sum_{0 < s < T} \sum_{m=1}^n e^{-\alpha_1 J(s-)} F(\mathbf{p}_s^+; \alpha_1, f_1, \dots, \alpha_m, f_m; \alpha_{m+1}) \\ & \quad \cdot G(\theta_s \mathbf{p}; \alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n) + e^{-\alpha_1 J(T-)} F(\mathbf{p}_T^-; 0; \alpha_1, f_1, \dots, \alpha_n, f_n) \end{aligned} \quad (4.53)$$

Here we compute the expectations of the random variables appearing in the last formula.

$$\mathbf{n}^+ \{F(w; \alpha_1, f_1, \dots, \alpha_m, f_m; \alpha_{m+1})\} = \hat{\mu}_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \dots G_{\alpha_{m-1}}^0 f_{m-1} G_{\alpha_m}^0 f_m u_{\alpha_{m+1}}). \quad (4.54)$$

When $m = n$, the last factor $u_{\alpha_{n+1}}$ in the above expression is understood to be $u_0 = \varphi$. In fact, the left hand side equals

$$\begin{aligned} & \mathbf{n} \left\{ \int_{0 < t_1 < \dots < t_m < \zeta(w)} \dots \int \prod_{k=1}^m \left(e^{-\alpha_k(t_k - t_{k-1})} f_k(w(t_k)) \right) e^{-\alpha_{m+1}(\zeta(w) - t_m)} dt_1 \dots dt_m; W_a^+ \right\} \\ &= \int_{0 < t_1 < \dots < t_m < \infty} \dots \int \mathbf{n} \left\{ \prod_{k=1}^m \left(e^{-\alpha_k(t_k - t_{k-1})} f_k(w(t_k)) \right) u_{\alpha_{m+1}}(w(t_m)); \zeta > t_m \right\}, \end{aligned}$$

which can be seen to coincide with the right hand side of (4.54) by (4.4).

We further have for any constant time $s > 0$,

$$E \{G(\theta_s \mathbf{p}; \alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n)\} = G(\alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n). \quad (4.55)$$

On the other hand, we have in view of §4.2

$$\begin{aligned} E \{F(\mathbf{p}_T^-; 0; \alpha_1, f_1, \dots, \alpha_n, f_n)\} &= L(m_0, \psi)^{-1} \mathbf{n}^- \{F(w; 0; \alpha_1, f_1, \dots, \alpha_n, f_n)\} \\ &= L(m_0, \psi)^{-1} \hat{\mu}_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \cdots G_{\alpha_{n-1}}^0 f_{n-1} G_{\alpha_n}^0 f_n \psi), \\ E \left\{ \int_0^T e^{-\alpha_1 J(s)} ds \right\} &= \frac{1}{\alpha_1(u_{\alpha_1}, \varphi) + L(m_0, \psi)}, \end{aligned} \quad (4.56)$$

$$E \left\{ e^{-\alpha_1 J(T-)} \right\} = \frac{L(m_0, \psi)}{\alpha_1(u_{\alpha_1}, \varphi) + L(m_0, \psi)}. \quad (4.57)$$

We can now get from (4.53) that

$$\begin{aligned} G(\alpha_1, f_1, \dots, \alpha_n, f_n) &= E \{F(X; 0; \alpha_1, f_1, \dots, \alpha_n, f_n)\} \\ &= \sum_{m=1}^n E \left\{ \int_0^T e^{-\alpha_1 J(s)} ds \right\} \mathbf{n}^+ \{F(w; \alpha_1, f_1, \dots, \alpha_m; \alpha_{m+1})\} \\ &\quad \times G(\alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n) + E \left\{ e^{-\alpha_1 J(T-)} \right\} E \{F(\mathbf{p}^-; 0; \alpha_1, f_1, \dots, \alpha_n, f_n)\} \\ &= \sum_{m=1}^{n-1} \frac{1}{\alpha_1(u_{\alpha_1}, \varphi) + L(m_0, \psi)} \hat{\mu}_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \cdots G_{\alpha_{m-1}}^0 f_{m-1} G_{\alpha_m}^0 f_m u_{\alpha_{m+1}}) \\ &\quad \times G(\alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n) + \frac{1}{\alpha_1(u_{\alpha_1}, \varphi) + L(m_0, \psi)} \hat{\mu}_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \cdots G_{\alpha_{n-1}}^0 f_{n-1} G_{\alpha_n}^0 f_n \varphi) \\ &+ \frac{L(m_0, \psi)}{\alpha_1(u_{\alpha_1}, \varphi) + L(m_0, \psi)} L(m_0, \psi)^{-1} \hat{\mu}_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \cdots G_{\alpha_{n-1}}^0 f_{n-1} G_{\alpha_n}^0 f_n \psi) \\ &= \frac{1}{\alpha_1(u_{\alpha_1}, \varphi) + L(m_0, \psi)} \sum_{m=1}^n \hat{\mu}_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \cdots G_{\alpha_{m-1}}^0 f_{m-1} G_{\alpha_m}^0 f_m u_{\alpha_{m+1}}) \\ &\quad \cdot G(\alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n). \end{aligned}$$

In the above and in what follows, we use the convention that

$$u_{\alpha_{m+1}} = G(\alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n) = 1$$

for $m = n$. This combined with (4.1) and (4.30) eventually leads us to

$$\begin{aligned} G(\alpha_1, f_1, \dots, \alpha_n, f_n) &= \sum_{m=1}^n G_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \cdots G_{\alpha_{m-1}}^0 f_{m-1} G_{\alpha_m}^0 f_m u_{\alpha_{m+1}})(a) \\ &\quad \cdot G(\alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n). \end{aligned} \quad (4.58)$$

Based on this formula, we shall prove the desired identity (4.43), namely,

$$G(\alpha_1, f_1, \dots, \alpha_n, f_n) = G_{\alpha_1} f_1 G_{\alpha_2} f_2 \cdots G_{\alpha_n} f_n(a) \quad (4.59)$$

by induction in n .

(1). When $n = 1$, (4.59) is just (4.30).

(2). Suppose (4.59) holds up to $n - 1$. Then

$$G(\alpha_{m+1}, f_{m+1}, \dots, \alpha_n, f_n) = (G_{\alpha_{m+1}} f_{m+1} \cdots G_{\alpha_n} f_n)(a),$$

and (4.58) can be written as

$$G(\alpha_1, f_1, \dots, \alpha_n, f_n) = \sum_{m=1}^n G_{\alpha_1}(f_1 G_{\alpha_2}^0 f_2 \cdots G_{\alpha_{m-1}}^0 f_{m-1} G_{\alpha_m}^0 f_m u_{\alpha_{m+1}})(a) \cdot (G_{\alpha_{m+1}} f_{m+1} \cdots G_{\alpha_n} f_n)(a). \quad (4.60)$$

Let us rewrite the right hand side of (4.59) by applying the formula (4.31) to the operation G_{α_2} in getting

$$\begin{aligned} (G_{\alpha_1} f_1 G_{\alpha_2} f_2 \cdots G_{\alpha_n} f_n)(a) &= (G_{\alpha_1} f_1 G_{\alpha_2}^0 f_2 G_{\alpha_3} f_3 \cdots G_{\alpha_n} f_n)(a) \\ &+ (G_{\alpha_1} f_1 u_{\alpha_2})(a) (G_{\alpha_2} f_2 \cdots G_{\alpha_n} f_n)(a). \end{aligned}$$

Apply the same procedure to the operation G_{α_3} to see that the right hand side of (4.59) equals

$$\begin{aligned} &(G_{\alpha_1} f_1 G_{\alpha_2}^0 f_2 G_{\alpha_3}^0 f_3 G_{\alpha_4} f_4 \cdots G_{\alpha_n} f_n)(a) \\ &+ (G_{\alpha_1} f_1 G_{\alpha_2}^0 f_2 u_{\alpha_3})(a) (G_{\alpha_3} f_3 \cdots G_{\alpha_n} f_n)(a) \\ &+ (G_{\alpha_1} f_1 u_{\alpha_2})(a) (G_{\alpha_2} f_2 \cdots G_{\alpha_n} f_n)(a). \end{aligned}$$

Repeating the same procedures, we finally find that the right hand side of (4.59) coincides with the right hand side of (4.60) as was to be proved.

(ii). For $t_1 > 0, \dots, t_n > 0$, let

$$F(t_1, \dots, t_n) = E \left\{ \prod_{k=1}^n f_k(X_{t_1+\dots+t_k}); \zeta_\omega > t_1 + \dots + t_n \right\},$$

$$G(t_1, \dots, t_n) = (p_{t_1} f_1 p_{t_2} f_2 \cdots p_{t_n} f_n)(a).$$

(4.43) is then equivalent to

$$\begin{aligned} &\int_0^\infty \cdots \int_0^\infty e^{-\alpha_1 t_1 - \cdots - \alpha_n t_n} F(t_1, \dots, t_n) dt_1 \cdots dt_n \\ &= \int_0^\infty \cdots \int_0^\infty e^{-\alpha_1 t_1 - \cdots - \alpha_n t_n} G(t_1, \dots, t_n) dt_1 \cdots dt_n. \end{aligned} \quad (4.61)$$

Clearly $F(t_1, \dots, t_n)$ is right continuous. Further, by virtue of Lemma 4.11, we can easily see that $G(t_1, \dots, t_n)$ is separately right continuous. Consequently, (4.61) implies

$$F(t_1, \dots, t_n) = G(t_1, \dots, t_n)$$

the desired Markov property of X . □

We add a lemma saying that the point a is regular for itself with respect to (X_t, P) .

Lemma 4.12. (i) $P(\eta_a = 0) = 1$, where $\eta_a = \inf\{t > 0 : X_t = a\}$.
(ii) $\mathbf{n}^+(W_a) = \infty$.

Proof. (i). In view of the proof of Proposition 4.3, $\lim_{t \downarrow 0} u_1(X_t) = 1$. Hence, if we put $\eta_{a,\epsilon} = \inf\{t > \epsilon : X_t = a\}$, then owing to the Markov property

$$\begin{aligned} E(e^{-\eta_a}) &= \lim_{\epsilon \downarrow 0} E(e^{-\eta_{a,\epsilon}}) \\ &= \lim_{\epsilon \downarrow 0} E(e^{-\epsilon} u_1(X_\epsilon); \zeta_\omega > \epsilon) = 1. \end{aligned}$$

(ii). By the construction of X_t , the point a is evidently instantaneous in the sense that

$$P(\tau_a = 0) = 1, \quad \text{where } \tau_a = \inf\{t > 0 : X_t \in S_0\}.$$

Hence (i) holds if and only if the domain $D_{\mathbf{p}^+}$ of the Poisson point process \mathbf{p}^+ accumulates at 0 P -a.s., which is also equivalent to (ii) (cf. [15, §4]). \square

4.5 A symmetric extension \tilde{X} of X^0

In §4.1, we have started with an m -symmetric diffusion

$$X^0 = \{X_t^0, 0 \leq t < \zeta^0, P_x^0, x \in S_0\}$$

on S_0 , where P_x^0 , $x \in S_0$, are probability measures on a certain sample space, say Ω^0 .

In §4.2, we have constructed a continuous process

$$X = \{X_t, 0 \leq t < \zeta_\omega, P\}$$

on S by piecing together the excursions, where P is a probability measure on another sample space Ω to define the excursion valued Poisson point processes.

For convenience, we assume that Ω^0 contains an extra point ω^a with $P_x^0(\{\omega^a\}) = 0$, $x \in S_0$, and we set $P_a^0 = \delta_{\omega^a}$, ω^a representing a path taking value a at any time.

We now let

$$\tilde{\Omega} = \Omega^0 \times \Omega, \quad \tilde{P}_x = P_x^0 \times P, \quad x \in S. \quad (4.62)$$

For $\tilde{\omega} = (\omega^0, \omega) \in \tilde{\Omega}$, let us define $\tilde{X}_t = \tilde{X}_t(\tilde{\omega})$ as follows:

(1) When $\omega^0 \in \Omega^0 \setminus \{\omega^a\}$,

$$\tilde{X}_t(\tilde{\omega}) = \begin{cases} X_t^0(\omega^0) & 0 \leq t < \zeta^0(\omega^0) \leq \sigma_a(\omega^0) \leq \infty \\ X_{t-\sigma_a(\omega^0)}(\omega) & \sigma_a(\omega^0) \leq t < \sigma_a(\omega^0) + \zeta_\omega, \text{ if } \sigma_a(\omega^0) < \infty. \end{cases} \quad (4.63)$$

(2) When $\omega^0 = \omega^a$,

$$\tilde{X}_t(\tilde{\omega}) = X_t(\omega) \quad 0 \leq t < \zeta_\omega. \quad (4.64)$$

The life time $\tilde{\zeta}$ of \tilde{X}_t is defined by

$$\tilde{\zeta} = \begin{cases} \zeta^0 & \text{if } \sigma_a(\omega^0) = \infty, \\ \sigma_a(\omega^0) + \zeta_\omega & \text{if } \sigma_a(\omega^0) < \infty. \end{cases} \quad (4.65)$$

Lemma 4.13. $\tilde{X} = \{\tilde{X}_t, 0 \leq t < \tilde{\zeta}, \tilde{P}_x, x \in S\}$ is a Markov process on S with transition function $\{p_t\}$ defined by (4.35) and (4.36).

Proof. This is an easy consequence of the Markov property of (X_t^0, P_x^0) and the Markov property of (X_t, P) proved in Proposition 4.4. To see this, we put, for any $0 < s_1 < s_2 < \dots < s_n$, $f_1, f_2, \dots, f_n \in \mathcal{B}(S)$,

$$I_k = \tilde{E}_x \left(f_1(\tilde{X}_{s_1}) \cdots f_{k-1}(\tilde{X}_{s_{k-1}}) f_k(\tilde{X}_{s_k}) \cdots f_n(\tilde{X}_{s_n}); s_{k-1} < \sigma_a \leq s_k \right),$$

for $1 \leq k \leq n$ with $s_0 = 0$, and

$$J = \tilde{E}_x (f_1(\tilde{X}_{s_1}) \cdots f_n(\tilde{X}_{s_n}); s_n < \sigma_a).$$

Using the definition of \tilde{X} , Proposition 4.4, the Markov property of X^0 and (4.36) successively, we are led to

$$\begin{aligned} I_k &= E_x^0 \left(f_1(X_{s_1}^0) \cdots f_{k-1}(X_{s_{k-1}}^0) E(f_k(X_{s_k - \sigma_a}) \cdots f_n(X_{s_n - \sigma_a})); s_{k-1} < \sigma_a \leq s_k \right) \\ &= E_x^0 \left(f_1(X_{s_1}^0) \cdots f_{k-1}(X_{s_{k-1}}^0) p_{s_k - \sigma_a} (f_k p_{s_{k+1} - s_k} f_{k+1} \cdots p_{s_n - s_{n-1}} f_n)(a); s_{k-1} < \sigma_a \leq s_k \right) \\ &= E_x^0 \left\{ f_1(X_{s_1}^0) \cdots f_{k-1}(X_{s_{k-1}}^0) \right. \\ &\quad \cdot E_{X_{s_{k-1}}^0}^0 \left(p_{s_k - s_{k-1} - \sigma_a} (f_k p_{s_{k+1} - s_k} f_{k+1} \cdots p_{s_n - s_{n-1}} f_n); \sigma_a \leq s_k - s_{k-1} \right); s_{k-1} < \sigma_a \leq s_k \left. \right\} \\ &= E_x^0 \left(f_1(X_{s_1}^0) \cdots f_{k-1}(X_{s_{k-1}}^0) \right. \\ &\quad \cdot (p_{s_k - s_{k-1}} - p_{s_k - s_{k-1}}^0) (f_k p_{s_{k+1} - s_k} f_{k+1} \cdots p_{s_n - s_{n-1}} f_n)(X_{s_{k-1}}^0); s_{k-1} < \sigma_a \leq s_k \left. \right). \end{aligned}$$

By the Markov property of X^0 , we thus get

$$\begin{aligned} I_k &= p_{s_1}^0 f_1 \cdots p_{s_{k-1} - s_{k-2}}^0 f_{k-1} p_{s_k - s_{k-1}} f_k p_{s_{k+1} - s_k} f_{k+1} \cdots p_{s_n - s_{n-1}} f_n(x) \\ &\quad - p_{s_1}^0 f_1 \cdots p_{s_{k-1} - s_{k-2}}^0 f_{k-1} p_{s_k - s_{k-1}}^0 f_k p_{s_{k+1} - s_k} f_{k+1} \cdots p_{s_n - s_{n-1}} f_n(x). \end{aligned}$$

Clearly we also have

$$J = E_x^0 (f_1(X_{s_1}^0) \cdots f_n(X_{s_n}^0); s_n < \sigma_a) = p_{s_1}^0 f_1 \cdots p_{s_n - s_{n-1}}^0 f_n.$$

Hence we arrive at

$$\tilde{E}_x (f_1(\tilde{X}_{s_1}) f_2(\tilde{X}_{s_2}) \cdots f_n(\tilde{X}_{s_n})) = \sum_{k=1}^n I_k + J = p_{s_1} f_1 p_{s_2 - s_1} f_2 \cdots p_{s_n - s_{n-1}} f_n(x),$$

the desired Markov property of \tilde{X} . □

We now state main theorems of the present paper. In this section, we have started with an m -symmetric diffusion X^0 on S_0 satisfying conditions **A.1, A.2, A.3, A.4** and constructed a Markov process \tilde{X} on S . The resolvent $\{G_\alpha\}_{\alpha > 0}$ of the Markov process \tilde{X} is defined by

$$G_\alpha f(x) = \tilde{E}_x \left(\int_0^\infty e^{-\alpha t} f(\tilde{X}_t) dt \right), \quad f \in \mathcal{B}(S). \quad (4.66)$$

The resolvent of X^0 was denoted by G_α^0 .

Theorem 4.1. *The process \tilde{X} enjoys the following properties:*

(1) \tilde{X} is an m -symmetric diffusion process on S . It admits no killing inside S and is a Hunt process on S in the sense that

$$\tilde{X}_{\tilde{\zeta}(\tilde{\omega})-}(\tilde{\omega}) = \Delta \quad \text{if } \tilde{\zeta}(\tilde{\omega}) < \infty.$$

(2) X^0 is identical in law with the process obtained from \tilde{X} by killing upon the hitting time σ_a of the point a .

Further the resolvent of \tilde{X} admits the next expression for $f \in \mathcal{B}(S)$:

$$G_\alpha f(x) = G_\alpha^0 f(x) + u_\alpha(x) \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)}, \quad x \in S_0, \quad (4.67)$$

$$G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)}, \quad (4.68)$$

where $L(m_0, \psi)$ is the energy functional of the X^0 -excessive measure $m_0 = \varphi \cdot m$ and the X^0 -excessive function $\psi = 1 - \varphi$.

Proof of Theorem 4.1. By Lemma 4.6, (4.37) and Lemma 4.13, we see that \tilde{X} is a Markov process on S with the m -symmetric resolvent (4.67),(4.68).

On account of **A.1**, we may assume that

$$X_t^0(\omega^0) \text{ is continuous in } t \in [0, \zeta^0(\omega^0)) \text{ and } X_{\zeta^0(\omega^0)-}(\omega^0) = a \cup \Delta$$

for every $\omega^0 \in \Omega^0$. We have already chosen Ω in a way that

$$X_t(\omega) \text{ is continuous in } t \in [0, \zeta_\omega) \text{ and } X_0(\omega) = a.$$

Hence the path $\tilde{X}_t(\tilde{\omega})$ defined by (4.63),(4.64),(4.65) is continuous on $[0, \tilde{\zeta})$.

Consider a function $u = G_\alpha f$ on S for $f \in C_b(S)$. By the assumptions **A.2,A.3** and the expression (4.67),(4.68), $u(X_t^0(\omega^0))$ is then continuous in $t \in [0, \sigma_a)$ for any $\omega^0 \in \Omega^0$. By the proof of Proposition 4.3, $u(X_t(\omega))$ is continuous in $t \in [0, \zeta_\omega)$ for any $\omega \in \Omega$. Hence $u(\tilde{X}_t(\tilde{\omega}))$ is right continuous in $t \in [0, \tilde{\zeta}(\tilde{\omega}))$ for any $\tilde{\omega} \in \tilde{\Omega}$. (In view of (4.33), we even know that $u(\tilde{X}_t)$ is continuous in $t \in [0, \tilde{\zeta})$ \tilde{P}_x -a.s. for any $x \in S$). Therefore we can conclude that \tilde{X} is a strong Markov process with continuous sample paths, namely, a diffusion process on S (cf.[2]). Clearly \tilde{X} is of no killing inside S and a Hunt process on S . The property (2) is also evident from the construction of \tilde{X} . \square

Remark 4.1. A prime reason for us to impose a regularity condition **A.4** on the given process X^0 on S_0 is in that it implies an important property in Lemma 4.3 of the excursion law \mathbf{n} of (4.4), which is essential in deriving the continuity near the point a of the process X constructed in §4.2.

Given a standard process \tilde{X} on S for which the point a is recurrent, K.Itô [15] associated with \tilde{X} a Poisson point process \mathbf{p} of excursions in the manner of §3.1 and gave a list of necessary conditions for the charactersitic measure \mathbf{n} of \mathbf{p} should obey. Conversely T.S. Salisbury [25], [26] constructed a right process on S for which a is recurrent by means of X^0 and an excursion law \mathbf{n} satisfying Itô's conditions being strengthened by adding the property as in Lemma 4.3 and some others.

Remark 4.2. By invoking the work of P.A. Meyer[20] on the absorbed Poisson point process and by adopting a similar argument to §4.2, we can show that Theorem 3.1 of §3.1 remains true without assuming condition **B.3** on the recurrence of the point $\{a\}$.

In this general case, the right continuous inverse $S(s)$ of the local time $L(t)$ at $\{a\}$ of the given process X on S is defined for $s \geq L(\infty)$ as $S(L(\infty)) = \infty$, and we see from Lemma 2.3 and by letting $\alpha \downarrow 0$ in (2.21) that $L(\infty)$ has an exponential distribution with mean $L(m_0, \psi)^{-1}$.

Let

$$\begin{aligned} D_{\mathbf{p}} &= \{s : S(s) - S(s-) > 0\}, \\ \mathbf{p}_s(t) &= X_{S(s-)+t}, \quad s \in D_{\mathbf{p}}, \quad 0 \leq t < S(s) - S(s-). \end{aligned}$$

Then $D_{\mathbf{p}} \subset (0, L(\infty)]$, $L(\infty) \in D_{\mathbf{p}}$ and $\{\mathbf{p}_s, s > 0\}$ is a point process with values in the space W_a defined by (4.13) instead of (3.6). Moreover, if we define the spaces W_a^+ , W_a^- by (4.14), (4.15) respectively, then

$$\mathbf{p}_s \in W_a^+ \text{ for } s \in D_{\mathbf{p}} \cap (0, L(\infty)), \quad \mathbf{p}_{L(\infty)} \in W_a^-.$$

By Theorem 5 of Meyer[20], $\{\mathbf{p}_s, s > 0\}$ is an *absorbed Poisson point process*. More precisely, on a certain probability space $(\tilde{\Omega}, \tilde{P})$, there is a Poisson point process $\{\tilde{\mathbf{p}}_s, s > 0\}$ on W_a with domain $D_{\tilde{\mathbf{p}}}$ and with the following properties.

(a) Let $\tilde{\zeta} = \inf\{s > 0 : \tilde{\mathbf{p}}_s \in W_a^-\}$ and consider the stopped point process $\{\tilde{\mathbf{p}}_s, s > 0\}$:

$$\tilde{\mathbf{p}}_s = \tilde{\mathbf{p}}_s \text{ for } s \in D_{\tilde{\mathbf{p}}} = D_{\tilde{\mathbf{p}}} \cap (0, \tilde{\zeta}].$$

Then the point process $\{\mathbf{p}_s, s > 0\}$ and $\{\tilde{\mathbf{p}}_s, s > 0\}$ are equivalent in law.

(b) Let \mathbf{n} be the characteristic measure of $\{\tilde{\mathbf{p}}_s, s > 0\}$. Then $\{w(t), \mathbf{n}\}$ is Markovian with respect to the transition function p_t^0 of X^0 . Let $\{\nu_t\}$ be the entrance law associated with \mathbf{n} . Then ν_t is a finite measure for each $t > 0$ and $\int_0^\infty e^{-t} \nu_t dt$ has a total mass not greater than 1.

We now prove that Theorem 3.1 *remains valid for this $\{\nu_t\}$ and for the entrance law $\{\mu_t\}$ specified by the equation (2.22).*

Take a bounded Borel function f on S and define $\hat{f}_\alpha(w)$, $w \in W_a$, $\alpha > 0$, as in the proof of Proposition 4.2. We have, almost surely with respect to P_a ,

$$\begin{aligned} \int_0^\zeta e^{-\alpha t} f(X_t) dt &= \sum_{s < L(\infty)} \int_{S(s-)}^{S(s)} e^{-\alpha t} f(X_t) dt + \int_{S(L(\infty)-)}^\infty e^{-\alpha t} f(X_t) dt \\ &= \sum_{s < L(\infty)} e^{-\alpha S(s-)} \hat{f}_\alpha(\mathbf{p}_s) + e^{-\alpha S(L(\infty)-)} \hat{f}_\alpha(\mathbf{p}_{L(\infty)}), \end{aligned}$$

which is equivalent in law to

$$\sum_{s < \tilde{\zeta}} e^{-\alpha \tilde{S}(s-)} \hat{f}_\alpha(\tilde{\mathbf{p}}_s^+) + e^{-\alpha \tilde{S}(\tilde{\zeta}-)} \hat{f}_\alpha(\tilde{\mathbf{p}}_{\tilde{\zeta}}), \quad (4.69)$$

where $\{\tilde{\mathbf{p}}_s^+, s > 0\}$ is a Poisson point process defined by $\tilde{\mathbf{p}}_s^+ = \tilde{\mathbf{p}}_s$ for $s \in D_{\tilde{\mathbf{p}}^+} = D_{\tilde{\mathbf{p}}} \cap \{s : \tilde{\mathbf{p}}_s \in W_a^+\}$ and $\tilde{S}(s) = \sum_{r \leq s} \zeta(\tilde{\mathbf{p}}_r^+)$. The characteristic measure of $\{\tilde{\mathbf{p}}_s^+, s > 0\}$ is the

restriction \mathbf{n}^+ of \mathbf{n} on W_a^+ . In the same way as in the proof of Lemma 4.5, we can prove that

$$\tilde{E}(e^{-\alpha\tilde{S}(s)}) = \exp(-\alpha\hat{\nu}_\alpha(\varphi)), \quad \hat{\nu}_\alpha = \int_0^\infty e^{-\alpha t} \nu_t dt.$$

Now the value $G_\alpha f(a)$ equals the expectation of the random variable (4.69) with respect to \tilde{P} , which can be evaluated by taking into account of the following facts.

- (i) The three objects $\{\tilde{\mathbf{p}}_s^+, s > 0\}$, $\tilde{\zeta}$ and $\tilde{\mathbf{p}}_\zeta$ are independent.
- (ii) $\tilde{\zeta}$ has an exponential distribution with mean $L(m_0, \psi)^{-1}$.
- (iii) The law of $\tilde{\mathbf{p}}_\zeta$ is $L(m_0, \psi)^{-1} \mathbf{n}^-$ where \mathbf{n}^- is the restriction of \mathbf{n} on W_a^- .

Indeed, exactly the same computation as in the proof of Propostion 4.2 leads us to

$$G_\alpha f(a) = \frac{\hat{\nu}_\alpha(f)}{\alpha\hat{\nu}_\alpha(\varphi) + L(m_0, \psi)}, \quad (4.70)$$

which combined with (2.15) and Lemma 2.2 (ii) yields

$$\frac{\hat{\nu}_\alpha(f)}{\alpha\hat{\nu}_\alpha(\varphi) + L(m_0, \psi)} = \frac{\hat{\mu}_\alpha(f)}{\alpha\hat{\mu}_\alpha(\varphi) + L(m_0, \psi)}$$

Therefore for each $\alpha > 0$ there is a constant c_α such that $\hat{\nu}_\alpha = c_\alpha \hat{\mu}_\alpha$. Inserting this into the above equation, we easily obtain $c_\alpha = 1$ and so $\nu_t = \mu_t$, $t > 0$.

5 Uniqueness of the symmetric extension and expression of its Dirichlet form

In the preceding section, we have started with an m -symmetric diffusion X^0 on S_0 satisfying conditions **A.1, A.2, A.3, A.4**, and constructed a process \tilde{X} on S satisfying properties **(1), (2)** stated in Theorem 4.1. Let us call a process on S satisfying conditions **(1), (2)** a *symmetric extension of X^0* . In this section, we are concerned with the uniqueness of a symmetric extension of X^0 and explicit expression of its Dirichlet form on $L^2(S; m)$. We aim at proving the following:

Theorem 5.1. *Assume that an m -symmetric diffusion X^0 on S_0 satisfies conditions **A.1, A.2**. Let \hat{X} be a symmetric extension of X^0 and $(\mathcal{E}, \mathcal{F})$ be the Dirichlet form on $L^2(S; m)$ of \hat{X} .*

- (i) \hat{X} admits the resolvent identical with (4.67), (4.68).
- (ii) $(\mathcal{E}, \mathcal{F})$ admits the expression

$$\mathcal{F}_e = \{w = u_0 + c\varphi : u_0 \in \mathcal{F}_{0,e}, c \text{ constant}\}, \quad \mathcal{F} = \mathcal{F}_e \cap L^2(S; m), \quad (5.1)$$

$$\mathcal{E}(w, w) = \mathcal{E}(u_0, u_0) + c^2 \mathcal{E}(\varphi, \varphi), \quad \mathcal{E}(\varphi, \varphi) = L(m_0, \psi), \quad (5.2)$$

where $(\mathcal{F}_{0,e}, \mathcal{E})$ is the extended Dirichlet space of X^0 and $L(m_0, \psi)$ is the energy functional of $m_0 = \varphi \cdot m$ and ψ with respect to X^0 .

- (iii) X^0 satisfies **(A.3)** automatically: $u_\alpha \in L^1(S; m)$, $\alpha > 0$.
- (iv) $\tilde{P}_a(\sigma_a = 0, \tau_a = 0) = 1$
where $\sigma_a = \inf\{t > 0 : X_t = a\}$, $\tau_a = \inf\{t > 0 : X_t \in S_0\}$.
- (v) $(\mathcal{E}, \mathcal{F})$ is irreducible.

Corollary 5.1. *Under the conditions **A.1, A.2** for an m -symmetric diffusion X^0 on S_0 , the symmetric extension of X^0 is unique in law.*

Corollary 5.1 follows from Theorem 5.1 (i). We prepare a lemma before the proof of Theorem 5.1.

Assume that $X = (X_t, P_x)$ is an m -symmetric Hunt process on S and $(\mathcal{E}, \mathcal{F})$ is the associated Dirichlet form on $L^2(S; m)$. No regularity for the Dirichlet form $(\mathcal{E}, \mathcal{F})$ is assumed in advance.

In accordance with [19], we set for a closed set $F \subset S$,

$$\mathcal{F}_F = \{u \in \mathcal{F} : u = 0 \text{ } m\text{-a.e. on } S \setminus F\},$$

and call an increasing family $\{F_n\}$ of closed subsets of S an \mathcal{E} -nest if the space $\cup_{n=1}^{\infty} \mathcal{F}_{F_n}$ is \mathcal{E}_1 -dense in \mathcal{F} . A set N is called \mathcal{E} -exceptional if $N \subset \cap_{n=1}^{\infty} F_n^c$ for some \mathcal{E} -nest $\{F_n\}$. On the other hand, we call a set $N \subset S$ an X -exceptional set if there exists a Borel set $B_1 \supset B$ with

$$P_m(\sigma_{B_1} < \infty) = 0.$$

A nearly Borel set $N \subset S$ is called X -properly exceptional if $m(N) = 0$ and $S \setminus N$ is X -invariant in the sense that

$$P_x(X_t \in S_{\Delta} \setminus N \text{ or } X_{t-} \in S_{\Delta} \setminus N \exists t \geq 0) = 1, \quad \forall x \in S \setminus N.$$

Lemma 5.1. (i) *The following properties of a set $N \subset S$ are equivalent each other:*
 α . N is \mathcal{E} -exceptional.
 β . N is X -exceptional.
 γ . N is contained in an X -properly exceptional Borel set.
(ii) *If $\{F_n\}$ is an \mathcal{E} -nest, then*

$$P_x \left(\lim_{n \rightarrow \infty} \sigma_{S \setminus F_n} \geq \zeta \right) = 1 \quad \text{q.e.}, \quad (5.3)$$

where q.e. means ‘except on a set $N \subset S$ satisfying one of the properties in (i)’.

(iii) $(\mathcal{E}, \mathcal{F})$ is a quasi-regular Dirichlet form on $L^2(S; m)$ in the sense of [19, §IV 3].

Proof. (i). The equivalences $\alpha \Leftrightarrow \beta$ and $\beta \Leftrightarrow \gamma$ were proved in [19, Th.5.29] and in [9, Th.4.1.1] respectively.

(ii). Put $\sigma = \lim_{n \rightarrow \infty} \sigma_{S \setminus F_n}$. On account of [19, Th.2.11, Th.5.4], we have for a strictly positive bounded m -integrable function f on S ,

$$E_x \left(\int_{\sigma \wedge \zeta}^{\zeta} e^{-s} f(X_s) ds \right) = 0 \quad m\text{-a.e. } x \in S.$$

Since the function of x on the left hand side of the above equation is X -excessive, it is finely continuous on S and hence the above equation holds q.e. by [9, Lemma 4.1.5].

(iii) Since $(\mathcal{E}, \mathcal{F})$ is associated with a Hunt process X , it must be quasi-regular by virtue of [19, Th.5.1]. \square

Proof of Theorem 5.1. Since \hat{X} is not only a diffusion process but also a Hunt process on S , the Dirichlet form $(\mathcal{E}, \mathcal{F})$ of \hat{X} is quasi-regular by the above lemma.

Consequently we can invoke [3, Th.3.7] to find a regular Dirichlet space $(S', m', \mathcal{F}', \mathcal{E}')$ related to the quasi-regular Dirichlet space $(S, m, \mathcal{F}, \mathcal{E})$ by a quasi-homeomorphism q : there exist an \mathcal{E} -nest $\{F_n\}$ on S and an \mathcal{E}' -nest $\{F'_n\}$ on S' such that q is a one to one mapping from $S_1 = \cup_{n=1}^{\infty} F_n$ onto $S'_1 = \cup_{n=1}^{\infty} F'_n$ and its restriction on each F_n is homeomorphic to F'_n . Further, m' is the image measure of m by q and the space $(\mathcal{F}', \mathcal{E}')$ is also the image of $(\mathcal{F}, \mathcal{E})$ by q . Thus, if we put $(\Phi u)(x') = u(q^{-1}(x'))$, $x' \in S'_1$, then

$$\int_{S'} (\Phi u) dm' = \int_S u dm, \quad \forall u \geq 0; \quad \mathcal{F}' = \Phi(\mathcal{F}), \quad \mathcal{E}'(\Phi u, \Phi v) = \mathcal{E}(u, v), \quad u, v \in \mathcal{F}. \quad (5.4)$$

We note that $S \setminus S_1$ (resp. $S' \setminus S'_1$) is \mathcal{E} - (resp. \mathcal{E}' -) exceptional and, when $N' = q(N)$, N is \mathcal{E} -exceptional if and only if N' is \mathcal{E}' -exceptional (cf.[3, Cor.3.6].)

For a Borel set $B \subset S$, we denote by B_{Δ} the subset $B \cup \Delta$ of S_{Δ} with induced topology. The above q can then be extended to a homeomorphism between $(F_n)_{\Delta}$ and $(F'_n)_{\Delta'}$ for each n , where Δ' denotes the point at infinity of S' (which is added as an isolated point when S' is compact).

We now apply Lemma 5.1 to the above \mathcal{E} -nest $\{F_n\}$ in finding an \hat{X} -properly exceptional Borel set $\hat{N} \subset S$ containing $S \setminus S_1$ such that (5.3) holds for any $x \in S \setminus \hat{N}$. q is then a one to one mapping between $S \setminus \hat{N}$ and $S' \setminus \hat{N}'$, where

$$\hat{N}' = (S' \setminus S'_1) \cup q(S \cap \hat{N}).$$

In view of condition **A.2** for X^0 , condition **(2)** for \hat{X} and the above observation, the one point set $\{a\}$ is not \hat{X} -exceptional and consequently it is not \mathcal{E} -exceptional by virtue of Lemma 5.1. Therefore a must be located in $S \setminus \hat{N}$ and furthermore

$$\{a'\} \text{ is not } \mathcal{E}' \text{ - exceptional}, \quad (5.5)$$

where $a' = q(a) \in S' \setminus \hat{N}'$.

The restriction of \hat{X} to $S \setminus \hat{N}$ is a diffusion with no killing inside $S \setminus \hat{N}$ and we denote it again by

$$\hat{X} = \left(\Omega, \mathcal{F}_t, \hat{X}_t, \hat{\zeta}, \hat{P}_x \right).$$

Let us transfer \hat{X} to a process

$$\hat{X}' = \left(\Omega, \mathcal{F}_t, \hat{X}'_t, \hat{\zeta}', \hat{P}'_x \right)$$

on $S' \setminus \hat{N}'$ by the mapping q :

$$\hat{X}'_t(\omega) = q(\hat{X}_t(\omega)), \quad \hat{\zeta}'(\omega) = \hat{\zeta}(\omega), \quad \omega \in \Omega, \quad t \geq 0,$$

$$\hat{P}'_x(\Lambda) = \hat{P}_{q^{-1}x}(\Lambda) \quad x \in S' \setminus \hat{N}', \quad \Lambda \in \mathcal{F}_{\infty}.$$

We may extend the state space of \hat{X}' to S' by making each point of \hat{N}' trap. It is then easy to see that \hat{X}' is a diffusion process on S' with no killing inside S' in the sense that

$$\hat{P}'_x \left(\hat{\zeta}' < \infty, \hat{X}'_{\hat{\zeta}'_-} = \Delta \right) = \hat{P}'_x(\hat{\zeta}' < \infty). \quad (5.6)$$

Further \hat{X}' is associated with the Dirichlet form $(\mathcal{E}', \mathcal{F}')$ which is regular. Since \hat{X}' is a diffusion without killing inside S' , $(\mathcal{E}', \mathcal{F}')$ must be strongly local (cf.[9, Th.4.5.3]).

By (5.5) and Lemma 5.1, we see that the one point set $\{a'\}$ is not \hat{X}' -exceptional and consequently it has a positive capacity with respect to $(\mathcal{E}', \mathcal{F}')$ in virtue of [9, Th.4.2.1].

Therefore $(\mathcal{E}', \mathcal{F}')$ and \hat{X}' fit the setting of §2 and they satisfy all the properties stated in Theorem 2.1 of §2. In particular, we have the next expressions of the resolvent and $(\mathcal{E}', \mathcal{F}')$ of \hat{X}' in terms of the part $\hat{X}'^{',0}$ of \hat{X}' on $S'_0 = S' \setminus \{a'\}$: if we denote the transition function and the resolvent of \hat{X}' (resp. $\hat{X}'^{',0}$) by p'_t, G'_α (resp. $p_t^{',0}, G_\alpha^{',0}$), then

$$G'_\alpha g(a') = \frac{(u'_\alpha, g)_{m'}}{\alpha(u'_\alpha, \varphi')_{m'} + L'(m'_0, \psi')} \quad (5.7)$$

$$\mathcal{E}'(\varphi', \varphi') = L'(m'_0, \psi'), \quad (5.8)$$

where φ' (resp. u'_α) is the hitting (resp. α -order hitting) probability of $\{a'\}$ of the process \hat{X}' , $\psi' = 1 - \varphi'$ and

$$L'(m'_0, \psi') = \lim_{t \downarrow 0} \frac{1}{t} (\varphi' - p_t^{',0} \varphi', \psi')_{m'}. \quad (5.9)$$

Notice that the part $(\mathcal{E}', \mathcal{F}'_0)$ of $(\mathcal{E}', \mathcal{F}')$ on S'_0 is associated with $\hat{X}'^{',0}$ which can be sent from X^0 on S_0 by the mapping q in the same way as above on account of the property **(2)** of \hat{X} . Hence we have for $x \in S' \setminus \hat{N}'$

$$\begin{aligned} \Phi(G_\alpha f)(x) &= G'_\alpha(\Phi f)(x), \quad \Phi(G_\alpha^0 f)(x) = G_\alpha^{',0}(\Phi f)(x), \quad \Phi(p_t^0 f)(x) = p_t^{',0}(\Phi f)(x), \\ \Phi(\varphi)(x) &= \varphi'(x), \quad \Phi(u_\alpha)(x) = u'_\alpha(x). \end{aligned} \quad (5.10)$$

(5.4),(5.7),(5.8),(5.9) and (5.10) now imply $L'(m'_0, \psi') = L(m_0, \psi)$ and furthermore

$$\mathcal{E}(\varphi, \varphi) = L(m_0, \psi), \quad G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, \varphi) + L(m_0, \psi)}. \quad (5.11)$$

We have obtained the expression (4.68) of the resolvent G_α of \hat{X} . It then satisfies (4.67) for all $x \in S_0$ because of the property **(2)** of \hat{X} . We can also readily get the assertions (ii) and (iii) of Theorem 5.1 using (5.4) and (5.10). As for (iv), we have obviously

$$\hat{P}_a(\sigma_a = 0, \tau_a = 0) = \hat{P}'_{a'}(\sigma_{a'} = 0, \tau_{a'} = 0),$$

and the right hand side equals 1 by virtue of Theorem 2.1. From the expression (4.67) of the resolvent of \hat{X} , we have

$$(I_A, G_\alpha I_B) > 0 \quad \text{for any } A, B \in \mathcal{B}(S) \text{ with } m(A) > 0, m(B) > 0.$$

This property is equivalent to the irreducibility of the Dirichlet form $(\mathcal{E}, \mathcal{F})$ proving (v). \square

Remark 5.1. For the symmetric extension \tilde{X} of X^0 constructed in §4, not only the expression (4.67),(4.68) of its resolvent but also the property (iv) in Theorem 5.1 have been directly proved in Lemma 4.12.

6 Examples

Example 6.1. Let X be the Brownian motion on \mathbb{R} , X^0 be the absorbed Brownian motion on $\mathbb{R} \setminus \{0\}$ and m be the Lebesgue measure dx on \mathbb{R} . Then X is the unique m -symmetric extension of X^0 (in the sense that X satisfies conditions **(1),(2)** of Theorem 4.1) in accordance with Corollary 5.1.

Let $L(t)$ be the local time of X at 0 and Z be an independent exponential random variable with mean δ^{-1} . The process X_δ obtained from X killed upon the first time that $L(t) \geq Z$ is a diffusion process extending X^0 but not a symmetric extension of X^0 in the present sense because it violates the above condition **(1)**.

For $\gamma > 0$, let X^γ be the process on \mathbb{R} obtained from X by a time change with respect to the inverse of its additive functional $t + \gamma L(t)$. X^γ is then a diffusion on \mathbb{R} with a canonical scale $2dx$ and the speed measure $m(dx) = dx + \gamma \delta_0(dx)$. X^γ extends X^0 but violates our assumption that $m(\{0\}) = 0$.

The resolvents and Dirichlet forms of X_δ , X^γ have been exhibited in Remark 2.2.

Example 6.2. Let D be a bounded open set in \mathbb{R}^d , ($d \geq 1$), and $L^2(D)$ be the L^2 -space based on the Lebesgue measure on D . Denote by $H_0^1(D)$ the closure of $C_0^1(D)$ in the Sobolev space

$$H^1(D) = \{u \in L^2(D) : \frac{\partial u}{\partial x_i} \in L^2(D), 1 \leq i \leq n\}$$

and put

$$\mathbf{D}(u, v) = \int_D \nabla u \cdot \nabla v(x) dx, \quad u, v \in H_0^1(D).$$

Then $(\frac{1}{2}\mathbf{D}, H_0^1(D))$ is a strongly local Dirichlet form on $L^2(D)$ satisfying the Poincaré inequality (3.13). The associated symmetric diffusion $X^0 = (X_t^0, 0 \leq t < \zeta^0, P_x^0)$ on D is the absorbing Brownian motion.

Let $D^* = D \cup \{a\}$ be the one point compactification of D . Regarding D as a subspace of D^* , we have then

$$\varphi(x) = P_x^0(\zeta^0 < \infty, X_{\zeta^0-}^0 = a) = 1, \quad \psi(x) = 1 - \varphi(x) = 0, \quad \forall x \in D, \quad (6.1)$$

$$u_\alpha(x) = E_x^0(e^{-\alpha \zeta^0}; X_{\zeta^0-}^0 = a) \text{ is continuous in } x \in D, \quad (\alpha > 0). \quad (6.2)$$

Obviously $u_\alpha \in L^1(D)$. Hence conditions **A.1, A.2, A.3, A.4** are satisfied by X^0 and we can construct a diffusion \tilde{X} on D^* as in §4. By virtue of Theorem 4.1, the resolvent of \tilde{X} is expressed as

$$G_\alpha f(x) = G_\alpha^0 f(x) + u_\alpha(x) \frac{(u_\alpha, f)}{\alpha(u_\alpha, 1)}, \quad x \in D, \quad G_\alpha f(a) = \frac{(u_\alpha, f)}{\alpha(u_\alpha, 1)},$$

and in particular, \tilde{X} is conservative.

$L^2(D^*)$ denotes the L^2 -space based on the 0-extension of the Lebesgue measure on D to D^* . By virtue of Theorem 4.1 and Theorem 5.1, \tilde{X} is symmetric with respect to this measure and its Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(D^*)$ is describable as

$$\mathcal{F} = H_0^1(D) + \text{constant functions on } D^*, \quad (6.3)$$

$$\mathcal{E}(w_1, w_2) = \frac{1}{2} \mathbf{D}(f_1, f_2), \quad w_i = f_i + c_i, \quad f_i \in H_0^1(D), \quad c_i \text{ constant}, \quad i = 1, 2. \quad (6.4)$$

On account of Theorem 3.2 and a related observation in §3.1, this is a regular, strongly local and irreducible recurrent Dirichlet form. This Dirichlet form first appeared in [8].

The entrance law $\{\mu_t\}_{t>0}$ governing the charactersitic measure of the excursion valued Poisson point process attached to \tilde{X} is given by

$$\mu_t(B)dt = \int_B P_x^0(\zeta^0 \in dt)dx, \quad B \in \mathcal{B}(D) \quad (6.5)$$

in view of (3.9). Let $D = \cup_i D_i$ be the decomposition of the open set D into connected components. The above identity tells us that the sample path of \tilde{X} entering from the point a is distributed among $\{D_i\}$ proportionally to their volumes and enters in D_i according to the restriction of μ_t to D_i . As was observed in §3.1, \tilde{X} is irreducible recurrent.

According to (2.24), the Lévy measure of the inverse local time of \tilde{X} at the point a is given by $-d\mu_t(D)$.

Example 6.3. We consider a finite number of disjoint rays $\ell_i, i = 1, \dots, N$, on \mathbb{R}^2 merging at a point $a \in \mathbb{R}^2$. Each ray ℓ_i is homeomorphic to the open half line $(0, \infty)$ and the point a is the boundary of each ray at 0-side. We put

$$S_0 = \sum_{i=1}^N \ell_i, \quad S = S_0 + a.$$

S is endowed with the induced topology as a subset of \mathbb{R}^2 .

Let m be a positive Radon measure on S_0 with $\text{Supp}[m] = S_0$. m is extended to S by setting $m(\{a\}) = 0$. The restriction of m to ℓ_i is denoted by m_i . For any function g on S_0 , its restriction to ℓ_i will be denoted by g_i . We consider a diffusion process $X^0 = \{X_t^0, \zeta^0, P_x^0\}$ on S_0 such that its restriction $X^{0,i}$ to each open half line $\ell_i \sim (0, \infty)$ is the absorbing diffusion governed by the speed measure m_i and a canonical scale, say s_i .

We notice that X^0 satisfies **A.2, A.3** if and only if 0 is a regular boundary in Feller's sense for each diffusion $X^{0,i}$ on ℓ_i , $1 \leq i \leq N$. Indeed, **A.2** holds if and only if 0 is exit (in the terminology used by [16]). If 0 is additionally non-entrance, then $m_i((0, 1)) = \infty$ and **A.3** is not satisfied. If 0 is regular, then $m_i((0, 1)) < \infty$ and $u_{\alpha,i}$ is m_i integrable on $(0, 1)$, while $u_{\alpha,i}$ is always m_i -integrable on $[1, \infty)$ (cf.[16, p 130].)

Thus we assume that 0 is regular for every $X^{0,i}$ so that **A.1, A.2, A.3** are satisfied by X^0 . **A.4** is also clearly satisfied. m is finite on any compact neighbourhood of a .

Therefore, a diffusion \tilde{X} on S can be constructed as in §4 and it is a unique m -symmetric extension of X^0 with no killing inside S according to Theorem 5.1. The resolvent of \tilde{X} has the expression

$$G_\alpha f(a) = \frac{\sum_i (u_{\alpha,i}, f_i)_{m_i}}{\alpha \sum_i (u_{\alpha,i}, \varphi_i)_{m_i} + \sum_i L(\varphi_i \cdot m_i, \psi_i)}.$$

The Dirichlet form $(\mathcal{E}, \mathcal{F})$ of \tilde{X} on $L^2(S; m)$ is regular, strongly local, irreducible and can be described as follows:

$$\mathcal{F}_e = \{w = u_0 + c\varphi : u_0 \in \mathcal{F}_{0,e}, c \text{ constant}\},$$

$$\mathcal{E}(w, w) = \mathcal{E}(u_0, u_0) + c^2 \mathcal{E}(\varphi, \varphi),$$

$$\mathcal{E}(\varphi, \varphi) = \sum_i L(\varphi_i \cdot m_i, \psi_i),$$

where

$$\mathcal{F}_{0,e} = \{u : u_i \text{ is absolutely continuous with respect to } s_i, \\ \int_0^\infty \left(\frac{du_i}{ds_i}\right)^2 ds_i < \infty, u_i(0) = 0, u_i(\infty) = 0, \text{ whenever } \infty \text{ is regular, } 1 \leq i \leq n\},$$

$$\mathcal{E}(u, u) = \sum_i \int_0^\infty \left(\frac{du_i}{ds_i}\right)^2 ds_i \quad u \in \mathcal{F}_{0,e}.$$

Related Dirichlet forms and diffusions first appeared in [13].

The entrance law from a is describable as

$$\mu_t(f)dt = \sum_i P_{f_i \cdot m_i}^{0,i} \left(\zeta^{0,i} \in dt, X_{\zeta^{0,i}-}^{0,i} = 0 \right). \quad (6.6)$$

We have a freedom of choice of the entrance law (6.6) in the following sense. Choose any positive numbers $\{p_1, \dots, p_N\}$ and observe that the absorbed diffusion X^0 on S_0 is unchanged if we replace $m_i, s_i, 1 \leq i \leq N$, by

$$\hat{m}_i = p_i \cdot m_i, \quad \hat{s}_i = p_i^{-1} \cdot s_i, \quad 1 \leq i \leq N,$$

respectively. Let \hat{m} be the measure on S whose restriction to ℓ_i equals \hat{m}_i for each $i = 1, 2, \dots, N$, with $\hat{m}(\{a\}) = 0$. Then we can consider the \hat{m} -symmetric extension \hat{X} of X^0 whose entrance law $\hat{\mu}$ from a is given by (6.6) but with the replacement of m_i by \hat{m}_i for $1 \leq i \leq N$.

Example 6.4. Let G_1, G_2 be open sets of \mathbb{R}^d , ($d \geq 1$), such that

$$\overline{G_1} \subset G_2, \quad \overline{G_1} \text{ is compact.}$$

We let $S_0 = G_2 \setminus \overline{G_1}$. We consider the space $S = S_0 \cup \{a\}$ equipped with the topology where a set U containing a is defined to be an open set if

$$U \setminus \{a\} = \{\text{open subset of } G_2 \text{ containing } \overline{G_1}\} \setminus \overline{G_1}.$$

Let X^0 be the absorbing Brownian motion on S_0 . Then conditions **A.1, A.2, A.3, A.4** are satisfied by X^0 . **A.3** can be verified by a comparison with the Brownian motion on \mathbb{R}^d .

Let m be the Lebesgue measure on S_0 extended to S by $m(\{a\}) = 0$. Let \tilde{X} be the m -symmetric diffusion on S as is constructed in §4. Then, by Theorem 5.1, its Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(S; m)$ is expressed as

$$\mathcal{F} = \mathcal{F}_e \cap L^2(S; m), \quad \mathcal{F}_e = \{w = u_0 + c\varphi : u_0 \in H_{0,e}^1(S_0), c \text{ constant}\},$$

$$\mathcal{E}(w, w) = \frac{1}{2} \mathbf{D}(u_0, u_0) + c^2 L(\varphi \cdot m, \psi),$$

where $H_{0,e}^1(S_0)$ denotes the extended Dirichlet space of $H_0^1(S_0)$.

$(\mathcal{E}, \mathcal{F})$ is a quasi-regular Dirichlet form on $L^2(S; m)$ but may not be regular. It is a regular Dirichlet space if each point of ∂G_1 is a regular boundary point of S_0 with respect to the Dirichlet problem for $(\alpha - \frac{1}{2}\Delta)$ on S_0 .

References

- [1] R.M. Blumenthal, *Excursions of Markov processes*, Birkhäuser, Boston, 1992
- [2] R.M. Blumenthal and R.K. Gettoor, *Markov processes and potential theory*, Academic Press, New York, 1968
- [3] Z.-Q. Chen, Z.-M. Ma and M. Röckner, Quasi-homeomorphisms of Dirichlet forms, Nagoya Math. J. 136(1994), 1-15
- [4] C. Dellacherie et P.A. Meyer, *Probabilités et potentiel*, Chap. XII, Hermann, Paris, 1987
- [5] C. Dellacherie, B. Maisonneuve et P.A. Meyer, *Probabilités et potentiel*, Chap. XVII-XXIV, Hermann, Paris, 1992
- [6] E.B. Dynkin, An application of flows to time shift and time reversal in stochastic processes, Trans. Amer. Math. Soc. 287(1985), 613-619
- [7] P.J. Fitzsimmons, On the excursions of Markov processes in classical duality, Probab. Th. Rel. Fields 75(1987), 159-178
- [8] M. Fukushima, On boundary conditions for multi-dimensional Brownian motions with symmetric resolvent densities, J. Math. Soc. Japan 21(1969), 58-93
- [9] M. Fukushima, Y. Oshima and M. Takeda, *Dirichlet Forms and Symmetric Markov Processes*, Walter de Gruyter, 1994
- [10] R.K. Gettoor, Excursions of a Markov process, Ann. Probability, 7(1979), 244-266
- [11] R.K. Gettoor, *Excessive measures*, Birkhäuser, 1990
- [12] R.K. Gettoor and M.J. Sharpe, Excursions of dual processes, adv. Math. 45(1982), 259-309
- [13] N. Ikeda and S. Watanabe, The local structure of a class of diffusions and related problems, in: Lecture Notes in Math. Vol 330, 1973, pp124-159
- [14] N. Ikeda and S. Watanabe, *Stochastic differential equations and diffusion processes*, North-Holland/Kodansha, 1981
- [15] K. Itô, Poisson point processes attached to Markov processes, in: Proc. Sixth Berkeley Symp. Math. Stat. Probab. III, 1970, pp225-239
- [16] K. Itô and H.P. McKean, *Diffusion processes and their sample paths*, Springer, 1970
- [17] D. Kim, On spectral gaps and exit time distributions for a non-smooth domain, to appear in Forum Math.
- [18] Y. Le Jan, Dual markovian semigroups and processes, in: Functional Analysis in Markov processes, Proceedings, Katata and Kyoto 1981, Lecture Notes in Math. Vol 923, 1982, pp47-75

- [19] Z.M. Ma and M. Röckner, *Introduction to the theory of (non-symmetric) Dirichlet forms*, Springer-Verlag, 1992
- [20] P.A. Meyer, Processus de Poisson ponctuels, d'après K.Itô, *Séminaire de Probab. V*, in: Lecture Notes in Math., Vol,191, Springer, Berlin, 1971,pp.177-190
- [21] P.A. Meyer, Note sur l'interprétation des mesures d'équilibre, *Séminaire de Probab. VII*, in: Lecture Notes in Math., Vol. 321, 1973, pp. 210-216
- [22] J. Mitro, Time reversal depending on local time, *Stochastic process. Appl.* 18(1984), 171-177
- [23] M. Nagasawa, Time reversion of Markov processes, *Nagoya Math. J.* 24(1964), 177-204
- [24] L.C.G. Rogers, Itô excursion theory via resolvents, *Z. Wahrsch. Verw. Gebiete* 63(1983), 237-255
- [25] Thomas S. Salisbury, On the Itô excursion process, *Probab.Theory Related Fields* 73(1986), 319-350
- [26] Thomas S. Salisbury, Construction of right processes from excursions, *Probab.Theory Related Fields* 73(1986), 351-367