

Research Report

KSTS/RR-04/002

Jan. 29, 2004

**Poles and α -points of meromorphic solutions of
the first Painlevé hierarchy**

by

Shun Shimomura

<p>Shun Shimomura Department of Mathematics Keio University</p>

Department of Mathematics
Faculty of Science and Technology
Keio University

©2004 KSTS

3-14-1 Hiyoshi, Kohoku-ku, Yokohama, 223-8522 Japan

Poles and α -points of Meromorphic Solutions of the First Painlevé Hierarchy

By
SHUN SHIMOMURA *

Abstract

The first Painlevé hierarchy, which is a sequence of higher order analogues of the first Painlevé equation, follows from the singular manifold equations for the mKdV hierarchy. For meromorphic solutions of the first Painlevé hierarchy, we give a lower estimate for the number of poles; which is regarded as an extension of one corresponding to the first Painlevé equation, and which indicates a conjecture on the growth order. From our main result, two corollaries follow: one is the transcendency of meromorphic solutions, and the other is a lower estimate for the frequency of α -points. An essential part of our proof is estimation of certain sums concerning the poles of each meromorphic solution.

§1. Introduction

For a meromorphic function $f(z)$ in the whole complex plane \mathbf{C} , the counting function for poles of $f(z)$ is defined by

$$N(r, f) := \int_0^r (n(\rho, f) - n(0, f)) \frac{d\rho}{\rho} + n(0, f) \log r,$$

where $n(r, f)$ denotes the number of poles inside the disk $|z| \leq r$, each counted according to its multiplicity. Moreover, we use the notation (cf. [6], [8]):

$$m(r, f) := \frac{1}{2\pi} \int_0^{2\pi} \log^+ |f(re^{i\phi})| d\phi, \quad \log^+ x := \max\{0, \log x\},$$

$$T(r, f) := m(r, f) + N(r, f)$$

Communicated by T. Kawai. Received May 15, 2003.

2000 Mathematics Subject Classification(s): 34M55, 34M05, 30D35

*Department of Mathematics, Keio University, 3-14-1, Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan

e-mail: shimomur@math.keio.ac.jp

denoting, respectively, the proximity and the characteristic functions; the growth order of $f(z)$ is defined by

$$\sigma(f) := \limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r}.$$

Let $w(z)$ be an arbitrary solution of the first Painlevé equation

$$(PI) \quad w'' = 6w^2 + z$$

($' = d/dz$). Then, $w(z)$ is a transcendental meromorphic function. By a well-known argument in the Nevanlinna theory, $\limsup_{r \rightarrow \infty} (N(r, w)/\log r) = \infty$ (cf. Remark 1.2), which implies that $w(z)$ admits infinitely many poles. This fact is quantitatively represented as

$$(1.1) \quad \limsup_{r \rightarrow \infty} \frac{\log N(r, w)}{\log r} \geq \frac{5}{2}$$

(cf. [11]); and, more precisely,

$$(1.2) \quad N(r, w) \gg r^{5/2}/\log r$$

(cf. [5, §7], [13]). These combined with $T(r, w) \ll r^{5/2}$ (cf. [5, §8], [12], [14]) imply that the growth order of $w(z)$ is equal to $5/2$. (For real-valued functions $\phi(r)$ and $\psi(r)$ on the interval $(r_0, +\infty)$, we write $\phi(r) \ll \psi(r)$ or $\psi(r) \gg \phi(r)$ if $\phi(r) = O(\psi(r))$ as $r \rightarrow +\infty$. In the case where $h(z)$ is a function of $z \in \mathbf{C}$, we also write $|h(z)| \ll \psi(r)$ if $|h(z)| = O(\psi(r))$ as $|z| = r \rightarrow +\infty$.)

A sequence of higher order analogues of (PI) is given in the following manner (cf. [5, §16], [7]). Let $d_\nu[w]$ ($\nu = 0, 1, 2, \dots$) be differential polynomials in w determined by the recursion relation

$$(1.3) \quad d_0[w] = 1,$$

$$(1.4) \quad Dd_{\nu+1}[w] = (D^3 - 8wD - 4w')d_\nu[w], \quad D = d/dz, \quad \nu \in \mathbf{N} \cup \{0\}$$

(cf. Lemma 2.6 with its proof). Some of them are written in the form

$$\begin{aligned} d_1[w]/4 &= -w + C_{10}, \\ d_2[w]/4 &= -w'' + 6w^2 + C_{21}d_1[w] + C_{20}, \\ d_3[w]/4 &= -w^{(4)} + 20ww'' + 10(w')^2 - 40w^3 + C_{32}d_2[w] + C_{31}d_1[w] + C_{30}, \\ d_4[w]/4 &= -w^{(6)} + 28ww^{(4)} + 56w'w^{(3)} + 42(w'')^2 - 280(w^2w'' + w(w')^2 - w^4) \\ &\quad + C_{43}d_3[w] + C_{42}d_2[w] + C_{41}d_1[w] + C_{40}, \end{aligned}$$

where C_{ij} are arbitrary constants. Consider a sequence of 2ν -th order equations of the form

$$(PI_{2\nu}) \quad d_{\nu+1}[w] + 4z = 0, \quad \nu \in \mathbf{N},$$

which is called *the first Painlevé hierarchy*. Equation (PI_2) essentially coincides with (PI) . These equations follow from the singular manifold equations for the mKdV hierarchy (cf. [7], [9], [15]). As in the case of (PI) , it is basic and interesting to study analytic properties of meromorphic solutions of $(PI_{2\nu})$, for example, to determine the growth order of them.

The purpose of this paper is to show the following, which is an extension of (1.1), and which is a first step toward this question:

Theorem 1.1. *Suppose that $w_\nu(z)$ be a meromorphic solution of $(PI_{2\nu})$. Then we have*

$$(1.5) \quad \limsup_{r \rightarrow \infty} \frac{\log N(r, w_\nu)}{\log r} \geq \frac{2\nu + 3}{\nu + 1},$$

namely the growth order of $w_\nu(z)$ is not less than $(2\nu + 3)/(\nu + 1)$.

Corollary 1.2. *Equation $(PI_{2\nu})$ admits no rational solutions.*

Furthermore, the frequency of α -points is estimated as follows:

Corollary 1.3. *For each $\alpha \in \mathbf{C}$,*

$$(1.6) \quad \limsup_{r \rightarrow \infty} \frac{\log N(r, 1/(w_\nu - \alpha))}{\log r} \geq \frac{2\nu + 3}{\nu + 1}.$$

Theorem 1.1 with the special case $\nu = 1$ leads us to the following:

Conjecture. *The growth order of $w_\nu(z)$ is equal to $(2\nu + 3)/(\nu + 1)$.*

Remark 1.1. Since $N(r, w_\nu) \ll n(r, w_\nu) \log r$, the quantity $N(r, w_\nu)$ in (1.5) can be replaced by $n(r, w_\nu)$.

Remark 1.2. For a meromorphic function $f(z)$, the deficiency of ∞ is defined by $\delta(\infty, f) := \liminf_{r \rightarrow \infty} (m(r, f)/T(r, f))$ ([6], [8]). For every solution $w(z)$ of (PI) , we have $\delta(\infty, w) = 0$; and this fact combined with the transcendency of $w(z)$ implies $\limsup_{r \rightarrow \infty} (N(r, w)/\log r) = \infty$ (cf. e.g. [5, §10]). For $w_\nu(z)$, analogous deficiency relations are valid: $\delta(\infty, w_\nu) = 0$, and for each $\alpha \in \mathbf{C}$, $\delta(\alpha, w_\nu) := \delta(\infty, 1/(w_\nu - \alpha)) = 0$ (cf. (3.7) and (4.1) of the proofs of Theorem 1.1 and Corollary 1.3).

Remark 1.3. The second Painlevé equation

$$(PII) \quad w'' = 2w^3 + zw + a, \quad a \in \mathbf{C}$$

belongs to the second Painlevé hierarchy $(PII_{2\nu})$ ($\nu \in \mathbf{N}$) (cf. [1], [2], [3], [7]). Value distribution properties of solutions of $(PII_{2\nu})$ are studied by Gromak and He ([4]) and by Li and He ([10]); for example, every transcendental meromorphic solution $w_{II,\nu}(z)$ satisfies $\delta(\infty, w_{II,\nu}) = 0$.

Theorem 1.1 and its corollaries are proved in Sections 3 and 4. In the proofs, we need some basic facts in the Nevanlinna theory and some properties of differential polynomials. They are reviewed or explained in Section 2. To prove Theorem 1.1, we deal with certain sums concerning the poles of each meromorphic solution, which are essential in the proof; and these sums are evaluated in the final section.

§2. Basic Facts

§2.1. Nevanlinna theory

We review basic facts in the Nevanlinna theory which are necessary in the proofs of our results (cf. [5, Appendix B], [6], [8, Chapters 1 and 2]). Let $f(z)$ be an arbitrary non-constant meromorphic function.

Lemma 2.1. *For an arbitrary $\alpha \in \mathbf{C}$, $T(r, 1/(f - \alpha)) = T(r, f) + O(1)$.*

- Lemma 2.2.** (i) $T(r, f)$ is a monotone increasing function of r .
(ii) $T(r, f) \ll \log r$ if and only if $f(z)$ is a rational function.
(iii) If $f(z)$ is transcendental, then $T(r, f)/\log r \rightarrow \infty$ as $r \rightarrow \infty$.

The following lemmas ([5, Lemmas B.11 and B.12], [8, Lemma 2.4.2 and Proposition 9.2.3]) are useful in the study of differential equations.

Lemma 2.3. *Let $f(z)$ be a non-constant meromorphic function satisfying $f^{\lambda+1} = P(z, f)$ ($\lambda \in \mathbf{N}$), where $P(z, u)$ is a polynomial in z, u and derivatives of u . Suppose that the total degree of $P(z, u)$ with respect to u and its derivatives does not exceed λ . Then $m(r, f) \ll \log T(r, f) + \log r$ as $r \rightarrow \infty$ outside an exceptional set with finite linear measure.*

Lemma 2.4. *Let $f(z)$ be a non-constant meromorphic function satisfying $F(z, f) = 0$, where $F(z, u)$ is a polynomial in z, u and derivatives of u . Suppose that $\alpha \in \mathbf{C}$ satisfies $F(z, \alpha) \not\equiv 0$. Then $m(r, 1/(f - \alpha)) \ll \log T(r, f) + \log r$ as $r \rightarrow \infty$ outside an exceptional set with finite linear measure.*

For real-valued functions, we have ([5, Lemma B.10], [8, Lemma 1,1,1])

Lemma 2.5. *Let $\phi(r)$ and $\psi(r)$ be real-valued monotone increasing functions on $(0, +\infty)$. Suppose that $\phi(r) \leq \psi(r)$ outside an exceptional set with finite linear measure. Then $\phi(r) \leq \psi(2r)$ on $(r_0, +\infty)$, where r_0 is some positive number.*

§2. Differential polynomials

For an arbitrary nonnegative integer $p \in \mathbf{N} \cup \{0\}$, set

$$[w, w', \dots, w^{(p)}]^\nu := \prod_{\kappa=0}^p (w^{(\kappa)})^{\nu_\kappa}, \quad w^{(0)} = w,$$

where $\boldsymbol{\iota} = (\iota_0, \iota_1, \dots, \iota_p) \in (\mathbf{N} \cup \{0\})^{p+1}$. For the index $\boldsymbol{\iota}$, we put

$$\|\boldsymbol{\iota}\| := \sum_{\kappa=0}^p (\kappa + 2)\iota_\kappa.$$

Consider the differential polynomial

$$\varphi[w] = \sum_{\boldsymbol{\iota} \in I_\varphi} c_\boldsymbol{\iota}[w, w', \dots, w^{(p)}]^\boldsymbol{\iota}, \quad c_\boldsymbol{\iota} \in \mathbf{C} \setminus \{0\},$$

where $I_\varphi \subset (\mathbf{N} \cup \{0\})^{p+1}$ is a finite set of indices. For $\varphi[w] (\neq 0)$, we define its weight by

$$\text{wt}(\varphi[w]) = \max\{\|\boldsymbol{\iota}\| \mid \boldsymbol{\iota} \in I_\varphi\};$$

in particular, if $\varphi[w] \equiv c_0 \in \mathbf{C} \setminus \{0\}$, then $\text{wt}(c_0) = 0$. For any integer $q > p$, the differential polynomial $\varphi[w]$ admits another expression

$$\tilde{\varphi}[w] = \sum_{\boldsymbol{\iota} \in I_\varphi} c_\boldsymbol{\iota}[w, w', \dots, w^{(p)}, \dots, w^{(q)}]^{(\boldsymbol{\iota}, \mathbf{o})}$$

with $(\boldsymbol{\iota}, \mathbf{o}) = (\iota_0, \iota_1, \dots, \iota_p, 0, \dots, 0)$. Then, $\text{wt}(\varphi[w]) = \text{wt}(\tilde{\varphi}[w])$, namely the definition of the weight is independent of the choice of the size q of the index. Let $\varphi[w] (\neq 0)$ and $\psi[w] (\neq 0)$ be arbitrary differential polynomials. Then,

$$\begin{aligned} \text{wt}(\varphi[w] + \psi[w]) &= \max\{\text{wt}(\varphi[w]), \text{wt}(\psi[w])\}, \\ \text{wt}(\varphi[w]\psi[w]) &= \text{wt}(\varphi[w]) + \text{wt}(\psi[w]). \end{aligned}$$

Remark 2.1. As will be shown later (cf. Lemma 2.7), every pole of $w_\nu(z)$ is double. Hence, for every pole of $[w_\nu, w'_\nu, \dots, w_\nu^{(2\nu)}]^\boldsymbol{\iota}$, the multiplicity of it is equal to $\|\boldsymbol{\iota}\|$. This fact is a background of the definition of the weight of differential polynomials.

We note the following:

Lemma 2.6. *For each $\nu \in \mathbf{N} \cup \{0\}$, $d_{\nu+1}[w]$ is expressible in the form*

$$(2.1) \quad d_{\nu+1}[w] = \gamma_{\nu+1}w^{\nu+1} + \sum_{\substack{\|\boldsymbol{\iota}\| \leq 2(\nu+1) \\ \iota_0 \leq \nu}} c_\boldsymbol{\iota}[w, w', \dots, w^{(2\nu)}]^\boldsymbol{\iota},$$

where

- (i) $\boldsymbol{\iota} = (\iota_0, \iota_1, \dots, \iota_{2\nu}) \in (\mathbf{N} \cup \{0\})^{2\nu+1}$,
- (ii) $\gamma_{\nu+1} \in \mathbf{C} \setminus \{0\}$, $c_\boldsymbol{\iota} \in \mathbf{C}$.

Proof. By (1.4), for every $\nu \in \mathbf{N}$,

$$\sum_{\mu=0}^{\nu} d_{\nu-\mu}[w] D d_{\mu+1}[w] = \sum_{\mu=0}^{\nu} d_{\nu-\mu}[w] (D^3 - 8wD - 4w') d_\mu[w],$$

and hence, by (1.3),

$$\begin{aligned} Dd_{\nu+1}[w] &= - \sum_{\mu=0}^{\nu-1} d_{\nu-\mu}[w] Dd_{\mu+1}[w] \\ &\quad + \sum_{\mu=0}^{\nu} (d_{\nu-\mu}[w] D^3 - 8wd_{\nu-\mu}[w] D - 4w'd_{\nu-\mu}[w]) d_{\mu}[w]. \end{aligned}$$

Substituting the identities

$$\begin{aligned} \sum_{\mu=0}^{\nu-1} d_{\nu-\mu}[w] Dd_{\mu+1}[w] &= \frac{1}{2} D \left(\sum_{\mu=0}^{\nu-1} d_{\nu-\mu}[w] d_{\mu+1}[w] \right), \\ \sum_{\mu=0}^{\nu} d_{\nu-\mu}[w] D^3 d_{\mu}[w] &= D \left(\sum_{\mu=0}^{\nu} d_{\nu-\mu}[w] D^2 d_{\mu}[w] - \frac{1}{2} \sum_{\mu=0}^{\nu} Dd_{\nu-\mu}[w] \cdot Dd_{\mu}[w] \right), \\ w \sum_{\mu=0}^{\nu} d_{\nu-\mu}[w] Dd_{\mu}[w] &= \frac{1}{2} D \left(w \sum_{\mu=0}^{\nu} d_{\nu-\mu}[w] d_{\mu}[w] \right) - \frac{w'}{2} \sum_{\mu=0}^{\nu} d_{\nu-\mu}[w] d_{\mu}[w], \end{aligned}$$

we have

$$\begin{aligned} Dd_{\nu+1}[w] &= D \left(\sum_{\mu=0}^{\nu} \left(d_{\nu-\mu}[w] D^2 - \frac{1}{2} Dd_{\nu-\mu}[w] \cdot D - 4wd_{\nu-\mu}[w] \right) d_{\mu}[w] \right) \\ &\quad - \frac{1}{2} D \left(\sum_{\mu=0}^{\nu-1} d_{\nu-\mu}[w] d_{\mu+1}[w] \right), \end{aligned}$$

which implies that

$$\begin{aligned} (2.2) \quad d_{\nu+1}[w] &= \sum_{\mu=0}^{\nu} \left(d_{\nu-\mu}[w] D^2 - \frac{1}{2} Dd_{\nu-\mu}[w] \cdot D - 4wd_{\nu-\mu}[w] \right) d_{\mu}[w] \\ &\quad - \frac{1}{2} \sum_{\mu=0}^{\nu-1} d_{\nu-\mu}[w] d_{\mu+1}[w] + C_{\nu}, \end{aligned}$$

where C_{ν} is an arbitrary constant. By (2.2) combined with (1.3), $d_{\nu+1}[w]$ is a differential polynomial. Moreover, in $d_{\nu+1}[w]$, the derivative with the highest order is $w^{(2\nu)}$. Indeed, this fact is inductively checked by using (1.3) and (1.4). Hence, for every $\nu \in \mathbf{N} \cup \{0\}$, $d_{\nu+1}[w]$ is written in the form

$$(2.3) \quad d_{\nu+1}[w] = \sum_{\iota \in I_{\nu}} \tilde{c}_{\iota}[w, w', \dots, w^{(2\nu)}]^{\iota}, \quad \iota = (\iota_0, \iota_1, \dots, \iota_{2\nu}), \quad \tilde{c}_{\iota} \in \mathbf{C} \setminus \{0\},$$

where $I_{\nu} \subset (\mathbf{N} \cup \{0\})^{2\nu+1}$ is a finite set of indices. We prove (2.1) by induction on ν . Clearly it is valid for $\nu = 0$. Suppose that (2.1) is valid for every $\nu \leq N$;

namely, $\text{wt}(d_{\nu+1}[w]) \leq 2(\nu + 1)$ for every $\nu \leq N$, and $\gamma_{N+1} \in \mathbf{C} \setminus \{0\}$. Since, for $\iota \neq \mathbf{o}$,

$$\begin{aligned} \text{wt}(D([w, w', \dots, w^{(2\nu)}]^\iota)) &= \text{wt}\left([w, w', \dots, w^{(2\nu)}]^\iota \sum_{\mu=0}^{2\nu} \iota_\mu w^{(\mu+1)} / w^{(\mu)}\right) \\ &= \max\left\{\text{wt}(w^{\iota_0} (w')^{\iota_1} \dots (w^{(2\nu)})^{\iota_{2\nu}} w^{(\mu+1)} / w^{(\mu)}) \mid 0 \leq \mu \leq 2\nu, \iota_\mu \neq 0\right\} \\ &= \max\left\{|\iota| + (\mu + 2)(-1) + (\mu + 3) \cdot 1 \mid 0 \leq \mu \leq 2\nu, \iota_\mu \neq 0\right\} = |\iota| + 1 \\ &= \text{wt}([w, w', \dots, w^{(2\nu)}]^\iota) + 1, \end{aligned}$$

we have $\text{wt}(D^l d_{\nu+1}[w]) = l + \text{wt}(d_{\nu+1}[w]) \leq l + 2(\nu + 1)$ for $\nu \leq N$ and for $l = 1, 2$. Hence, by (2.2) with $\nu = N + 1$, $\text{wt}(d_{N+2}[w]) \leq 2(N + 1) + 2 = 2(N + 2)$. Furthermore, by (1.4),

$$\begin{aligned} Dd_{N+2}[w] &= -8wD(\gamma_{N+1}w^{N+1}) - 4w' \cdot \gamma_{N+1}w^{N+1} + \dots \\ &= -4\gamma_{N+1}(2N + 3)w^{N+1}w' + \dots, \end{aligned}$$

which implies that $\gamma_{N+2} = -4(2N + 3)(N + 2)^{-1}\gamma_{N+1} \in \mathbf{C} \setminus \{0\}$. Hence, (2.1) is valid for $\nu = N + 1$ as well. This completes the proof. \square

Lemma 2.7. *For a meromorphic solution $w_\nu(z)$ of $(\text{PI}_{2\nu})$, let a_0 be an arbitrary pole of it. Then, around $z = a_0$,*

$$w_\nu(z) = c(a_0)(z - a_0)^{-2} + O(1),$$

where $c(a_0) = k(a_0)(k(a_0) + 1)/2$ for some integer $k(a_0) \in \{1, \dots, \nu\}$.

Proof: Around the pole $z = a_0$, we write $w_\nu(z) = b(z - a_0)^{-\sigma} + \dots$, $b \neq 0$. Suppose that $\sigma \geq 3$. It is inductively shown that $d_k[w_\nu](z) = b_k(z - a_0)^{-\sigma k} + \dots$, $b_k \neq 0$ for every $k \in \mathbf{N}$, because this formula with k implies

$$\begin{aligned} Dd_{k+1}[w_\nu](z) &= (D^3 - 8w_\nu(z)D - 4w'_\nu(z))d_k[w_\nu](z) \\ &= (-8(-\sigma k) - 4(-\sigma))bb_k(z - a_0)^{-\sigma(k+1)-1} + \dots \\ &= 4(2k + 1)\sigma bb_k(z - a_0)^{-\sigma(k+1)-1} + \dots, \end{aligned}$$

namely

$$d_{k+1}[w_\nu](z) = -4(2k + 1)(k + 1)^{-1}bb_k(z - a_0)^{-\sigma(k+1)} + \dots.$$

Hence, if $\sigma \geq 3$, substitution of $d_{\nu+1}[w_\nu](z) = b_{\nu+1}(z - a_0)^{-\sigma(\nu+1)} + \dots$ into $(\text{PI}_{2\nu})$ yields a contradiction. Supposing that $\sigma = 1$, by an analogous argument, we can show that $d_{\nu+1}[w_\nu](z) = b'_{\nu+1}(z - a_0)^{-(2\nu+1)} + \dots$, $b'_{\nu+1} \neq 0$, and also derive a contradiction. Therefore, $z = a_0$ is a double pole. Put $w_\nu(z) = b_0(z - a_0)^{-2} + \dots$, $b_0 \neq 0$. Since substitution of $d_k[w_\nu](z) = A_k(z - a_0)^{-2k} + \dots$, $k \in \mathbf{N}$ into (1.4) yields that $d_{k+1}[w_\nu](z) = A_{k+1}(z - a_0)^{-2(k+1)} + \dots$ with

$$A_{k+1} = -4(2k + 1)(k + 1)^{-1}(b_0 - k(k + 1)/2)A_k.$$

By this fact,

$$A_{\nu+1} = B_{\nu+1} b_0 \prod_{k=1}^{\nu} (b_0 - k(k+1)/2), \quad B_{\nu+1} \neq 0.$$

Substituting $d_{\nu+1}[w_\nu](z)$ into $(PI_{2\nu})$, we have $b_0 = k(k+1)/2$ for some $k \in \{1, \dots, \nu\}$. Furthermore, the relation

$$\begin{aligned} Dd_{\nu+2}[w_\nu](z) &= (D^3 - 8w_\nu(z)D - 4w'_\nu(z))d_{\nu+1}[w_\nu](z) \\ &= (D^3 - 8w_\nu(z)D - 4w'_\nu(z))(-4z) = 32w_\nu(z) + 16zw'_\nu(z) \\ &= 16D(zw_\nu(z)) + 16w_\nu(z), \end{aligned}$$

namely

$$16w_\nu(z) = D(d_{\nu+2}[w_\nu](z) - 16zw_\nu(z))$$

means that the residue of $w_\nu(z)$ at the pole $z = a_0$ vanishes. This completes the proof. \square

§3. Proof of Theorem 1.1

To prove (1.5), we suppose the contrary:

$$(3.1) \quad \limsup_{r \rightarrow \infty} \frac{\log N(r, w_\nu)}{\log r} < \frac{2\nu + 3}{\nu + 1},$$

namely, for some $\varepsilon > 0$, $N(r, w_\nu) \ll r^{(2\nu+3)/(\nu+1)-\varepsilon}$, from which it follows that

$$(3.2) \quad n(r) = n(r, w_\nu) \ll r^{(2\nu+3)/(\nu+1)-\varepsilon},$$

because

$$N(2r, w_\nu) \geq \int_r^{2r} (n(\rho, w_\nu) - n(0, w_\nu)) \frac{d\rho}{\rho} \geq (n(r, w_\nu) + O(1)) \log 2.$$

Starting from (3.1), we would like to derive a contradiction. Let $\{a_j\}_{j \in J}$ be a sequence of all distinct poles of $w_\nu(z)$ arranged as $|a_1| \leq \dots \leq |a_j| \leq \dots$, where $J = \mathbf{N}$ or $\{1, \dots, p\}$ ($p \in \mathbf{N}$) or \emptyset . Clearly these poles do not accumulate at any point in \mathbf{C} . By Lemma 2.7, we write $w_\nu(z)$ in the form

$$(3.3) \quad w_\nu(z) = \Phi(z) + g(z),$$

$$(3.4) \quad \Phi(z) = \sum_{j \in J} c(a_j) ((z - a_j)^{-2} - a_j^{-2}),$$

where $g(z)$ is an entire function. In (3.4), we make the following conventions: (i) if $a_1 = 0$, then the term $(z - a_1)^{-2} - a_1^{-2}$ is replaced by z^{-2} ; (ii) if $J = \emptyset$, then $\Phi(z) \equiv 0$. In what follows, we may suppose that $\Phi(z) \not\equiv 0$, because the case

where $\Phi(z) \equiv 0$ is similarly treated by adding a slight modification. Under (3.2), we have the following lemmas whose proofs will be given afterward:

Lemma 3.1. *For every $r > 1$, there exists z_r satisfying $0.7r \leq |z_r| \leq r$,*

$$\sum_{|a_j| < 2r} |z_r - a_j|^{-2} \ll r^{1/(\nu+1)-\varepsilon/2}, \quad \sum_{|a_j| < 2r} |z_r - a_j|^{-3} \ll r^{(3/2)/(\nu+1)-\varepsilon}.$$

Lemma 3.2. *Let r be an arbitrary number satisfying $r > 1$. Then,*

$$\sum_{|a_j| \geq 2r} |(z - a_j)^{-2} - a_j^{-2}| \ll r^{1/(\nu+1)-\varepsilon}, \quad \sum_{|a_j| \geq 2r} |z - a_j|^{-3} \ll 1$$

for $|z| \leq r$, and

$$\sum_{0 < |a_j| < 2r} |a_j^{-2}| \ll r^{1/(\nu+1)-\varepsilon}.$$

Lemma 3.3. *There exists a set $E \subset (0, \infty)$ with finite linear measure such that*

$$\sum_{0 < |a_j| < \infty} |(z - a_j)^{-2} - a_j^{-2}| \ll |z|^9 \quad \text{for } |z| \in (0, \infty) \setminus E.$$

By Lemma 2.6, $w_\nu(z)$ satisfies the equation

$$(3.5) \quad -\gamma_{\nu+1} w^{\nu+1} = \sum_{\substack{\|\boldsymbol{\iota}\| \leq 2(\nu+1) \\ \iota_0 \leq \nu}} c_\boldsymbol{\iota} w^{\iota_0} (w')^{\iota_1} \cdots (w^{(2\nu)})^{\iota_{2\nu}} + 4z.$$

For each term on the right-hand side, note that

$$(3.6) \quad \sum_{\kappa=0}^{2\nu} \iota_\kappa \leq \nu,$$

because $2 \sum_{\kappa=0}^{2\nu} \iota_\kappa = \|\boldsymbol{\iota}\| - \sum_{\kappa=0}^{2\nu} \kappa \iota_\kappa = 2(\nu+1)$ is valid if and only if $\boldsymbol{\iota} = (\iota_0, \iota_1, \dots, \iota_{2\nu}) = (\nu+1, 0, \dots, 0)$. By Lemma 2.3, there exists a set $E^* \subset (0, \infty)$ with finite linear measure such that

$$(3.7) \quad m(r, w_\nu) \ll \log T(r, w_\nu) + \log r$$

as $r \rightarrow \infty$, $r \notin E^*$. (Note that $(\text{PI}_{2\nu})$ does not admit a constant solution.) Combining this with (3.1), we have $T(r, w_\nu) \ll r^{(2\nu+3)/(\nu+1)}$ and $m(r, w_\nu) \ll \log r$ for $r \notin E^*$. By Lemma 3.3, for $r \notin E \cup E^*$,

$$T(r, g) = m(r, g) = m(r, w_\nu - \Phi) \leq m(r, w_\nu) + m(r, \Phi) \ll \log r.$$

By Lemmas 2.5 and 2.2, this is valid for r approaching ∞ without an exceptional set, and hence $g(z)$ is a polynomial.

By Lemmas 3.1 and 3.2, for every $r > 1$, there exists z_r , $0.7r \leq |z_r| \leq r$ satisfying

$$(3.8) \quad |\Phi(z_r)| \ll \sum_{|a_j| < 2r} |z_r - a_j|^{-2} + \sum_{0 < |a_j| < 2r} |a_j^{-2}| \\ + \sum_{|a_j| \geq 2r} |(z_r - a_j)^{-2} - a_j^{-2}| \ll r^{1/(\nu+1)-\varepsilon/2}.$$

Then, also for every $\kappa = 1, 2, \dots, 2\nu$,

$$(3.9) \quad |\Phi^{(\kappa)}(z_r)| \ll r^{(\kappa/2+1)/(\nu+1)-\varepsilon}.$$

Indeed, observing that

$$|\Phi^{(\kappa)}(z_r)| \ll \sum_{|a_j| < 2r} |z_r - a_j|^{-2-\kappa} + \sum_{|a_j| \geq 2r} |z_r - a_j|^{-2-\kappa},$$

we have the following:

(i) if κ is odd,

$$|\Phi^{(\kappa)}(z_r)| \ll \left(\sum_{|a_j| < 2r} |z_r - a_j|^{-3} \right) \left(\sum_{|a_j| < 2r} |z_r - a_j|^{-2} \right)^{(\kappa-1)/2} + \sum_{|a_j| \geq 2r} |z_r - a_j|^{-3} \\ \ll r^{(3/2)/(\nu+1)-\varepsilon} r^{(1/(\nu+1)-\varepsilon/2)(\kappa-1)/2} \ll r^{(\kappa/2+1)/(\nu+1)-\varepsilon};$$

(ii) if κ is even,

$$|\Phi^{(\kappa)}(z_r)| \ll \left(\sum_{|a_j| < 2r} |z_r - a_j|^{-2} \right)^{\kappa/2+1} + \sum_{|a_j| \geq 2r} |z_r - a_j|^{-3} \\ \ll r^{(1/(\nu+1)-\varepsilon/2)(\kappa/2+1)} \ll r^{(\kappa/2+1)/(\nu+1)-\varepsilon}.$$

From (3.5), we have

$$(3.10) \quad |w_\nu(z_r)| \ll \left(|z_r| + \sum_{\substack{\|\mathbf{u}\| \leq 2(\nu+1) \\ \iota_0 \leq \nu}} |w_\nu(z_r)^{\iota_0} w'_\nu(z_r)^{\iota_1} \dots w_\nu^{(2\nu)}(z_r)^{\iota_{2\nu}}| \right)^{1/(\nu+1)}.$$

Now suppose that $\deg g(z) = \delta_0 \geq 1$. Substitute $w_\nu^{(\kappa)}(z_r) = g^{(\kappa)}(z_r) + \Phi^{(\kappa)}(z_r)$ ($\kappa = 0, 1, \dots, 2\nu$) into (3.10), and observe that

$$|w_\nu(z_r)| \geq |g(z_r)| - |\Phi(z_r)| = |g(z_r)| + O(r^{1/(\nu+1)}) \gg r^{\delta_0},$$

and that, for every \mathbf{u} satisfying $\iota_0 \leq \nu$ and $\|\mathbf{u}\| \leq 2(\nu+1)$,

$$|w_\nu(z_r)^{\iota_0} w'_\nu(z_r)^{\iota_1} \dots w_\nu^{(2\nu)}(z_r)^{\iota_{2\nu}}| \\ \ll (|g(z_r)| + |\Phi(z_r)|)^{\iota_0} (|g(z_r)| + |\Phi'(z_r)|)^{\iota_1} \dots (|g(z_r)| + |\Phi^{(2\nu)}(z_r)|)^{\iota_{2\nu}} \\ \ll |z_r|^{\delta_0(\iota_0 + \iota_1 + \dots + \iota_{2\nu})} \ll r^{\delta_0 \nu}$$

(cf. (3.6), (3.8) and (3.9)). Then, we have the contradiction $r^{\delta_0} \ll (|z_r| + r^{\delta_0\nu})^{1/(\nu+1)} \ll r^{\delta_0\nu/(\nu+1)}$; which implies that $g(z) \equiv C \in \mathbf{C}$. Substituting $w = w_\nu(z) = \Phi(z) + C$ into $(\text{PI}_{2\nu})$, and observing that, for every ι satisfying $0 < \|\iota\| \leq 2(\nu + 1)$,

$$\begin{aligned} & |w_\nu(z_r)^{\iota_0} w'_\nu(z_r)^{\iota_1} \cdots w_\nu^{(2\nu)}(z_r)^{\iota_{2\nu}}| \\ & \ll (|\Phi(z_r)| + |C|)^{\iota_0} |\Phi'(z_r)|^{\iota_1} \cdots |\Phi^{(2\nu)}(z_r)|^{\iota_{2\nu}} \ll r^{\chi(\iota) - \varepsilon/2} \end{aligned}$$

with $\chi(\iota) = \sum_{\kappa=0}^{2\nu} (\kappa/2 + 1)\iota_\kappa / (\nu + 1) = (\|\iota\|/2) / (\nu + 1) \leq 1$ (cf. (3.8) and (3.9)), we have

$$0.7r \leq |z_r| \ll |d_{\nu+1}[w_\nu](z_r)| \ll r^{1-\varepsilon/2},$$

which is a contradiction. We have thus proved (1.5).

§4. Proofs of Corollaries 1.2 and 1.3

Corollary 1.2 immediately follows from Theorem 1.1. To prove Corollary 1.3, note that $w \equiv \alpha \in \mathbf{C}$ is not a solution of $(\text{PI}_{2\nu})$. By Lemma 2.4,

$$(4.1) \quad m(r, 1/(w_\nu - \alpha)) \ll \log T(r, w_\nu) + \log r$$

as $r \rightarrow \infty$ for $r \notin E_1$, where $E_1 \subset (0, \infty)$ is a set with finite linear measure. Since $w_\nu(z)$ is transcendental, by Lemmas 2.1 and 2.2, we have

$$\frac{N(r, 1/(w_\nu - \alpha))}{T(r, w_\nu)} = 1 - \frac{m(r, 1/(w_\nu - \alpha)) + O(1)}{T(r, w_\nu)} \rightarrow 1 \quad \text{as } r \rightarrow \infty, r \notin E_1;$$

and hence

$$(4.2) \quad N(r, 1/(w_\nu - \alpha)) \geq (1/2)T(r, w_\nu) \quad \text{for } r \in (r_1, \infty) \setminus E_1,$$

for some $r_1 > 0$. On the other hand, by (3.7), $N(r, w_\nu)/T(r, w_\nu) \rightarrow 1$ as $r \rightarrow \infty$, $r \notin E^*$. Hence,

$$(4.3) \quad (1/2)N(r, w_\nu) \leq T(r, w_\nu) \quad \text{for } r \in (r_2, \infty) \setminus E^*,$$

for some $r_2 > 0$. Using Lemma 2.5, from (4.2) and (4.3), we derive that $N(r, w_\nu) \leq 4N(2r, 1/(w_\nu - \alpha))$ for $r \in (r_3, \infty)$, where $r_3 > 0$ is sufficiently large. This inequality combined with (1.5) yields the conclusion (1.6) of Corollary 1.3.

§5. Proofs of Lemmas 3.1, 3.2 and 3.3

5.1. Proof of Lemma 3.1

Put $D_r = \{z \mid |z| \leq r\}$ and $\Delta_0 = \mathbf{C} \setminus (\bigcup_{j \geq 1} U_j)$; where $U_j = \{z \mid |z - a_j| < |a_j|^{-(1/2)/(\nu+1)}\}$ if $a_j \neq 0$, and $U_1 = \{z \mid |z| < 1\}$ if $a_1 = 0$. Since, by (3.2),

$$\begin{aligned} \sum_{1 < |a_j| < r} |a_j|^{-1/(\nu+1)} &= \int_1^r \rho^{-1/(\nu+1)} dn(\rho) \\ &= \left[\rho^{-1/(\nu+1)} n(\rho) \right]_1^r + \frac{1}{(\nu+1)} \int_1^r \rho^{-1-1/(\nu+1)} n(\rho) d\rho \ll r^{2-\varepsilon}, \end{aligned}$$

we can take r_0 so large that $7\pi r^2/8 \leq \mu(\Delta_0 \cap D_r) < \pi r^2$ for every $r > r_0$, where $\mu(X)$ denotes the area of a set X . For every $r > 1$, if $|a_j| < 2r$, then

$$\iint_{D_r \setminus U_j} |z - a_j|^{-2} dx dy \leq \iint_{\substack{|a_j|^{-(1/2)/(\nu+1)} \leq \rho \leq 3r \\ 0 \leq \theta \leq 2\pi}} \rho^{-1} d\rho d\theta \ll \log r,$$

and

$$\iint_{D_r \setminus U_j} |z - a_j|^{-3} dx dy \leq \iint_{\substack{|a_j|^{-(1/2)/(\nu+1)} \leq \rho \leq 3r \\ 0 \leq \theta \leq 2\pi}} \rho^{-2} d\rho d\theta \ll r^{(1/2)/(\nu+1)}.$$

Hence,

$$(5.1) \quad \iint_{\Delta_0 \cap D_r} \sum_{|a_j| < 2r} |z - a_j|^{-2} dx dy \ll n(2r) \log r \leq K_0 r^{(2\nu+3)/(\nu+1)-\varepsilon/2},$$

$$(5.2) \quad \iint_{\Delta_0 \cap D_r} \sum_{|a_j| < 2r} |z - a_j|^{-3} dx dy \ll n(2r) r^{(1/2)/(\nu+1)} \leq K_0 r^{2+(3/2)/(\nu+1)-\varepsilon},$$

where K_0 is some positive number. Consider the sets

$$\begin{aligned} F_r^1 &= \left\{ z \in \Delta_0 \cap D_r \mid \sum_{|a_j| < 2r} |z - a_j|^{-2} \leq 8\pi^{-1} K_0 r^{1/(\nu+1)-\varepsilon/2} \right\}, \\ F_r^2 &= \left\{ z \in \Delta_0 \cap D_r \mid \sum_{|a_j| < 2r} |z - a_j|^{-3} \leq 8\pi^{-1} K_0 r^{(3/2)/(\nu+1)-\varepsilon} \right\}. \end{aligned}$$

Suppose that $\mu(F_r^1) < 3\pi r^2/4$. Then

$$\begin{aligned} \iint_{\Delta_0 \cap D_r \setminus F_r^1} \sum_{|a_j| < 2r} |z - a_j|^{-2} dx dy \\ > 8\pi^{-1} K_0 r^{1/(\nu+1)-\varepsilon/2} (7/8 - 3/4) \pi r^2 = K_0 r^{(2\nu+3)/(\nu+1)-\varepsilon/2}, \end{aligned}$$

which contradicts (5.1). This implies that $\mu(F_r^1) \geq 3\pi r^2/4$. By the same argument, we have $\mu(F_r^2) \geq 3\pi r^2/4$. Hence, $\mu(F_r^1 \cap F_r^2) \geq \pi r^2/2$. Observing that $\mu(\{z \mid |z| < 0.7r\}) = 0.49\pi r^2$, we have $\{z \mid 0.7r \leq |z| \leq r\} \cap (F_r^1 \cap F_r^2) \neq \emptyset$, which implies the conclusion.

§5.2. Proof of Lemma 3.2

For $|a_j| \geq 2r$, and for $z \in D_r$, observing that $|z/a_j| \leq 1/2$, we have $|z - a_j|^{-3} \leq 8|a_j|^{-3}$, and

$$|(z - a_j)^{-2} - a_j^{-2}| = 2|z||a_j|^{-3}|1 - (z/a_j)/2||1 - z/a_j|^{-2} \leq 10r|a_j|^{-3}.$$

Hence, by (3.2),

$$\begin{aligned} \sum_{|a_j| \geq 2r} |(z - a_j)^{-2} - a_j^{-2}| &\ll r \sum_{|a_j| \geq 2r} |a_j|^{-3} \ll r \int_{2r}^{\infty} t^{-3} dn(t) \\ &\ll r \int_{2r}^{\infty} t^{-4} n(t) dt \ll r^{1/(\nu+1)-\varepsilon}, \end{aligned}$$

$$\sum_{|a_j| \geq 2r} |z - a_j|^{-3} \ll \sum_{|a_j| \geq 2r} |a_j|^{-3} \ll \int_{2r}^{\infty} t^{-3} dn(t) \ll 1,$$

and

$$\begin{aligned} \sum_{0 < |a_j| < 2r} |a_j^{-2}| &\ll \int_1^{2r} t^{-2} dn(t) + O(1) \\ &\ll r^{1/(\nu+1)-\varepsilon} + \int_1^{2r} t^{-3} n(t) dt + O(1) \ll r^{1/(\nu+1)-\varepsilon}. \end{aligned}$$

Thus the lemma is proved.

§5.3. Proof of Lemma 3.3

We put

$$E = (0, |a_1| + 1) \cup \left(\bigcup_{j \in J \setminus \{1\}} (|a_j| - |a_j|^{-3}, |a_j| + |a_j|^{-3}) \right).$$

Since, by (3.2),

$$\sum_{j \in J \setminus \{1\}} |a_j|^{-3} \ll \int_1^{\infty} t^{-3} dn(t) + O(1) \ll \int_1^{\infty} t^{-4} n(t) dt + O(1) \ll 1,$$

the total length of E is finite. If $|z| \notin E$, then

$$\left(\sum_{0 < |a_j| < 2|z|} + \sum_{|a_j| \geq 2|z|} \right) |(z - a_j)^{-2} - a_j^{-2}| \ll (|z|^6 + 1)n(2|z|) + |z|^{1/(\nu+1)} \ll |z|^9.$$

This completes the proof.

References

- [1] Airault, H., Rational solutions of Painlevé equations, *Stud. Appl. Math.*, **61** (1979), 31–53 .
- [2] Flaschka, H. and Newell, A. C., Monodromy- and spectrum-preserving deformations, I, *Commun. Math. Phys.*, **76** (1980), 65–116.
- [3] Gromak, V. I., Nonlinear evolution equations and equations of P -type, *Differential Equations*, **20** (1984), 1435–1439.
- [4] Gromak, V. I. and He, Y., On the solutions of the second Painlevé equation of higher order, *Proc. Math. Inst. Belarus Natl. Acad. Sci.*, **4** (2000), 37–48.
- [5] Gromak, V. I., Laine, I. and Shimomura, S., *Painlevé Differential Equations in the Complex Plane*, Walter de Gruyter, Berlin, New York 2002.
- [6] Hayman, W. K., *Meromorphic Functions*, Clarendon Press, Oxford, 1964.
- [7] Kudryashov, N. A., The first and second Painlevé equations of higher order and some relations between them, *Phys. Lett. A*, **224** (1997), 353–360.
- [8] Laine, I., *Nevanlinna Theory and Complex Differential Equations* Walter de Gruyter, Berlin, New York, 1993.
- [9] Lax, P. D., Almost periodic solutions of the KdV equation, *SIAM Rev.*, **18** (1976), 351–375.
- [10] Li, Y. and He, Y., On analytic properties of higher analogs of the second Painlevé equation, *J. Math. Phys.*, **43** (2002), 1106–1115.
- [11] Mues, E. and Redheffer, R., On the growth of logarithmic derivatives, *J. London Math. Soc.*, **8** (1974), 412–425.
- [12] Shimomura, S., Growth of the first, the second and the fourth Painlevé transcendents, *Math. Proc. Cambridge Philos. Soc.*, **134** (2003), 259–269.
- [13] Shimomura, S., Lower estimates for the growth of Painlevé transcendents, *Funkcial. Ekvac.* **46** (2003), 287–295.
- [14] Steinmetz, N., Value distribution of Painlevé transcendents, *Israel J. Math.*, **128** (2002), 29–52.
- [15] Weiss, J., On classes of integrable systems and the Painlevé property, *J. Math. Phys.*, **25** (1984), 13–24.