

Iwasawa theory for Artin representations I

Ralph Greenberg & Vinayak Vatsal

In honor of Professor Kenkichi Iwasawa

Abstract.

We introduce a natural way to define Selmer groups and p -adic L -functions for modular forms of weight 1. The corresponding Galois representation ρ of $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ is a 2-dimensional Artin representation with odd determinant. Thus, the dimension d^+ of the $(+1)$ -eigenspace for complex conjugation is 1. Choose a prime p such that the restriction of ρ to the local Galois group $\text{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)$ has a 1-dimensional constituent ε with multiplicity 1. If we fix the choice of such an ε , we can define a Selmer group and a p -adic L -function. On the algebraic side, we prove that the Selmer group over the cyclotomic \mathbf{Z}_p -extension of \mathbf{Q} is a cotorsion module over the Iwasawa algebra Λ . That result is valid for an Artin representation of arbitrary dimension d under the assumption that $d^+ = 1$ and that such an ε can be chosen. On the analytic side, the corresponding complex L -function has no critical values and the definition of the p -adic L -function depends on deforming the Galois representation ρ by Hida theory.

§1. Introduction

Suppose that K is a finite Galois extension of \mathbf{Q} . Let Δ denote the Galois group $\text{Gal}(K/\mathbf{Q})$. Consider an irreducible representation

$$\rho: \Delta \rightarrow GL_d(\overline{\mathbf{Q}}),$$

where $\overline{\mathbf{Q}}$ is an algebraic closure of \mathbf{Q} and $d \geq 1$. Let χ denote the character of ρ . If we fix an embedding σ_∞ of $\overline{\mathbf{Q}}$ into \mathbf{C} , then we obtain a

Received March 27, 2018.

Revised August 12, 2019.

2010 *Mathematics Subject Classification.* 11F11, 11R23, 11R34.

Key words and phrases. Iwasawa theory, Artin representations.

d -dimensional representation of Δ over \mathbf{C} . One can then define the corresponding Artin L -function $L_\infty(s, \chi)$. It is defined by an Euler product for $\operatorname{Re}(s) > 1$ and can be analytically continued as a meromorphic function on the complex plane. Conjecturally, $L_\infty(s, \chi)$ is an entire function if χ is nontrivial. We will assume that χ is nontrivial throughout this paper.

Now suppose that p is a prime and that we fix an embedding σ_p of $\overline{\mathbf{Q}}$ into $\overline{\mathbf{Q}_p}$, an algebraic closure of the field \mathbf{Q}_p of p -adic numbers. Then ρ becomes a d -dimensional representation of Δ over $\overline{\mathbf{Q}_p}$. Its character χ now has values in $\overline{\mathbf{Q}_p}$. One can ask whether there is a natural way to define a function $L_p(s, \chi)$ attached to ρ and the fixed embedding, where s is now a p -adic variable.

This question has been studied previously when K is totally real. In that case, the values of $L_\infty(s, \chi)$ at negative odd integers are nonzero algebraic numbers. Using the fixed embeddings σ_∞ and σ_p , one can regard those values as elements of $\overline{\mathbf{Q}}$ and then as elements of $\overline{\mathbf{Q}_p}$. Such a p -adic L -function $L_p(s, \chi)$ can then be defined by an interpolation property involving those elements of $\overline{\mathbf{Q}_p}$. First of all, if $d = 1$, then χ can be identified with an even Dirichlet character and the construction of such a function $L_p(s, \chi)$ was done by Kubota and Leopoldt [24]. If $d > 1$, then there is a definition described in [11] which is based on the Deligne-Ribet construction of p -adic L -functions for Hecke characters of the intermediate fields for the extension K/\mathbf{Q} given in [6]. Furthermore, still assuming that K is totally real, one can give a precise interpretation of the zeros of $L_p(s, \chi)$. This is a generalization of a famous conjecture of Iwasawa (for the case where $d = 1$) and has been proven by Wiles [34].

Let v denote an archimedean prime of K and let K_v denote the v -adic completion of K , which we identify with either \mathbf{R} or \mathbf{C} . As usual, we identify $\operatorname{Gal}(K_v/\mathbf{R})$ with a subgroup Δ_v of Δ . Let d^+ denote the multiplicity of the trivial character of Δ_v in $\rho|_{\Delta_v}$. This does not depend on the choice of v . Of course, if K is totally real, then $d^+ = d$. Our main objective in this paper is to study the case where $d^+ = 1$ and $d \geq 1$. Under certain assumptions, we will define certain p -adic L -functions and discuss a conjectural interpretation of the zeros of those functions. Our theory reduces to the classical case where $d^+ = d$ when $d = 1$.

We will usually make the following three-part assumption. In the last part, $\Delta_{\mathfrak{p}}$ denotes the decomposition subgroup of Δ for a prime \mathfrak{p} lying over p .

Hypothesis A: *The degree $[K : \mathbf{Q}]$ is not divisible by p . Also, $d^+ = 1$. Furthermore, there exists a 1-dimensional representation $\varepsilon_{\mathfrak{p}}$ of $\Delta_{\mathfrak{p}}$ which occurs with multiplicity 1 in $\rho|_{\Delta_{\mathfrak{p}}}$.*

Our theory depends on choosing one such character $\varepsilon_{\mathfrak{p}}$ of $\Delta_{\mathfrak{p}}$. Also, if $d > 1$, then K must be totally complex and hence of even degree. We will therefore assume that p is odd from here on.

Since $d^+ = 1$, Frobenius Reciprocity implies that ρ occurs with multiplicity 1 in the \mathbf{Q} -representation of Δ induced from the trivial representation of Δ_v for any archimedean prime v of K . It follows that ρ itself can be realized by matrices with entries in $\mathbf{Q}(\chi)$, the field generated by the values of χ . Let \mathcal{F} be the field generated over \mathbf{Q}_p by the values of χ and of $\varepsilon_{\mathfrak{p}}$. We can then regard ρ as a representation of Δ over \mathcal{F} (via σ_p). Let V denote the underlying representation space. Thus, $\dim_{\mathcal{F}}(V) = d$. Also, the last part of Hypothesis A means that the maximal \mathcal{F} -subspace $V^{(\varepsilon_{\mathfrak{p}})}$ of V on which $\Delta_{\mathfrak{p}}$ acts by the character $\varepsilon_{\mathfrak{p}}$ has dimension 1.

The theory described in this paper depends not just on χ , but also on the choice of the character $\varepsilon_{\mathfrak{p}}$ of $\Delta_{\mathfrak{p}}$ satisfying Hypothesis A. Suppose that \mathfrak{p}' is another prime of K lying over p . Then $\mathfrak{p}' = \delta(\mathfrak{p})$ for some $\delta \in \Delta$. Thus, $\Delta_{\mathfrak{p}'} = \delta^{-1}\Delta_{\mathfrak{p}}\delta$. Conjugation by δ^{-1} defines an isomorphism $\Delta_{\mathfrak{p}'} \rightarrow \Delta_{\mathfrak{p}}$. One then obtains a character $\varepsilon_{\mathfrak{p}'}$ of $\Delta_{\mathfrak{p}'}$ from $\varepsilon_{\mathfrak{p}}$ by composing with this isomorphism. One sees easily that $\varepsilon_{\mathfrak{p}'}$ is well-defined and occurs with multiplicity 1 in $\rho|_{\Delta_{\mathfrak{p}'}}$. In effect, we have a family of characters $\{\varepsilon_{\mathfrak{p}}\}_{\mathfrak{p}|p}$ for the decomposition subgroups which are compatible under conjugation.

Let \mathcal{K} denote the completion of K at \mathfrak{p} . Then \mathcal{K} is a certain finite Galois extension of \mathbf{Q}_p . Consider the restriction map $\text{Gal}(\mathcal{K}/\mathbf{Q}_p) \rightarrow \Delta_{\mathfrak{p}}$. It is an isomorphism. Composing this map with the character $\varepsilon_{\mathfrak{p}}$ of $\Delta_{\mathfrak{p}}$ determines a character ε of $\text{Gal}(\mathcal{K}/\mathbf{Q}_p)$. It has values in \mathcal{F} . If \mathfrak{p}' is another prime of K lying over p , then the completion $\mathcal{K}_{\mathfrak{p}'}$ is isomorphic to \mathcal{K} . The character ε then determines a character of $\Delta_{\mathfrak{p}'}$. In fact, this character does not depend on the isomorphism and is just $\varepsilon_{\mathfrak{p}'}$ as defined above. Thus, the family $\{\varepsilon_{\mathfrak{p}}\}_{\mathfrak{p}|p}$ is determined by the single character ε of $\text{Gal}(\mathcal{K}/\mathbf{Q}_p)$.

We will first describe the algebraic side of our theory. We assume that Hypothesis A is satisfied. Let \mathcal{O} denote the ring of integers in \mathcal{F} . Let T denote an \mathcal{O} -lattice in V which is invariant under the action of Δ . Since $|\Delta|$ is prime to p and ρ is irreducible over \mathcal{F} , T is determined up to multiplication by an element of \mathcal{F}^{\times} . Let $D = V/T$. Thus, D is isomorphic to $(\mathcal{F}/\mathcal{O})^d$ as an \mathcal{O} -module and has an \mathcal{O} -linear action of Δ . Furthermore, for every prime \mathfrak{p} of K lying above p , we will denote

the image of $V^{(\varepsilon_{\mathfrak{p}})}$ in D by $D^{(\varepsilon_{\mathfrak{p}})}$. It is the maximal $\Delta_{\mathfrak{p}}$ -invariant \mathcal{O} -submodule of D on which $\Delta_{\mathfrak{p}}$ acts by $\varepsilon_{\mathfrak{p}}$ and is isomorphic to \mathcal{F}/\mathcal{O} as an \mathcal{O} -module.

We now define a “Selmer group” associated to χ and ε which we denote by $S_{\chi, \varepsilon}(\mathbf{Q})$. It will be a certain subgroup of $H^1(G_{\mathbf{Q}}, D)$, where $G_{\mathbf{Q}} = \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$. We will write $H^1(\mathbf{Q}, D)$ instead of $H^1(G_{\mathbf{Q}}, D)$, as usual. A similar notation will be used for other fields and Galois modules. Suppose that ℓ is a prime. Let $\overline{\mathbf{Q}}_{\ell}$ be an algebraic closure of \mathbf{Q}_{ℓ} . Pick (arbitrarily) an embedding of $\overline{\mathbf{Q}}$ into $\overline{\mathbf{Q}}_{\ell}$. We can then identify $G_{\mathbf{Q}_{\ell}} = \text{Gal}(\overline{\mathbf{Q}}_{\ell}/\mathbf{Q}_{\ell})$ with a subgroup of $G_{\mathbf{Q}}$. Let \mathbf{Q}_{ℓ}^{unr} be the maximal unramified extension of \mathbf{Q}_{ℓ} in $\overline{\mathbf{Q}}_{\ell}$. The inertia subgroup of $G_{\mathbf{Q}_{\ell}}$ is $G_{\mathbf{Q}_{\ell}^{unr}}$ and is also identified with a subgroup of $G_{\mathbf{Q}}$.

Note that if $\ell = p$, then the chosen embedding of $\overline{\mathbf{Q}}$ into $\overline{\mathbf{Q}}_p$ determines a prime of K lying above p , which we denote simply by \mathfrak{p} . Our Selmer group is defined by

$$S_{\chi, \varepsilon}(\mathbf{Q}) = \ker \left(H^1(\mathbf{Q}, D) \rightarrow \prod_{\ell \neq p} H^1(\mathbf{Q}_{\ell}^{unr}, D) \times H^1(\mathbf{Q}_p^{unr}, D/D^{(\varepsilon_{\mathfrak{p}})}) \right).$$

The “global-to-local” map occurring in this definition is induced by the restriction maps, where we identify the $G_{\mathbf{Q}_{\ell}^{unr}}$ ’s with subgroups of $G_{\mathbf{Q}}$ as above. However, as our notation suggests, $S_{\chi, \varepsilon}(\mathbf{Q})$ is completely determined by the \mathcal{F} -valued character χ of Δ (of degree d) and by the \mathcal{F} -valued character ε of $\text{Gal}(K/\mathbf{Q}_p)$ (of degree 1). It does not depend on the chosen embeddings and identifications.

The above definition of our Selmer group is suggested by the general notion of a Selmer group which was studied in [12] and which also depends on having a filtration on a representation space V for $G_{\mathbf{Q}}$ when restricted to the local Galois group $G_{\mathbf{Q}_p}$. In that paper, the local condition at the prime p is defined in terms of a certain subspace denoted by F^+V . That paper primarily considers the so-called p -critical case in which the dimension of F^+V is equal to d^+ . And so, if $d^+ = 1$, then F^+V should be of dimension 1. In the definition given above, the role of F^+V is now being played by $V^{(\varepsilon_{\mathfrak{p}})}$.

Now $S_{\chi, \varepsilon}(\mathbf{Q})$ is a subgroup of $H^1(\mathbf{Q}, D)$. Since D is an \mathcal{O} -module, so is $H^1(\mathbf{Q}, D)$. It is clear that $S_{\chi, \varepsilon}(\mathbf{Q})$ is an \mathcal{O} -submodule. The restriction map

$$H^1(\mathbf{Q}, D) \rightarrow H^1(K, D)^{\Delta} = \text{Hom}_{\Delta}(\text{Gal}(K^{ab}/K), D)$$

is an isomorphism. Here K^{ab} is the maximal abelian extension of K in $\overline{\mathbf{Q}}$. Also, the notation $\text{Hom}_{\Delta}(\cdot, \cdot)$ will always denote the group of continuous, Δ -equivariant homomorphisms. The above isomorphism follows

from the assumption that $|\Delta|$ is not divisible by p and the fact that G_K acts trivially on D . Thus, we can identify $S_{\chi,\varepsilon}(\mathbf{Q})$ with its image under the restriction map which we will now describe.

Let $\xi \in \text{Hom}_\Delta(\text{Gal}(K^{ab}/K), D)$. Let K_ξ be the fixed field for $\ker(\xi)$. Thus, K_ξ is a finite, abelian extension of K , Galois over \mathbf{Q} , and ξ defines a Δ -equivariant isomorphism of $\text{Gal}(K_\xi/K)$ to a subgroup of D . For ξ to be in the image of $S_{\chi,\varepsilon}(\mathbf{Q})$, the local conditions defining $S_{\chi,\varepsilon}(\mathbf{Q})$ are equivalent to the following conditions on ξ . First of all, the extension K_ξ/K is unramified at all primes of K not dividing p . Furthermore, if \mathfrak{p} is a prime of K lying over p , then the image of the corresponding inertia subgroup of $\text{Gal}(K_\xi/K)$ under the map ξ is contained in $D^{(\varepsilon_{\mathfrak{p}})}$. By using the above identification, we will prove the following result.

Theorem 1. $S_{\chi,\varepsilon}(\mathbf{Q})$ is finite.

The proof of this theorem uses the Baker-Brumer Theorem concerning the linear independence over $\overline{\mathbf{Q}}$ of the p -adic logarithms of algebraic numbers and is reminiscent of the proof of Leopoldt's conjecture for abelian extensions of \mathbf{Q} . Leopoldt's conjecture is usually formulated as a statement about the p -adic independence of units in a number field K . But, if K is Galois over \mathbf{Q} , then it has an equivalent formulation as a statement about the ρ -components of the unit group, where ρ varies over the absolutely irreducible Artin representations of $\text{Gal}(K/\mathbf{Q})$. As we will show in Section 2, the Baker-Brumer Theorem provides a proof of that conjecture when $d^+ = 1$. However, Theorem 1 is a stronger statement and not just a consequence of Leopoldt's conjecture for such a ρ . The proof requires a somewhat more refined application of the Baker-Brumer Theorem and depends on Hypothesis A.

Let \mathbf{Q}_∞ be the cyclotomic \mathbf{Z}_p -extension of \mathbf{Q} . For $n \geq 0$, let \mathbf{Q}_n denote the subfield of \mathbf{Q}_∞ of degree p^n over \mathbf{Q} . One can define Selmer groups $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ for all $n \geq 1$ and $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ in a similar way to the definition of $S_{\chi,\varepsilon}(\mathbf{Q})$ (the case $n = 0$). We will actually prove that $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ is finite for all $n \geq 0$. Let $\Gamma = \text{Gal}(\mathbf{Q}_\infty/\mathbf{Q})$ and let $\Lambda_{\mathcal{O}} = \mathcal{O}[[\Gamma]]$. Now Γ acts naturally on $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ and the action is \mathcal{O} -linear. Thus, we can regard $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ as a discrete $\Lambda_{\mathcal{O}}$ -module. Let $X_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ denote the Pontryagin dual of $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$. We can regard $X_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ as a compact $\Lambda_{\mathcal{O}}$ -module. The following result is a straightforward consequence of the finiteness of the $S_{\chi,\varepsilon}(\mathbf{Q}_n)$'s.

Theorem 2. The $\Lambda_{\mathcal{O}}$ -module $X_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is finitely-generated and torsion.

Alternatively, we would say that $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is a cofinitely-generated, co-torsion $\Lambda_{\mathcal{O}}$ -module.

We will next state a result on the algebraic side concerning the characteristic ideal of $X_{\chi,\varepsilon}(\mathbf{Q}_\infty)$. We will prove this result in a sequel to this paper (part II). By definition, the characteristic ideal is a principal ideal in $\Lambda_{\mathcal{O}}$. Assume that $d = 2$. We will then prove that the characteristic ideal has a generator with an interpolation property of a certain form. To simplify the statement and explanation, we will assume here that $\mathcal{F} = \mathbf{Q}_p$. Thus, we are assuming that both χ and ε have values in \mathbf{Q}_p .

Let $K_\infty = K\mathbf{Q}_\infty$, the cyclotomic \mathbf{Z}_p -extension of K . We can identify $\text{Gal}(K_\infty/K)$ with $\Gamma = \text{Gal}(\mathbf{Q}_\infty/\mathbf{Q})$ by the obvious restriction map, noting that $K \cap \mathbf{Q}_\infty = \mathbf{Q}$ because $[K : \mathbf{Q}]$ is prime to p . We may assume that K contains μ_p . Then $K_\infty = K(\mu_{p^\infty})$. The action of $\text{Gal}(K_\infty/K)$ on μ_{p^∞} defines an isomorphism $\kappa : \Gamma \rightarrow 1 + p\mathbf{Z}_p$ (since p is odd). Recall that \mathcal{K} denotes the completion of K at any one of the primes of K lying above p . We let \mathcal{K}_∞ denote the cyclotomic \mathbf{Z}_p -extension of \mathcal{K} . We can also identify $\text{Gal}(\mathcal{K}_\infty/\mathcal{K})$ with Γ . For $n \geq 0$, the n -th layers in K_∞/K and in $\mathcal{K}_\infty/\mathcal{K}$ will be denoted by K_n and \mathcal{K}_n , respectively. We also will identify $\text{Gal}(K_\infty/\mathbf{Q}_\infty)$ with Δ by the restriction map. We then have a canonical isomorphism $\text{Gal}(K_\infty/\mathbf{Q}) \cong \Delta \times \Gamma$. Similarly, $\text{Gal}(\mathcal{K}_\infty/\mathbf{Q}_p) \cong \mathcal{G} \times \Gamma$, where $\mathcal{G} = \text{Gal}(\mathcal{K}/\mathbf{Q}_p)$. We will denote $\text{Gal}(K_n/\mathbf{Q})$ by Δ_n .

For each $n \geq 0$, let \mathcal{U}_n denote the group of principal units in \mathcal{K}_n . Let U_n denote the group of units in K_n which are principal in all the completions of K_n at primes above p . Thus, we can pick embeddings $K_n \rightarrow \mathcal{K}_n$ in a compatible way and thereby obtain natural injective homomorphisms $U_n \rightarrow \mathcal{U}_n$. The lack of uniqueness will not affect the statement of the next theorem. Those homomorphisms commute with the natural global and local norm maps $U_m \rightarrow U_n$ and $\mathcal{U}_m \rightarrow \mathcal{U}_n$ for $m \geq n \geq 0$.

Since we are regarding \mathcal{G} as a subgroup of $\text{Gal}(\mathcal{K}_\infty/\mathbf{Q}_p)$, and the \mathcal{U}_n 's are abelian pro- p groups and hence \mathbf{Z}_p -modules, we can consider the \mathcal{U}_n 's as $\mathbf{Z}_p[\mathcal{G}]$ -modules. For each $n \geq 0$, let $\mathcal{U}_n^{(\varepsilon)}$ denote the ε -component of \mathcal{U}_n . We can then define the projection map $\mathcal{U}_n \rightarrow \mathcal{U}_n^{(\varepsilon)}$, which is a surjective Γ -equivariant map. We will use the somewhat peculiar notation $|\cdot|_\varepsilon$ for this projection map. Our motivation for this notation is to suggest an analogy with the classical Stark conjecture which involves the complex log of the absolute value of so-called Stark units. Theorem 3 below involves instead the p -adic log of $|\cdot|_\varepsilon$ applied to certain units.

Suppose that $\varphi : \Gamma \rightarrow \overline{\mathbf{Q}}_p^\times$ is a continuous group homomorphism. Then we can extend φ to a continuous \mathbf{Z}_p -algebra homomorphism from

Λ to $\overline{\mathbf{Q}}_p$, which we also denote by φ . Thus, its image $\varphi(\Lambda)$ is a compact subring of some finite extension of \mathbf{Q}_p . The elements of $\text{Hom}_{\text{cont}}(\Gamma, \overline{\mathbf{Q}}_p^\times)$ of finite order will be of particular interest. If φ has order p^n , then $\varphi(\Lambda) = \mathbf{Z}_p[\mu_{p^n}]$. One can also consider the elements of $\text{Hom}_{\text{cont}}(\Gamma, \overline{\mathbf{Q}}_p^\times)$ of the form κ^t , where $t \in \mathbf{Z}_p$. They have values in $1 + p\mathbf{Z}_p$ and we have $\kappa^t(\Lambda) = \mathbf{Z}_p$.

Now ε factors through the Galois group of a cyclic extension of \mathbf{Q}_p of degree prime to p . We have a canonical factorization $\varepsilon = \omega^a \beta$, where $0 \leq a < p - 1$ and β is unramified. Let b denote the order of β . Thus, β factors through $\text{Gal}(\mathbf{Q}_p(\mu_f)/\mathbf{Q}_p)$ where $f = p^b - 1$ and is a faithful character of that Galois group. Also, ω denotes the Teichmüller character and factors through $\text{Gal}(\mathbf{Q}_p(\mu_p)/\mathbf{Q}_p)$. If φ is a character of Γ of order p^n , then $\omega^a \varphi$ factors through $\text{Gal}(\mathbf{Q}_p(\mu_{p^{n+1}})/\mathbf{Q}_p)$. Thus, $\varepsilon \varphi$ can be regarded as a character of $\text{Gal}(\mathbf{Q}_p(\mu_{fp^{n+1}})/\mathbf{Q}_p)$. We choose a generator ζ_f for μ_f so that its image $\tilde{\zeta}_f$ in the residue field \mathbf{F}_{p^b} for $\mathbf{Q}_p(\mu_f)$ is part of a normal basis for $\mathbf{F}_{p^b}/\mathbf{F}_p$. Of course, this just means that $\{\tilde{\zeta}_f^{p^j}\}_{0 \leq j < b}$ is linearly independent over \mathbf{F}_p . The existence of such an element ζ_f is a theorem of Lenstra and Schoof (in [25]). For $n \geq 0$, we choose generators $\zeta_{fp^{n+1}}$ for $\mu_{fp^{n+1}}$ such that $\zeta_{fp^{n+1}}^p = \zeta_{fp^n}$. We define a “Gaussian sum” by

$$\mathfrak{s}(\varepsilon \varphi) = \sum_{\sigma} \varepsilon \varphi(\sigma^{-1}) \sigma(\zeta_{fp^{n+1}})$$

where σ varies over $\text{Gal}(\mathbf{Q}_p(\mu_{fp^{n+1}})/\mathbf{Q}_p)$. The nonvanishing of $\mathfrak{s}(\varepsilon \varphi)$ can be proved using our choice of ζ_f . We cannot say much else about $\mathfrak{s}(\varepsilon \varphi)$. One can extend ε (noncanonically!) to a global character ε' . Then $\varepsilon' \varphi$ can be regarded as a character of $\text{Gal}(\mathbf{Q}(\mu_{fp^{n+1}})/\mathbf{Q})$. The sum $\sum_{\sigma} \varepsilon' \varphi(\sigma^{-1}) \sigma(\zeta_{fp^{n+1}})$, where σ now varies over that global Galois group, would give a standard Gaussian sum (namely, for the inverse of $\varepsilon' \varphi$) and $\mathfrak{s}(\varepsilon \varphi)$ would just be part of that sum.

The next theorem will not be proven here, but will be the main result in the sequel. It gives an interpolation property for a generator of the characteristic ideal. As we will discuss in the sequel, we can also give an interpolation property for $\varphi_0(\theta_{\chi, \varepsilon})$ in most cases, where φ_0 is the trivial character of Γ .

Theorem 3. *Suppose that $d = 2$ and that assumption A is satisfied. Then the characteristic ideal of $X_{\chi, \varepsilon}(\mathbf{Q}_{\infty})$ has a generator $\theta_{\chi, \varepsilon}$ with the following property. There exists a norm-compatible sequence $\{\eta_n\}$, where*

$\eta_n \in U_n$ for all $n \geq 0$, such that

$$\varphi(\theta_{\chi,\varepsilon}) = \frac{1}{\mathfrak{s}(\varepsilon\varphi)} \cdot \sum_{\delta \in \Delta_n} \chi\varphi(\delta^{-1}) \log_p(|\delta(\eta_n)|_\varepsilon)$$

for all $n \geq 1$ and all $\varphi \in \text{Hom}_{\text{cont}}(\Gamma, \overline{\mathbf{Q}_p}^\times)$ of order p^n .

The proof of this theorem involves extending a classical result of Iwasawa to this setting. Indeed, we prove a certain four-term exact sequence involving the module $X_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ and certain inverse limits of local unit groups and global unit groups. The sequence $\{\eta_n\}$ defines an element η_∞ in the Λ -module $\mathcal{U}_\infty = \varprojlim \mathcal{U}_n$. The ε -component $\mathcal{U}_\infty^{(\varepsilon)}$ has Λ -rank 1, and is usually just isomorphic to Λ . The projection of η_∞ to $\mathcal{U}_\infty^{(\varepsilon)}$ generates a Λ -submodule of $\mathcal{U}_\infty^{(\varepsilon)}$ of rank 1. The corresponding quotient module has the same characteristic ideal as $X_{\chi,\varepsilon}(\mathbf{Q}_\infty)$. All of this will be discussed in the sequel. The arguments work for arbitrary d if we make some additional assumptions concerning ρ .

We are not entirely satisfied with the interpolation property in Theorem 3. For one thing, it depends on the choice of ζ_f . However, since we are free to multiply $\theta_{\chi,\varepsilon}$ by an arbitrary unit in $\Lambda_{\mathcal{O}}$, we are hoping to find more natural interpolation factors, perhaps depending on χ and ε in some canonical way. One simple (and canonical) choice would be $\tau(\omega^a\varphi)/p^{n+1}$ in place of $1/\mathfrak{s}(\varepsilon\varphi)$, where $\tau(\omega^a\varphi)$ is the standard Gaussian sum. However, that choice does not involve the unramified character β and corresponds to multiplying by a unit in a possibly larger ring $\Lambda_{\mathcal{O}'}$, instead of in $\Lambda_{\mathcal{O}}$, where \mathcal{O}' is the ring of integers in a certain unramified extension \mathcal{F}' of \mathcal{F} .

Now we will discuss the analytic side of our theory. Recall that d^+ denotes the multiplicity of the trivial character in $\rho|_{\Delta_v}$, where v is any archimedean prime of K and Δ_v is the corresponding decomposition subgroup of Δ . The multiplicity of the nontrivial character of Δ_v (if K is not totally real) is denoted by d^- . Of course, $d^- = d - d^+$. Our most general results concern the case where $d = 2$. Since we assume that $d^+ = 1$, we have $d^- = 1$ and ρ has odd determinant. The action of $\Delta_{\mathfrak{p}}$ on $V/V^{(\varepsilon_{\mathfrak{p}})}$ is given by a character $\varepsilon'_{\mathfrak{p}} : \Delta_{\mathfrak{p}} \rightarrow \mathcal{F}^\times$. As before, we can define a character $\varepsilon' : \text{Gal}(\mathcal{K}/\mathbf{Q}_p) \rightarrow \mathcal{F}^\times$. We assume the Artin conjecture for ρ and all of its twists by one-dimensional characters so that we can associate to ρ a certain modular form f_ρ of weight 1. We will define a p -adic L -function $L_p(s, \chi, \varepsilon)$ as the restriction of a 2-variable p -adic L -function to a certain line.

For simplicity, we will assume here that ε' is unramified. In general, we can reduce to that case by twisting by some power of ω . By using the fixed embeddings σ_∞ and σ_p , we can regard the q -expansion of f_ρ as having coefficients in \mathcal{F} . One can then apply a theorem of Wiles to prove the existence of a Hida family of modular forms with the following property (which we describe here under a certain simplifying assumption). The modular forms in the family will be given by q -expansions with coefficients in \mathcal{F} . Roughly speaking, in each weight $k \geq 2$, the corresponding p -adic Galois representation ρ_k has an unramified quotient (when considered as a representation of $G_{\mathbf{Q}_p}$) and the action on that quotient is given by a character ε'_k of $G_{\mathbf{Q}_p}$. As the weight k approaches 1 p -adically, ε'_k approaches ε' . The q -expansion of the corresponding modular forms f_k in the family approach the q -expansion of f_1 (coefficient by coefficient, as elements in the formal power series ring $\mathcal{F}[[q]]$).

The two-variable p -adic L -function for a Hida family was first constructed by Mazur under certain assumptions and then by Kitagawa in [23], and later by a different method in [19]. As above, we describe this function under a simplifying assumption which we mention below. Under some mild restrictions on ρ , the Hida family is uniquely determined by the choice of ε' . However, since ρ is 2-dimensional, the choice of ε determines ε' , and we will use ε in our notation. The corresponding two-variable p -adic L -function that we consider here is defined using the so-called canonical periods, following [7]. One of the variables is the weight variable k . The second variable is the cyclotomic variable s . If we restrict that function to a weight $k \geq 2$, we essentially obtain the p -adic L -function $L_p(s, f_k)$ attached to the modular form f_k (as defined in [26]). To be precise, the restriction is $u_k \cdot L_p(s, f_k)$, where the factor u_k is in \mathcal{O}^\times .

If we specialize to $k = 1$, we obtain a function which we denote by $L_p(s, \chi, \varepsilon)$ for $s \in \mathbf{Z}_p$. It is only well-defined up to a factor in \mathcal{O}^\times . Furthermore, there is an element $\theta_{\chi, \varepsilon}^{an}$ in $\Lambda_{\mathcal{O}}$ such that

$$(1) \quad L_p(s, \chi, \varepsilon) = \kappa^{1-s}(\theta_{\chi, \varepsilon}^{an})$$

for all $s \in \mathbf{Z}_p$. We now denote the element $\theta_{\chi, \varepsilon}$ occurring in Theorem 3 by $\theta_{\chi, \varepsilon}^{al}$. With this notation, we make the following conjecture:

Conjecture *The principal ideals in $\Lambda_{\mathcal{O}}$ generated by $\theta_{\chi, \varepsilon}^{al}$ and $\theta_{\chi, \varepsilon}^{an}$ are equal.*

That is, $\theta_{\chi, \varepsilon}^{an}$ is a generator for the characteristic ideal of $X_{\chi, \varepsilon}(\mathbf{Q}_\infty)$. This conjecture is partly inspired by the fact that, when $d = 1$, one of the equivalent forms of Iwasawa's classical Main Conjecture can be

formulated in precisely this way. (See the introduction of [10].) In that case, one has $\varepsilon_p = \chi|_{\Delta_p}$ and $L_p(s, \chi, \varepsilon)$ should be taken to be $L_p(s, \chi)$, the Kubota-Leopoldt p -adic L -function for χ . Furthermore, under some restrictive assumptions, the above conjecture for $d = 2$ can be deduced from the usual main conjecture for ordinary modular forms of weight $k \geq 2$, first by deducing the so-called two-variable main conjecture for the Hida family and then using a specialization argument at weight $k = 1$. The work of Ochiai in [28] is highly relevant here. We will discuss this approach fully in the sequel. However, at present, although it is clear that $\theta_{\chi, \varepsilon}^{al}$ is nonzero, we cannot prove in general that $\theta_{\chi, \varepsilon}^{an}$ is nonzero.

We have made one simplifying assumption in the above description. Let N denote the level of the weight 1 modular form f_ρ . Write $N = p^a M$, where $p \nmid M$. Let \mathbf{h}_∞ denote Hida's universal ordinary Hecke algebra of level M . One can view \mathbf{h}_∞ in a natural way as a finite flat algebra over a certain subring Λ' . The subring Λ' is isomorphic to a formal power series ring $\mathbf{Z}_p[[T]]$ in one variable. As we explain in Section 5 (using the theorem of Wiles mentioned previously), the form f_ρ together with the choice of ε singles out a certain domain R which is a quotient of the ring \mathbf{h}_∞ and which is a finite integral extension of Λ' . Our simplifying assumption is that $R = \Lambda'$. However, we do not make this assumption in Section 5.

Acknowledgements: *The first author would like to thank the NSF and the second author would like to thank NSERC for grants over the years supporting this research.*

§2. Galois cohomology for Artin representations

We will be quite general at first. Suppose that K is a finite Galois extension of \mathbf{Q} , that $\Delta = \text{Gal}(K/\mathbf{Q})$, and that $\rho : \Delta \rightarrow GL_d(\mathcal{F})$ is any absolutely irreducible representation over \mathcal{F} . We even allow $[K : \mathbf{Q}]$ to be divisible by p . Let χ denote the character of ρ . We use χ consistently in our notation. We assume that \mathcal{F} is a finite extension of \mathbf{Q}_p . Let \mathcal{O} be the ring of integers in \mathcal{F} . Let $D = V/T$, where V is the underlying \mathcal{F} -vector space for ρ and T is a Δ -invariant \mathcal{O} -lattice in V .

Let Σ be any finite set of primes containing p , ∞ , and the primes which are ramified in K/\mathbf{Q} . Thus, $K \subset \mathbf{Q}_\Sigma$, where \mathbf{Q}_Σ is the maximal extension of \mathbf{Q} unramified outside of Σ . We can regard D as a discrete \mathcal{O} -module with an \mathcal{O} -linear action of $\text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$. We denote d by $d(\chi)$ in this section and write $d(\chi) = d^+(\chi) + d^-(\chi)$, reflecting the action of

the decomposition subgroup Δ_v for an archimedean prime v of K . Note that if one assumes that the residual representation for ρ is irreducible, then D is determined up to isomorphism as a $\text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$ -module by χ .

We first discuss the Galois cohomology groups $H^i(\mathbf{Q}_\Sigma/\mathbf{Q}, D)$ for $i \geq 0$. They are discrete \mathcal{O} -modules. Let $h_i = \text{corank}_{\mathcal{O}}(H^i(\mathbf{Q}_\Sigma/\mathbf{Q}, D))$ for $i \geq 0$. As is well known, if $p \geq 3$, then $H^i(\mathbf{Q}_\Sigma/\mathbf{Q}, D) = 0$ for $i \geq 3$. For $p = 2$, those cohomology groups are finite and have exponent 2. Thus, $h_i = 0$ if $i \geq 3$, and so we just consider $i \in \{0, 1, 2\}$. We will be primarily interested in h_1 . The Poitou-Tate formula for the Euler-Poincaré characteristic implies that

$$h_1 = d^-(\chi) + h_2 + h_0$$

If χ is the trivial character, then one sees easily that $h_0 = h_1 = 1$ and hence $h - 2 = 0$. If we assume that χ is not the trivial character, then $h_0 = 0$. The following result is an immediate consequence of the above remarks.

Proposition 2.1. *Assume that χ is nontrivial. Then*

$$\text{corank}_{\mathcal{O}}(H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, D)) \geq d^-(\chi)$$

and equality holds if and only if $\text{corank}_{\mathcal{O}}(H^2(\mathbf{Q}_\Sigma/\mathbf{Q}, D)) = 0$.

It is reasonable to conjecture that $\text{corank}_{\mathcal{O}}(H^2(\mathbf{Q}_\Sigma/\mathbf{Q}, D)) = 0$ for any nontrivial, irreducible Artin representation. This is closely related to Leopoldt's conjecture, as we now explain. Let

$$\mathcal{U}_p = \prod_{\mathfrak{p}|p} \mathcal{U}_{\mathfrak{p}} ,$$

where $\mathcal{U}_{\mathfrak{p}}$ is the group of principal units in the completion $K_{\mathfrak{p}}$. Let U_K consists of the units of K which are principal units at all $\mathfrak{p}|p$. Thus, U_K is a subgroup of the unit group of K of finite index. Consider the natural diagonal map

$$(2) \quad U_K \rightarrow \mathcal{U}_p .$$

The $\mathcal{U}_{\mathfrak{p}}$'s, and hence \mathcal{U}_p , are \mathbf{Z}_p -modules. The map (2) is obviously injective and extends to a map

$$(3) \quad \lambda_p : U_K \otimes_{\mathbf{Z}} \mathbf{Z}_p \rightarrow \mathcal{U}_p .$$

which is \mathbf{Z}_p -linear. The image of λ_p is the closure of the image of the map (2) and will sometimes be denoted simply by \overline{U}_K . Furthermore, Δ acts naturally on both $U_K \otimes_{\mathbf{Z}} \mathbf{Z}_p$ and \mathcal{U}_p . The map λ_p is Δ -equivariant. Tensoring the above objects with \mathcal{F} , we again get a Δ -equivariant map which we denote by $\lambda_{p,\mathcal{F}}$. Then, restricting to the χ -component, we obtain a Δ -equivariant map

$$(4) \quad \lambda_{p,\mathcal{F}}^{(\chi)} : (U_K \otimes_{\mathbf{Z}} \mathcal{F})^{(\chi)} \rightarrow (\mathcal{U}_p \otimes_{\mathbf{Z}_p} \mathcal{F})^{(\chi)} .$$

Leopoldt's conjecture (for K and p) asserts that λ_p is injective. It is equivalent to saying that $\lambda_{p,\mathcal{F}}$ is injective. For each χ , one can conjecture that $\lambda_{p,\mathcal{F}}^{(\chi)}$ is injective. This obviously follows from Leopoldt's conjecture. Conversely, if $\lambda_{p,\mathcal{F}}^{(\chi)}$ is injective for all the absolutely irreducible character χ of Δ , then Leopoldt's conjecture holds for K and p . Here one should choose \mathcal{F} so that all of the absolutely irreducible representations of $\Delta = \text{Gal}(K/\mathbf{Q})$ are realizable over \mathcal{F} .

Let $r(\chi)$ denote the multiplicity of χ in the Δ -representation space $U_K \otimes_{\mathbf{Z}} \mathcal{F}$. The proof of Dirichlet's Unit Theorem allows one to determine $r(\chi)$. In fact, if χ is nontrivial, then $r(\chi) = d^+(\chi)$. If χ is trivial, then $r(\chi) = 0$. On the other hand, the Δ -representation space $\mathcal{U}_p \otimes_{\mathbf{Z}_p} \mathcal{F}$ is isomorphic to the regular representation of Δ over \mathcal{F} and hence χ has multiplicity $d(\chi)$. The image of $\lambda_{p,\mathcal{F}}^{(\chi)}$ is the χ -component of $\text{Im}(\lambda_{p,\mathcal{F}})$. Let $r_p(\chi)$ denote the multiplicity of χ in $\text{Im}(\lambda_{p,\mathcal{F}})$. The conjecture that $\lambda_{p,\mathcal{F}}^{(\chi)}$ is injective can be equivalently stated as follows.

LC(χ, p): *We have $r_p(\chi) = r(\chi)$.*

It is obvious that $r_p(\chi) \leq r(\chi)$. One can reformulate **LC**(χ, p) in terms of Galois cohomology.

Proposition 2.2. *The assertion **LC**(χ, p) holds if and only if the equality $\text{corank}_{\mathcal{O}}(H^2(\mathbf{Q}_{\Sigma}/\mathbf{Q}, D)) = 0$ holds.*

Proof. Consider the restriction map

$$\alpha : H^1(\mathbf{Q}_{\Sigma}/\mathbf{Q}, D) \rightarrow H^1(\mathbf{Q}_{\Sigma}/K, D)^{\Delta} .$$

Since Δ is finite, it follows that the kernel and cokernel of α are both finite. Hence $H^1(\mathbf{Q}_{\Sigma}/\mathbf{Q}, D)$ and $H^1(\mathbf{Q}_{\Sigma}/K, D)^{\Delta}$ have the same \mathcal{O} -corank. Furthermore,

$$H^1(\mathbf{Q}_{\Sigma}/K, D)^{\Delta} = \text{Hom}_{\Delta}(\text{Gal}(M/K), D) ,$$

where M is the maximal abelian pro- p extension of K contained in \mathbf{Q}_Σ . Clearly, M is Galois over \mathbf{Q} and so Δ acts naturally on $\text{Gal}(M/K)$. Also, $\text{Gal}(M/K)$ can be considered as a \mathbf{Z}_p -module and is finitely generated.

Let \tilde{K} denote the compositum of all \mathbf{Z}_p -extensions of K . Then $K \subset \tilde{K} \subseteq M$ and M/\tilde{K} is a finite extension. Furthermore, \tilde{K} is Galois over \mathbf{Q} and hence Δ also acts naturally on $\text{Gal}(\tilde{K}/K)$. These remarks imply the first of the following equalities. The second equality is easily verified.

$$\begin{aligned} \text{corank}_{\mathcal{O}}(H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, D)) &= \text{corank}_{\mathcal{O}}(\text{Hom}_{\Delta}(\text{Gal}(\tilde{K}/K) \otimes_{\mathbf{Z}_p} \mathcal{O}, D)) \\ &= \dim_{\mathcal{F}}(\text{Hom}_{\Delta}(\text{Gal}(\tilde{K}/K) \otimes_{\mathbf{Z}_p} \mathcal{F}, V)). \end{aligned}$$

Hence, the \mathcal{O} -corank of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, D)$ is just the multiplicity of χ in the Δ -representation space $\text{Gal}(\tilde{K}/K) \otimes_{\mathbf{Z}_p} \mathcal{F}$. We next describe that multiplicity in terms of $r_p(\chi)$.

By class field theory, there is a homomorphism

$$\mathcal{U}_p/\text{Im}(\lambda_p) \rightarrow \text{Gal}(\tilde{K}/K)$$

whose kernel and cokernel are finite. The multiplicity of χ in $\mathcal{U}_p \otimes_{\mathbf{Z}_p} \mathcal{F}$ is $d(\chi)$. The multiplicity of χ in $\text{Im}(\lambda_p) \otimes_{\mathbf{Z}_p} \mathcal{F}$ is $r_p(\chi)$. Hence the multiplicity of χ in $\text{Gal}(\tilde{K}/K) \otimes_{\mathbf{Z}_p} \mathcal{F}$ is $d(\chi) - r_p(\chi)$.

On the other hand, as mentioned above, if $\chi \neq \chi_0$, then one has $r(\chi) = d^+(\chi)$. This follows from the well-known fact that

$$U_K \otimes_{\mathbf{Z}} \mathbf{Q} \cong \text{Ind}_{\Delta_v}^{\Delta}(\theta_0)/V_0$$

where v is an archimedean prime of K , θ_0 is the trivial character of the decomposition subgroup Δ_v , and V_0 is the underlying \mathbf{Q} -representation space for the trivial character χ_0 of Δ over \mathbf{Q} . If $\chi \neq \chi_0$, then Frobenius Reciprocity implies that $r(\chi)$ is equal to the multiplicity of θ_0 in $\rho|_{\Delta_v}$, which is indeed equal to $d^+(\chi)$. Of course, we have $r(\chi_0) = 0$.

Using the formula for the Euler-Poincaré characteristic, we find that

$$h_2 = (d(\chi) - r_p(\chi)) - d^-(\chi) - h_0 = d^+(\chi) - r_p(\chi) - h_0$$

where, as previously, h_i denotes the \mathcal{O} -corank of $H^i(\mathbf{Q}_\Sigma/\mathbf{Q}, D)$. It follows that if $\chi \neq \chi_0$, then $r(\chi) - r_p(\chi) = h_2$. This gives the stated equivalence in Proposition 2.1. For $\chi = \chi_0$, the result is obvious. ■

Remark 2.3. If p is an odd prime, then the p -cohomological dimension of $\text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$ is 2. Hence $H^3(\mathbf{Q}_\Sigma/\mathbf{Q}, D[p]) = 0$. Since D is divisible by p , it follows that $H^2(\mathbf{Q}_\Sigma/\mathbf{Q}, D)$ is also divisible by p . Thus,

its Pontryagin dual is a finitely-generated, torsion-free \mathbf{Z}_p -module, and hence is free. It is free as an \mathcal{O} -module. In particular, if the \mathcal{O} -corank of $H^2(\mathbf{Q}_\Sigma/\mathbf{Q}, D)$ is 0, then we actually have $H^2(\mathbf{Q}_\Sigma/\mathbf{Q}, D) = 0$. If $p = 2$, one can just show that $pH^2(\mathbf{Q}_\Sigma/\mathbf{Q}, D)$ is 0.

One useful positive result concerning $\mathbf{LC}(\chi, p)$ is the following.

Proposition 2.4. *If $r(\chi) \geq 1$, then $r_p(\chi) \geq 1$.*

The proof makes use of the following lemma. We regard $U_K \otimes_{\mathbf{Z}} \overline{\mathbf{Q}}$ as a module over the group ring $\overline{\mathbf{Q}}[\Delta]$, where $\overline{\mathbf{Q}}$ is the algebraic closure of \mathbf{Q} in $\overline{\mathbf{Q}_p}$. We regard $\text{Im}(\lambda_p) \otimes_{\mathbf{Z}_p} \overline{\mathbf{Q}_p}$ as a module over $\overline{\mathbf{Q}_p}[\Delta]$ and hence over the subring $\overline{\mathbf{Q}}[\Delta]$.

Lemma 2.5. *Suppose that $\theta \in \overline{\mathbf{Q}}[\Delta]$ and that θ does not annihilate $U_K \otimes_{\mathbf{Z}} \overline{\mathbf{Q}}$. Then θ does not annihilate $\text{Im}(\lambda_p) \otimes_{\mathbf{Z}_p} \overline{\mathbf{Q}_p}$.*

Proof. Let \mathcal{F} be a finite extension of \mathbf{Q}_p containing the coefficients of θ . As before, we let \mathcal{K} be the completion of K at one of the primes of K lying above p . Let \mathcal{U} be the group of principal units in \mathcal{K} . We consider the corresponding projection map

$$pr : \mathcal{U}_p \otimes_{\mathbf{Z}_p} \mathcal{F} = \prod_{\mathfrak{p}|p} (\mathcal{U}_{\mathfrak{p}} \otimes_{\mathbf{Z}_p} \mathcal{F}) \rightarrow \mathcal{U} \otimes_{\mathbf{Z}_p} \mathcal{F}$$

It suffices to show that $\text{Im}(\lambda_{p,\mathcal{F}})^\theta$ has a nontrivial image under the map pr .

By assumption, there exists an $\eta \in U_K$ such that η^θ is nontrivial (as an element in $U_K \otimes_{\mathbf{Z}} \mathcal{F}$). Now $\theta = \sum_{\delta \in \Delta} \alpha_\delta \delta$, where the α_δ 's are in $\mathcal{F} \cap \overline{\mathbf{Q}}$. The set of conjugates $\{\delta(\eta)\}_{\delta \in \Delta}$ generates a subgroup Υ of U_K . Let t denote its rank. One can choose a subset Δ' of Δ of cardinality t so that $\{\delta'(\eta)\}_{\delta' \in \Delta'}$ generates a subgroup of finite index in Υ . Viewing Υ as a \mathbf{Z} -module, the set $\{\delta'(\eta)\}_{\delta' \in \Delta'}$ is a \mathbf{Z} -basis for that subgroup. Thus, replacing θ by an integral multiple if necessary, we have

$$\eta^\theta = \eta^\psi$$

where $\psi = \sum_{\delta' \in \Delta'} \beta_{\delta'} \delta'$ and the $\beta_{\delta'}$'s are in $\mathcal{F} \cap \overline{\mathbf{Q}}$. Since η^ψ is nontrivial, the $\beta_{\delta'}$'s are not all zero.

The map $\log_p : \mathcal{U} \rightarrow \mathcal{K}$ extends to an \mathcal{F} -linear map from $\mathcal{U} \otimes_{\mathbf{Z}_p} \mathcal{F}$ to $\mathcal{K} \otimes_{\mathbf{Z}_p} \mathcal{F}$. The image of η^ψ under the map

$$\log_p \circ pr \circ \lambda_{p,\mathcal{F}} : U_K \otimes_{\mathbf{Z}} \mathcal{F} \rightarrow \mathcal{K} \otimes_{\mathbf{Z}_p} \mathcal{F} .$$

is $\gamma = \sum_{\delta' \in \Delta'} \log_p(\delta'(\eta)) \otimes \beta_{\delta'}$. The inclusions $\mathcal{K}, \mathcal{F} \subset \overline{\mathbf{Q}}_p$ induce a map from $\mathcal{K} \otimes_{\mathbf{Z}_p} \mathcal{F}$ to $\overline{\mathbf{Q}}_p$. The image of γ is $\sum_{\delta' \in \Delta'} \beta_{\delta'} \log_p(\delta'(\eta))$. The set $\{\log_p(\delta'(\eta))\}_{\delta' \in \Delta'}$ is linearly independent over \mathbf{Q} . The Baker-Brumer Theorem [2] implies that $\{\log_p(\delta'(\eta))\}_{\delta' \in \Delta'}$ is linearly independent over $\overline{\mathbf{Q}}$. It follows that $\gamma \neq 0$. This implies that the image of η^θ under $\lambda_{p, \mathcal{F}}$ is nontrivial. Since $\lambda_{p, \mathcal{F}}$ is Δ -equivariant, it indeed follows that θ is not an annihilator of $\text{Im}(\lambda_p) \otimes_{\mathbf{Z}_p} \overline{\mathbf{Q}}_p$. ■

Remark 2.6. The proof of the above lemma shows that if $\theta \in \mathcal{F}[\Delta]$ has coefficients in $\mathcal{F} \cap \overline{\mathbf{Q}}$ and doesn't annihilate $U_K \otimes_{\mathbf{Z}} \mathcal{F}$, then $pr(\text{Im}(\lambda_{p, \mathcal{F}})^\theta)$ is a nontrivial \mathcal{F} -subspace of $\mathcal{U} \otimes_{\mathbf{Z}_p} \mathcal{F}$. This will be important later.

Proposition 2.4 follows from Lemma 2.5. One takes θ to be the idempotent for χ in $\mathcal{F}[\Delta]$ which does have coefficients in $\mathcal{F} \cap \overline{\mathbf{Q}}$. The following corollary follows immediately since $r(\chi) = d^+(\chi)$ when $\chi \neq \chi_0$.

Corollary 2.7. *If $d^+(\chi) = 1$, then $\text{LC}(\chi, p)$ is true.*

A very closely related result is proved in [8]. The above corollary is implicit in that work. In addition, those authors give an interesting class of examples where Leopoldt's conjecture can be proved. In fact, their examples are an illustration of the above corollary. Suppose that $\Delta \cong A_4$, the alternating group of order 12. Suppose also that K is not totally real. Then U_K has rank 5. There are four absolutely irreducible representations of Δ , up to isomorphism, three of dimension 1 and one of dimension 3. One finds easily that the three nontrivial representations are the constituents in $U_K \otimes_{\mathbf{Z}} \mathcal{F}$ as a representation space for Δ , two of dimension 1 and the one of dimension 3, and that they all satisfy $d^+ = 1$. And so Leopoldt's conjecture for K and p follows from the above corollary.

The rest of this section concerns the Galois cohomology groups $H^i(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)$. They are discrete \mathcal{O} -modules with a continuous action of $\Gamma = \text{Gal}(\mathbf{Q}_\infty/\mathbf{Q})$ and so can be regarded as discrete $\Lambda_{\mathcal{O}}$ -modules, where $\Lambda_{\mathcal{O}} = \mathcal{O}[[\Gamma]]$. They are cofinitely generated as $\Lambda_{\mathcal{O}}$ -modules. Their $\Lambda_{\mathcal{O}}$ -coranks can be determined without assuming any conjectures. The results below are essentially consequences of theorems of Iwasawa, notably Theorems 17 and 18 in [22]. We also refer to [12] and [16].

Proposition 2.8. *The $\Lambda_{\mathcal{O}}$ -corank of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)$ is equal to $d^-(\chi)$. If p is odd, then $H^2(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D) = 0$. If $p = 2$, then $pH^2(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D) = 0$.*

Proof. The theorems of Poitou and Tate yield a formula for the Euler-Poincaré characteristic in terms of the $\Lambda_{\mathcal{O}}$ -coranks of the cohomology groups $H^i(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$. The alternating sum of those $\Lambda_{\mathcal{O}}$ -coranks for $0 \leq i \leq 2$ is $-d^-(\chi)$. (See Proposition 3 in [12].) Since the $\Lambda_{\mathcal{O}}$ -corank of $H^0(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$ is clearly 0, one obtains the formula

$$\text{corank}_{\Lambda_{\mathcal{O}}}(H^1(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)) = d^-(\chi) + \text{corank}_{\Lambda_{\mathcal{O}}}(H^2(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)) .$$

The proposition amounts to the assertion that $H^2(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$ has $\Lambda_{\mathcal{O}}$ -corank 0, the so-called *weak Leopoldt conjecture* for the Galois module D .

Let $K_{\infty} = K\mathbf{Q}_{\infty}$, the cyclotomic \mathbf{Z}_p -extension of K . Consider the restriction map

$$H^2(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D) \rightarrow H^2(\mathbf{Q}_{\Sigma}/K_{\infty}, D) .$$

Note that $\text{Gal}(\mathbf{Q}_{\Sigma}/K_{\infty})$ acts trivially on D . Thus, as a $\text{Gal}(\mathbf{Q}_{\Sigma}/K_{\infty})$ -module, D is isomorphic to $(\mathbf{Q}_p/\mathbf{Z}_p)^{d(\chi)}$ where the Galois group acts trivially. Furthermore, the kernel of the above map has exponent dividing the degree $[K_{\infty} : \mathbf{Q}_{\infty}]$ and hence has Λ -corank 0. Thus, it suffices to show that $H^2(\mathbf{Q}_{\Sigma}/K_{\infty}, \mathbf{Q}_p/\mathbf{Z}_p)$ has Λ -corank 0.

Now $H^1(\mathbf{Q}_{\Sigma}/K_{\infty}, \mathbf{Q}_p/\mathbf{Z}_p) = \text{Hom}(\text{Gal}(M_{\infty}^{\Sigma}/K_{\infty}), \mathbf{Q}_p/\mathbf{Z}_p)$. Here M_{∞}^{Σ} denotes the maximal, abelian pro- p extension of K_{∞} contained in \mathbf{Q}_{Σ} . One can regard $\text{Gal}(M_{\infty}^{\Sigma}/K_{\infty})$ as a Λ -module. According to a well-known theorem of Iwasawa, the Λ -rank of $\text{Gal}(M_{\infty}^{\Sigma}/K_{\infty})$ is equal to $r_2(K)$, the number of complex primes of K . (See Theorem 17 in [22] for the case where Σ is the set of primes Σ_0 lying above p or ∞ . For larger Σ , one can use the fact that there are only finitely many primes of K_{∞} lying above primes in $\Sigma - \Sigma_0$ to show that $\text{Gal}(M_{\infty}^{\Sigma}/M_{\infty}^{\Sigma_0})$ has finite \mathbf{Z}_p -rank and hence is a torsion Λ -module.) Thus, the $\Lambda_{\mathcal{O}}$ -corank of $H^1(\mathbf{Q}_{\Sigma}/K_{\infty}, \mathbf{Q}_p/\mathbf{Z}_p)$ is $r_2(K)$. Again, the Poitou-Tate Duality Theorems imply that the Euler-Poincaré characteristic (which is defined as the alternating sum of the $\Lambda_{\mathcal{O}}$ -coranks of the $H^i(\mathbf{Q}_{\Sigma}/K_{\infty}, \mathbf{Q}_p/\mathbf{Z}_p)$'s for $0 \leq i \leq 2$) is equal to $-r_2(K)$. It follows that $H^2(\mathbf{Q}_{\Sigma}/K_{\infty}, \mathbf{Q}_p/\mathbf{Z}_p)$ indeed has $\Lambda_{\mathcal{O}}$ -corank 0.

If p is odd, then Proposition 4 in [12] asserts that the Pontryagin dual of $H^2(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$ is a free Λ -module. Since its rank is 0, it must vanish. Hence $H^2(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D) = 0$. By considering the restriction map to an imaginary quadratic extension of \mathbf{Q}_{∞} , the argument in [12] shows that $H^2(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$ has exponent 1 or 2 when $p = 2$. ■

One other basic result about the structure of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)$ and a $\Lambda_{\mathcal{O}}$ -module concerns the phenomenon of *purity*, namely that the torsion submodule of the Pontryagin dual of that $\Lambda_{\mathcal{O}}$ -module has support which is purely of codimension 1. This is the content of the following proposition.

Proposition 2.9. *Assume that p is odd. Then the Pontryagin dual of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)$ has no nontrivial, finite $\Lambda_{\mathcal{O}}$ -submodules.*

Proof. This follows immediately from Proposition 5 in [12], the crucial assumption being that $H^2(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)$ vanishes. If the image of ρ has order prime to p , one could alternatively use the fact that the Λ -module $\text{Gal}(M_\infty^\Sigma/K_\infty)$ has no nontrivial, finite Λ -submodules. This is a consequence of Theorem 18 in [22]. See also Theorem 11.3.11, part (i), in [27], where the assertion is formulated equivalently in terms of projective dimension. ■

§3. Selmer groups for Artin representations 1

We will prove Theorem 1 in this section. More generally, we prove the finiteness of the Selmer group over \mathbf{Q}_n for all $n \geq 0$. Suppose that ρ is an irreducible Artin representation of $G_{\mathbf{Q}}$ over \mathcal{F} satisfying Hypothesis A in the introduction. Since $d^+ = 1$, one sees easily that ρ is absolutely irreducible. We assume that ρ factors through $\Delta = \text{Gal}(K/\mathbf{Q})$, where K is a finite Galois extension of \mathbf{Q} of degree prime to p . We consider the Selmer groups $S_{\chi, \varepsilon}(\mathbf{Q}_n)$ for all $n \geq 0$. The definition of the above Selmer groups can be described as follows. As usual, the description of the local condition at a prime ν of a number field of residue field characteristic ℓ always implicitly involves choosing a fixed embedding of \mathbf{Q} into \mathbf{Q}_ℓ which induces the prime ν on the number field. The Selmer group is independent of those choices.

If ℓ is any finite prime and ν is a prime of \mathbf{Q}_n lying over ℓ , we denote the completion of \mathbf{Q}_n at ν by $\mathbf{Q}_{n, \nu}$. If ν is a prime of \mathbf{Q}_∞ lying over ℓ , then we let $\mathbf{Q}_{\infty, \nu} = \bigcup_{n \geq 0} \mathbf{Q}_{n, \nu}$. Note that if $\ell \neq p$, then $\mathbf{Q}_{\infty, \nu} \subset \mathbf{Q}_\ell^{unr}$. For such a prime ℓ , we define

$$\mathcal{H}_\ell(\mathbf{Q}_n, D) = \prod_{\nu|\ell} \text{Im}(H^1(\mathbf{Q}_{n, \nu}, D) \rightarrow H^1(\mathbf{Q}_\ell^{unr}, D))$$

for all $n \geq 0$. There is a unique prime π of \mathbf{Q}_∞ lying over p . Thus, our chosen embedding of \mathbf{Q} into $\overline{\mathbf{Q}}_p$ induces π on \mathbf{Q}_∞ . Let \mathfrak{p} be the prime of K induced by that embedding. We then define

$$(5) \quad \mathcal{H}_p(\mathbf{Q}_n, D) = \text{Im} \left(H^1(\mathbf{Q}_{n,\pi}, D) \rightarrow H^1(\mathbf{Q}_{n,\pi}^{unr}, D/D^{(\varepsilon_{\mathfrak{p}})}) \right).$$

We can then define the Selmer groups to be considered by

$$(6) \quad S_{\chi,\varepsilon}(\mathbf{Q}_n) = \text{Ker} \left(H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_n, D) \rightarrow \prod_{\ell \in \Sigma} \mathcal{H}_\ell(\mathbf{Q}_n, D) \right).$$

for any $n \geq 0$. Here Σ is a finite set of primes of \mathbf{Q} containing p and ∞ chosen so that $K \subset \mathbf{Q}_\Sigma$. The global-to-local map occurring in this definition is defined by the various restriction maps. Since we are taking p to be odd, the 1-cocycle classes are certainly trivial at the archimedean primes of \mathbf{Q}_n . And so the product in (6) is just over the nonarchimedean primes in Σ .

Note that if χ is trivial, then $\varepsilon_{\mathfrak{p}}$ is trivial and $D^{(\varepsilon_{\mathfrak{p}})} = D$. It is then obvious that

$$(7) \quad \text{Hom}(\text{Gal}(\mathbf{Q}_\infty/\mathbf{Q}_n), D) \subseteq S_{\chi,\varepsilon}(\mathbf{Q}_n)$$

and this Selmer group is therefore infinite. Actually, one can easily show that equality holds in (7). However, as we stated in the introduction, we are assuming throughout this paper that χ is nontrivial. We will now prove the following result.

Proposition 3.1. *Assume that Hypothesis A holds. Then $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ is finite for all $n \geq 0$.*

Proof. Since $p \nmid [K : \mathbf{Q}]$, we have $K \cap \mathbf{Q}_n = \mathbf{Q}$. Let $K_n = K\mathbf{Q}_n$, the n -th layer of the cyclotomic \mathbf{Z}_p -extension of K . Then we can identify $\text{Gal}(K_n/\mathbf{Q}_n)$ with Δ and $\text{Gal}(K_n/K)$ with $\text{Gal}(\mathbf{Q}_n/\mathbf{Q})$, which in turn can be identified with Γ/Γ_n . Here $\Gamma_n = \Gamma^{p^n} = \text{Gal}(\mathbf{Q}_\infty/\mathbf{Q}_n)$. Let $\Delta_n = \text{Gal}(K_n/\mathbf{Q})$. We have a canonical isomorphism $\Delta_n \cong \Delta \times \Gamma/\Gamma_n$. Each prime \mathfrak{p} of K lying over p is totally ramified in K_n/K . Let \mathfrak{p}_n be the unique prime of K_n lying over \mathfrak{p} and let $\mathcal{U}_{n,\mathfrak{p}}$ denote the group of principal units in the completion of K_n at \mathfrak{p}_n . The decomposition subgroups of $\text{Gal}(K_n/\mathbf{Q}_n)$ and Δ_n for \mathfrak{p}_n are then identified with $\Delta_{\mathfrak{p}}$ and $\Delta_{\mathfrak{p}} \times \Gamma/\Gamma_n$, respectively. Both of these decomposition groups act on $\mathcal{U}_{n,\mathfrak{p}}$. There is a natural action of Δ_n on

$$\mathcal{U}_{n,p} = \prod_{\mathfrak{p}|p} \mathcal{U}_{n,\mathfrak{p}}.$$

As in Section 2, we have a natural Δ_n -equivariant map

$$\lambda_{n,p} : U_n \otimes_{\mathbf{Z}} \mathbf{Z}_p \rightarrow \mathcal{U}_{n,p} ,$$

where U_n is the group of units of K_n which are principal units at \mathfrak{p}_n for all $\mathfrak{p}|p$. The image of $\lambda_{n,p}$ will be denoted by $\overline{U}_{n,p}$. It is just the closure of the image of U_n under the natural diagonal map from U_n into $\mathcal{U}_{n,p}$.

We consider the Selmer groups $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ for $n \geq 0$. The restriction map

$$H^1(\mathbf{Q}_n, D) \rightarrow H^1(K_n, D)^\Delta = \text{Hom}_\Delta(G_{K_n}^{ab}, D) ,$$

is an isomorphism and the image of $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ under that map is contained in $\text{Hom}_\Delta(\text{Gal}(M_n/K_n), D)$, where M_n is the maximal abelian pro- p extension of K_n unramified outside of the set of primes lying over p . This assertion captures the local conditions defining $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ at all primes $\ell \neq p$. Suppose that $\xi \in \text{Hom}_\Delta(\text{Gal}(M_n/K_n), D)$. Noting that the local degree $[K_{n,\mathfrak{p}_n} : \mathbf{Q}_{n,\pi}]$ is prime to p , one sees that ξ is in the image of $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ if and only if the image of the inertia subgroup $I_{n,\mathfrak{p}}$ of $\text{Gal}(M_n/K_n)$ for \mathfrak{p}_n under ξ is contained in $D^{(\varepsilon_{\mathfrak{p}})}$. Note that if this condition is satisfied for one $\mathfrak{p}|p$, it is satisfied for all those \mathfrak{p} 's. Also, $I_{n,\mathfrak{p}}$ is $\Delta_{\mathfrak{p}}$ -invariant. Now we can extend the map ξ to a Δ -equivariant map

$$\xi_{\mathcal{O}} : \text{Gal}(M_n/K_n)_{\mathcal{O}} \rightarrow D ,$$

where the subscript \mathcal{O} here (and elsewhere) denotes the tensor product over \mathbf{Z}_p with \mathcal{O} , a ring which is finite and flat over \mathbf{Z}_p . The map $\xi_{\mathcal{O}}$ is continuous and \mathcal{O} -linear. Furthermore, with this notation, we have a $\Delta_{\mathfrak{p}}$ -decomposition

$$I_{n,\mathfrak{p},\mathcal{O}} = I_{n,\mathfrak{p},\mathcal{O}}^{(\varepsilon_{\mathfrak{p}})} \times J_{n,\mathfrak{p},\mathcal{O}}$$

where $J_{n,\mathfrak{p},\mathcal{O}}$ is the direct product of all of the $\Delta_{\mathfrak{p}}$ -components of $I_{n,\mathfrak{p},\mathcal{O}}$ apart from the $\varepsilon_{\mathfrak{p}}$ -component. In terms of this decomposition, ξ is in the image of $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ under the restriction map if and only if $\xi_{\mathcal{O}}(J_{n,\mathfrak{p},\mathcal{O}}) = 0$ for one (and hence for all) primes \mathfrak{p} lying over p . Thus, the Selmer condition for ξ is that $\xi_{\mathcal{O}}$ factors through the quotient $\text{Gal}(M_n/K_n)_{\mathcal{O}}/J_{n,\mathfrak{p},\mathcal{O}}$, where $J_{n,\mathfrak{p},\mathcal{O}}$ denotes the \mathcal{O} -submodule of $\text{Gal}(M_n/K_n)_{\mathcal{O}}$ generated by all the $J_{n,\mathfrak{p},\mathcal{O}}$'s for $\mathfrak{p}|p$.

For each $\mathfrak{p}|p$, we have $\mathcal{U}_{n,\mathfrak{p},\mathcal{O}} = \mathcal{U}_{n,\mathfrak{p},\mathcal{O}}^{(\varepsilon_{\mathfrak{p}})} \times \mathcal{V}_{n,\mathfrak{p},\mathcal{O}}$, where the second factor is the direct product of the $\Delta_{\mathfrak{p}}$ -components of $\mathcal{U}_{n,\mathfrak{p},\mathcal{O}}$ apart from the $\varepsilon_{\mathfrak{p}}$ -component. We then have $\mathcal{U}_{n,p,\mathcal{O}} = \mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]} \times \mathcal{V}_{n,p,\mathcal{O}}$, where we

use the following notation:

$$\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]} = \prod_{\mathfrak{p}|p} \mathcal{U}_{n,p,\mathcal{O}}^{(\varepsilon_{\mathfrak{p}})}, \quad \mathcal{V}_{n,p,\mathcal{O}} = \prod_{\mathfrak{p}|p} \mathcal{V}_{n,p,\mathcal{O}}$$

The projection map from $\mathcal{U}_{n,p,\mathcal{O}}$ to $\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]}$ will be denoted by $\pi_{n,\varepsilon}$. We should point out that Δ_n acts naturally on $\mathcal{U}_{n,p,\mathcal{O}}$ and that both $\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]}$ and $\mathcal{V}_{n,p,\mathcal{O}}$ are invariant under the action of Δ_n . The map $\pi_{n,\varepsilon}$ is Δ_n -equivariant and \mathcal{O} -linear.

Let L_n denote the p -Hilbert class field of K_n , the maximal, abelian, unramified p -extension of K_n . We have $K_n \subseteq L_n \subset M_n$. Class field theory gives an exact sequence

$$(8) \quad 1 \rightarrow \overline{U}_{n,p} \rightarrow \mathcal{U}_{n,p} \xrightarrow{\alpha} \text{Gal}(M_n/K_n) \rightarrow \text{Gal}(L_n/K_n) \rightarrow 1.$$

The image of $\mathcal{U}_{n,p}$ (as a factor in $\mathcal{U}_{n,p}$) under the reciprocity map α is $I_{n,p}$. Tensoring the above exact sequence with \mathcal{O} , the \mathcal{O} -submodule $J_{n,p,\mathcal{O}}$ of $\text{Gal}(M_n/K_n)_{\mathcal{O}}$ is the image of $\mathcal{V}_{n,p,\mathcal{O}}$ under the map $\alpha_{\mathcal{O}}$. The above observations then give us the obvious inclusion

$$H_{unr}^1(\mathbf{Q}_n, D) \subseteq S_{\chi,\varepsilon}(\mathbf{Q}_n)$$

as well as the following canonical isomorphism

$$S_{\chi,\varepsilon}(\mathbf{Q}_n)/H_{unr}^1(\mathbf{Q}_n, D) \cong \text{Hom}_{\mathcal{O}[\Delta]}(\mathcal{U}_{n,p,\mathcal{O}}/\mathcal{V}_{n,p,\mathcal{O}}\overline{U}_{n,p,\mathcal{O}}, D).$$

Note that $\text{Gal}(\mathbf{Q}_n/\mathbf{Q}) = \Gamma/\Gamma_n$ acts naturally on $H_{unr}^1(\mathbf{Q}_n, D)$ and $S_{\chi,\varepsilon}(\mathbf{Q}_n)$. Also, since Δ_n is identified with $\Delta \times \Gamma/\Gamma_n$, there is also an action of Γ/Γ_n on the second group. The isomorphism is Γ/Γ_n -equivariant.

The restriction map defines an isomorphism

$$H_{unr}^1(\mathbf{Q}_n, D) \cong \text{Hom}_{\Delta}(\text{Gal}(L_n/K_n), D)$$

and this group is finite. We also have a canonical isomorphism

$$\mathcal{U}_{n,p,\mathcal{O}}/\mathcal{V}_{n,p,\mathcal{O}}\overline{U}_{n,p,\mathcal{O}} \cong \mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]}/\pi_{n,\varepsilon}(\overline{U}_{n,p,\mathcal{O}})$$

as $\mathcal{O}[\Delta_n]$ -modules. For brevity, we will denote $\pi_{n,\varepsilon}(\overline{U}_{n,p,\mathcal{O}})$ by $\overline{U}_{n,p,\mathcal{O}}^{[\varepsilon]}$. Thus,

$$\overline{U}_{n,p,\mathcal{O}}^{[\varepsilon]} = \text{Im}(\pi_{n,\varepsilon} \circ \lambda_{n,p,\mathcal{O}}).$$

It is an $\mathcal{O}[\Delta_n]$ -submodule of $\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]}$.

In summary, $S_{\chi, \varepsilon}(\mathbf{Q}_n)$ contains the finite \mathcal{O} -submodule $H_{unr}^1(\mathbf{Q}_n, D)$ and the corresponding quotient is $\text{Hom}_{\mathcal{O}[\Delta]}(\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]}/\overline{\mathcal{U}}_{n,p,\mathcal{O}}^{[\varepsilon]}, D)$. In particular, $S_{\chi, \varepsilon}(\mathbf{Q}_n)$ is finite if and only if $(\overline{\mathcal{U}}_{n,p,\mathcal{O}}^{[\varepsilon]})^{(\chi)}$ has finite index in $(\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]})^{(\chi)}$. We will verify this by showing that both \mathcal{O} -modules have the same \mathcal{O} -rank. To have a more compact notation, we will denote the χ -components $(\overline{\mathcal{U}}_{n,p,\mathcal{O}}^{[\varepsilon]})^{(\chi)}$ and $(\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon]})^{(\chi)}$ by $\overline{\mathcal{U}}_{n,p,\mathcal{O}}^{[\varepsilon](\chi)}$ and $\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon](\chi)}$, respectively.

Let $\mu_{n,\mathfrak{p}}$ denote the torsion subgroup of $\mathcal{U}_{n,\mathfrak{p}}$. The p -adic log map on $\mathcal{U}_{n,\mathfrak{p}}$ is $\text{Gal}(\mathcal{K}_n/\mathbf{Q}_p)$ -equivariant and its image is open in the additive group of \mathcal{K}_n . Its kernel is $\mu_{n,\mathfrak{p}}$. Thus, $\mathcal{U}_{n,\mathfrak{p}} \otimes_{\mathbf{Z}_p} \mathbf{Q}_p$ is isomorphic to the regular representation of $\text{Gal}(\mathcal{K}_n/\mathbf{Q}_p)$ over \mathbf{Q}_p . Recall that $\text{Gal}(\mathcal{K}_n/\mathbf{Q}_p)$ is identified with $\Delta_{\mathfrak{p}} \times \Gamma/\Gamma_n$. Since $\varepsilon_{\mathfrak{p}}$ is 1-dimensional, it follows that $\mathcal{U}_{n,\mathfrak{p},\mathcal{O}}^{(\varepsilon_{\mathfrak{p}})} \otimes_{\mathcal{O}} \mathcal{F}$ is isomorphic to the regular representation of Γ/Γ_n over \mathcal{F} . In particular, one sees that the rank $\mathcal{O}(\mathcal{U}_{n,\mathfrak{p},\mathcal{O}}^{(\varepsilon_{\mathfrak{p}})}) = p^n$.

A subscript \mathcal{F} will indicate tensoring over \mathcal{O} with \mathcal{F} . For example, we denote $\mathcal{U}_{n,\mathfrak{p},\mathcal{O}} \otimes_{\mathcal{O}} \mathcal{F}$ by $\mathcal{U}_{n,\mathfrak{p},\mathcal{F}}$. With this notation, we have

$$\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]} = \prod_{\mathfrak{p}|p} \mathcal{U}_{n,\mathfrak{p},\mathcal{F}}^{(\varepsilon_{\mathfrak{p}})}.$$

As a representation space for $\Delta_n = \Delta \times \Gamma/\Gamma_n$ over \mathcal{F} , one can view $\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]}$ as an induced representation. For this purpose, we single out one choice of $\mathfrak{p}|p$ as in the introduction. The decomposition subgroup of Δ_n for \mathfrak{p} is $\Delta_{\mathfrak{p}} \times \Gamma/\Gamma_n$, which we denote by $\Delta_{n,\mathfrak{p}}$. Furthermore, $\mathcal{U}_{n,\mathfrak{p},\mathcal{F}}^{(\varepsilon_{\mathfrak{p}})}$ is a representation space for $\Delta_{n,\mathfrak{p}}$ of dimension p^n and is isomorphic to $\varepsilon_{\mathfrak{p}} \otimes_{\mathcal{F}} \sigma_n$, where σ_n denotes the regular representation of Γ/Γ_n over \mathcal{F} . We denote this representation of $\Delta_{n,\mathfrak{p}}$ more briefly by $\varepsilon_{\mathfrak{p}}\sigma_n$. We can also regard σ_n as a representation of Δ_n over \mathcal{F} factoring through Δ_n/Δ . Then we have

$$\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]} \cong \text{Ind}_{\Delta_{n,\mathfrak{p}}}^{\Delta_n}(\varepsilon_{\mathfrak{p}}\sigma_n) \cong \text{Ind}_{\Delta_{n,\mathfrak{p}}}^{\Delta_n}(\varepsilon_{\mathfrak{p}}) \otimes_{\mathcal{F}} \sigma_n.$$

One can regard $\text{Ind}_{\Delta_{n,\mathfrak{p}}}^{\Delta_n}(\varepsilon_{\mathfrak{p}})$ as the representation $\text{Ind}_{\Delta_{\mathfrak{p}}}^{\Delta}(\varepsilon_{\mathfrak{p}})$ composed with the restriction map $\Delta_n \rightarrow \Delta$.

Since $\varepsilon_{\mathfrak{p}}$ occurs with multiplicity 1 in $\chi|_{\Delta_{\mathfrak{p}}}$, Frobenius Reciprocity implies that ρ occurs with multiplicity 1 in $\text{Ind}_{\Delta_{\mathfrak{p}}}^{\Delta}(\varepsilon_{\mathfrak{p}})$. Now \mathcal{F} is generated over \mathbf{Q}_p by roots of unity of order prime to p and hence is unramified. The irreducible representations of Γ/Γ_n over \mathcal{F} are τ_j for $0 \leq j \leq n$, where τ_j has kernel Γ_j/Γ_n . Then τ_j has degree $d_j = p^j - p^{j-1}$

for $1 \leq j \leq n$ and degree $d_0 = 1$ for $j = 0$. We have

$$\sigma_n \cong \bigoplus_{j=0}^n \tau_j$$

which has degree p^n . These remarks imply that each of the irreducible representations $\rho \otimes \tau_j$ occurs in $\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]}$ with multiplicity 1. These are precisely the irreducible constituents in $\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon](\chi)}$ which therefore has \mathcal{F} -dimension dp^n . In contrast, recall that $\mathcal{U}_{n,p,\mathcal{F}}$ is isomorphic to the regular representation of Δ_n over \mathcal{F} . The multiplicity of each $\rho \otimes \tau_j$ in $\mathcal{U}_{n,p,\mathcal{F}}$ is d and therefore $\mathcal{U}_{n,p,\mathcal{F}}^{(\chi)}$ has \mathcal{F} -dimension d^2p^n .

Now consider the Δ_n -representation space $U_{n,\mathcal{F}} = U_n \otimes_{\mathbf{Z}} \mathcal{F}$. The absolutely irreducible constituents φ of each σ_n are 1-dimensional. Since \mathbf{Q}_n is totally real, each such φ is an even character. Hence $d^+(\chi\varphi) = 1$ and hence $\chi\varphi$ occurs with multiplicity 1 in $U_n \otimes_{\mathbf{Z}} \overline{\mathbf{Q}}_p$. It follows that $\chi\tau_j$ occurs with multiplicity 1 in $U_{n,\mathcal{F}}$ for each j , $0 \leq j \leq n$. Let $e_{\chi\tau_j}$ be the idempotent for $\chi\tau_j$ in $\mathcal{F}[\Delta_n]$ (and actually in the center of $\mathcal{F}[\Delta_n]$). Then $e_{\chi\tau_j}$ does not annihilate $U_{n,\mathcal{F}}$. Furthermore, if one restricts $\chi\tau_j$ to Δ , one obtains a multiple of χ . Suppose that \mathfrak{p} is a prime of K lying above p . It follows that $\varepsilon_{\mathfrak{p}}$ occurs as a constituent in $\chi\tau_j|_{\Delta_{\mathfrak{p}}}$. Let $e_{\varepsilon_{\mathfrak{p}}} \in \mathcal{F}[\Delta_{\mathfrak{p}}]$ be the corresponding idempotent. We can regard $e_{\varepsilon_{\mathfrak{p}}}$ as an element of $\mathcal{F}[\Delta_n]$. For $0 \leq j \leq n$, let $\theta_j = e_{\varepsilon_{\mathfrak{p}}} e_{\chi\tau_j}$. Thus $\theta_j \in \mathcal{F}[\Delta_n]$. Also, the coefficients of θ_j are in $\overline{\mathbf{Q}}$. Furthermore, the above remarks show that θ_j does not annihilate $U_{n,\mathcal{F}}$. According to remark 2.6, it follows that the projection of $\text{Im}(\lambda_{n,p,\mathcal{F}})^{\theta_j}$ to $\mathcal{U}_{n,p,\mathcal{F}}$ is nontrivial. Note that the image of that projection is actually contained in $\mathcal{U}_{n,p,\mathcal{F}}^{(\varepsilon_{\mathfrak{p}})}$.

The above remarks show that, for $0 \leq j \leq n$, there exists an element $\alpha_{n,j}$ of U_n such that $\beta_{n,j} = \lambda_{n,p}(\alpha_n)^{\theta_j}$ has the following properties. Clearly, $\beta_{j,n}$ is in $\mathcal{U}_{n,p,\mathcal{F}}^{(\chi)}$ for the action of Δ and, more precisely, in the $\chi\tau_j$ component of $\mathcal{U}_{n,p,\mathcal{F}}$ for the action of Δ_n . Also, the image of $\beta_{j,n}$ under the composite map

$$\mathcal{U}_{n,p,\mathcal{F}} \rightarrow \mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]} \rightarrow \mathcal{U}_{n,p,\mathcal{F}}^{(\varepsilon_{\mathfrak{p}})}$$

is nontrivial. Here the first map is $\pi_{n,\varepsilon,\mathcal{F}}$ and the second map is the obvious projection map. It follows that $\pi_{n,\varepsilon,\mathcal{F}}(\beta_{j,n})$ is nontrivial. It must be in the $\chi\tau_j$ -component of $\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]}$. Since $\chi\tau_j$ has multiplicity 1 in $\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]}$, the image of the $\chi\tau_j$ -component of $U_{n,\mathcal{F}}$ under the map $\pi_{n,\varepsilon,\mathcal{F}} \circ \lambda_{n,p,\mathcal{F}}$ must coincide with the $\chi\tau_j$ -component of $\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon]}$. Since this is so for all j , it follows that the image of $U_{n,\mathcal{F}}^{(\chi)}$ under that map is

precisely $\mathcal{U}_{n,p,\mathcal{F}}^{[\varepsilon](\chi)}$. This proves that the image of $U_{n,\mathcal{O}}^{(\chi)}$ under $\pi_{n,\varepsilon} \circ \lambda_{n,p,\mathcal{O}}$ has the same \mathcal{O} -rank as $\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon](\chi)}$ and hence has finite index. That means that $\overline{U}_{n,p,\mathcal{O}}^{[\varepsilon](\chi)}$ indeed has finite index in $\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon](\chi)}$. ■

One of the key observations in the proof of Proposition 3.1 is the following result concerning the structure of $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ as an $\mathcal{O}[\Gamma/\Gamma_n]$ -module.

Proposition 3.2. *We have an exact sequence of finite $\mathcal{O}[\Gamma/\Gamma_n]$ -modules*

$$0 \rightarrow H_{unr}^1(\mathbf{Q}_n, D) \rightarrow S_{\chi,\varepsilon}(\mathbf{Q}_n) \rightarrow \mathrm{Hom}_{\mathcal{O}[\Delta]}(\mathcal{U}_{n,p,\mathcal{O}}^{[\varepsilon](\chi)} / \overline{U}_{n,p,\mathcal{O}}^{[\varepsilon](\chi)}, D) \rightarrow 0 .$$

In the classical case where $d = d^+ = 1$, χ is just an even character of $\mathrm{Gal}(K/\mathbf{Q})$ of order prime to p , where K is a cyclic extension of \mathbf{Q} . One can omit the ε everywhere in this case because χ determines ε . Note that χ does not occur as a constituent in $\mathcal{V}_{n,p,\mathcal{F}}$ when $d = 1$. The exact sequence in Proposition 3.2 follows immediately from (8) by tensoring with \mathcal{O} and taking the χ -component of each term.

§4. Selmer groups for Artin representations 2

We will prove Theorem 2 in this section. Here $\mathbf{Q}_\infty = \bigcup_{n \geq 0} \mathbf{Q}_n$ is the cyclotomic \mathbf{Z}_p -extension of \mathbf{Q} and \mathbf{Q}_n is the subfield of degree p^n over \mathbf{Q} . We can define the Selmer group over \mathbf{Q}_∞ as a direct limit

$$S_{\chi,\varepsilon}(\mathbf{Q}_\infty) = \varinjlim_n S_{\chi,\varepsilon}(\mathbf{Q}_n)$$

where the direct limit is induced by the restriction maps

$$H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_n, D) \rightarrow H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_m, D) .$$

for $m \geq n \geq 0$. Alternatively, we can define this Selmer group as the kernel of a global-to-local map

$$S_{\chi,\varepsilon}(\mathbf{Q}_\infty) = \ker \left(H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D) \rightarrow \prod_{\ell \in \Sigma} \mathcal{H}_\ell(\mathbf{Q}_\infty, D) \right) .$$

Note that, for $\ell \neq p$, $\mathrm{Gal}(\mathbf{Q}_{\infty,\nu}^{unr}/\mathbf{Q}_{\infty,\nu})$ has profinite order prime to p and so an element in $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)$ is locally unramified at $\nu|\ell$ if and

only if it is locally trivial. Thus, we can define

$$\mathcal{H}_\ell(\mathbf{Q}_\infty, D) = \prod_{\nu|\ell} H^1(\mathbf{Q}_{\infty,\nu}, D)$$

for $\ell \neq p$. For $\ell = p$, one defines

$$\begin{aligned} \mathcal{H}_p(\mathbf{Q}_\infty, D) &= \text{Im} \left(H^1(\mathbf{Q}_{\infty,\pi}, D) \rightarrow H^1(\mathbf{Q}_{\infty,\pi}^{unr}, D/D^{(\varepsilon_p)}) \right) \\ &= H^1(\mathbf{Q}_{\infty,\pi}, D)/L(\mathbf{Q}_{\infty,\pi}, D) \end{aligned}$$

where $L(\mathbf{Q}_{\infty,\pi}, D)$ is a $\Lambda_{\mathcal{O}}$ -submodule of $H^1(\mathbf{Q}_{\infty,\pi}, D)$ which sits in the exact sequence

$$0 \rightarrow H^1(\mathbf{Q}_{\infty,\pi}, D^{(\varepsilon_p)}) \rightarrow L(\mathbf{Q}_{\infty,\pi}, D) \rightarrow H_{unr}^1(\mathbf{Q}_{\infty,\pi}, D') \rightarrow 0 .$$

Here we write D' for $D/D^{(\varepsilon_p)}$ for brevity and $H_{unr}^1(\mathbf{Q}_{\infty,\pi}, D')$ denotes the kernel of the restriction map $H^1(\mathbf{Q}_{\infty,\pi}, D') \rightarrow H^1(\mathbf{Q}_{\infty,\pi}^{unr}, D')$. Also, since $p \nmid |\Delta_p|$, we have $D \cong D^{(\varepsilon_p)} \oplus D'$. This implies the injectivity in the above sequence and also implies that the above sequence splits. Thus, $L(\mathbf{Q}_{\infty,\pi}, D)$ can be identified with

$$H^1(\mathbf{Q}_{\infty,\pi}, D^{(\varepsilon_p)}) \oplus H_{unr}^1(\mathbf{Q}_{\infty,\pi}, D') ,$$

considered in the obvious way as a $\Lambda_{\mathcal{O}}$ -submodule of $H^1(\mathbf{Q}_{\infty,\pi}, D)$.

Our study of $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ will be based on the following control theorem.

Proposition 4.1. *Assume that Hypothesis A is satisfied and that χ is not the trivial character. For any $n \geq 0$, we have an exact sequence of $\mathcal{O}[\Gamma/\Gamma_n]$ -modules:*

$$0 \rightarrow S_{\chi,\varepsilon}(\mathbf{Q}_n) \rightarrow S_{\chi,\varepsilon}(\mathbf{Q}_\infty)^{\Gamma_n} \rightarrow H^0(\Delta_p, D/D^{(\varepsilon_p)}) \rightarrow 0 .$$

Note that $H^0(\Delta_p, D/D^{(\varepsilon_p)}) \cong (\mathcal{F}/\mathcal{O})^t$, where t is the multiplicity of the trivial representation of Δ_p in $V/V^{(\varepsilon_p)}$ and the action of Γ/Γ_n on $(\mathcal{F}/\mathcal{O})^t$ is trivial.

Proof. We exclude the case where χ is the trivial character. We apply the snake lemma to the following commutative diagram.

$$\begin{array}{ccccc} S_{\chi,\varepsilon}(\mathbf{Q}_n) & \longrightarrow & H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_n, D) & \xrightarrow{\phi_n} & \prod_{\ell \in \Sigma} \mathcal{H}_\ell(\mathbf{Q}_n, D) \\ \downarrow s_n & & \downarrow h_n & & \downarrow r_n \\ S_{\chi,\varepsilon}(\mathbf{Q}_\infty, D)^{\Gamma_n} & \longrightarrow & H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)^{\Gamma_n} & \longrightarrow & \prod_{\ell \in \Sigma} \mathcal{H}_\ell(\mathbf{Q}_\infty, D)^{\Gamma_n} \end{array}$$

where the vertical maps are the obvious restriction maps and ϕ_n is the global-to-local map defining $S_{\chi, \varepsilon}(\mathbf{Q}_n)$. For our application of the snake lemma, we need the obvious fact that the first maps on each row are injective and also the fact that ϕ_n is surjective, which we prove below.

We first show that the \mathcal{O} -corank of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_n, D)$ is equal to $d^- p^n = (d-1)p^n$. To see this, we have an isomorphism

$$H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_n, D) \cong H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, \text{Ind}_{\mathbf{Q}_n}^{\mathbf{Q}}(D))$$

by Shapiro's Lemma. One can identify $\text{Ind}_{\mathbf{Q}_n}^{\mathbf{Q}}(D)$ with $D \otimes_{\mathcal{O}} \mathcal{O}[\Gamma/\Gamma_n]$ and its \mathcal{O} -corank is dp^n . Let $\mathcal{F}_n = \mathcal{F}(\mu_{p^n})$ and let \mathcal{O}_n denote the ring of integers in that local field. Tensoring with \mathcal{O}_n over \mathcal{O} , it suffices to show that $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, D \otimes_{\mathcal{O}} \mathcal{O}_n[\Gamma/\Gamma_n])$ has \mathcal{O}_n -corank equal to $d^- p^n$. Now $\text{Ind}_{\mathbf{Q}_n}^{\mathbf{Q}}(V) \otimes_{\mathcal{F}} \mathcal{F}_n$ is isomorphic to the direct sum of the irreducible representations $\rho \otimes \varphi$ over \mathcal{F}_n , where φ varies over the characters of Γ/Γ_n , all of which are d -dimensional and have $d^+ = 1$, $d^- = d-1$. For brevity, we let V_φ and T_φ denote the corresponding \mathcal{F}_n -representation spaces and Galois invariant \mathcal{O}_n -lattices. Let $D_\varphi = V_\varphi/T_\varphi$. Propositions 2.1 and 2.2 and corollary 2.7 imply that the \mathcal{O}_n -corank of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, D_\varphi)$ is d^- for all φ . Hence the \mathcal{O}_n -corank of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, \oplus_\varphi D_\varphi)$ is $d^- p^n$. There is a Δ -equivariant surjective homomorphism from $D \otimes \mathcal{O}_n[\Gamma/\Gamma_n]$ to $\oplus_\varphi D_\varphi$ with finite kernel. It follows that the \mathcal{O}_n -corank in question is indeed equal to $d^- p^n$.

We now show that $\mathcal{H}_p(\mathbf{Q}_n, D)$ also has \mathcal{O} -corank equal to $d^- p^n$ and that $\mathcal{H}_\ell(\mathbf{Q}_n, D)$ is finite for all $\ell \in \Sigma$, $\ell \neq p$. We use the Poitou-Tate formulas for local Euler-Poincaré characteristics. Let T^* denote the compact Galois module $\text{Hom}(D, \mu_{p^\infty})$. It is free as an \mathcal{O} -module. For any prime ℓ , let \mathcal{L} be a finite extension of \mathbf{Q}_ℓ . Then Poitou-Tate duality implies that $H^2(\mathcal{L}, D)$ is dual to $H^0(\mathcal{L}, T^*)$. Since the action of $G_{\mathcal{L}}$ on D factors through a finite quotient and the action on T^* factors through an infinite quotient, it follows that $H^0(\mathcal{L}, T^*) = 0$ and hence $H^2(\mathcal{L}, D) = 0$. The only assumption we need is that $G_{\mathcal{L}}$ acts on D through a finite quotient group.

Recall that $\mathcal{H}_p(\mathbf{Q}_n, D)$ is the image of the composite map

$$H^1(\mathbf{Q}_{n, \pi}, D) \longrightarrow H^1(\mathbf{Q}_{n, \pi}, D') \longrightarrow H^1(\mathbf{Q}_{n, \pi}^{unr}, D'),$$

where D' denote $D/D^{(\varepsilon_p)}$ as before. The first map is easily seen to be surjective and so we must just consider the \mathcal{O} -corank of the image of the second map. Now $\text{corank}_{\mathcal{O}}(D') = d-1 = d^-$. As above, we also have $H^2(\mathcal{L}, D') = 0$ for any finite extension \mathcal{L} of \mathbf{Q}_p . The Euler-Poincaré

characteristic formulas for D' imply that

$$\text{corank}_{\mathcal{O}}(H^1(\mathbf{Q}_{n,\pi}, D')) = d^- p^n + \text{corank}_{\mathcal{O}}(H^0(\mathbf{Q}_{n,\pi}, D'))$$

where π is the unique prime above p in \mathbf{Q}_n (or \mathbf{Q}_∞ as before). Here we are using the vanishing of $H^2(\mathbf{Q}_{n,\pi}, D')$. For brevity in the following discussion, if \mathcal{L} is any extension of \mathbf{Q}_p , then we denote $H^0(\mathcal{L}, D')$ simply by $D'(\mathcal{L})$. Since $G_{\mathbf{Q}_p}$ acts on D' through a finite quotient of order prime to p , it follows that $D'(\mathcal{L}) = D'(\mathbf{Q}_p)$ whenever \mathcal{L} is a finite p -extension of \mathbf{Q}_p , or even a pro- p extension of \mathbf{Q}_p . In particular, we have

$$D'(\mathbf{Q}_{n,\pi}) = D'(\mathbf{Q}_p) \cong (\mathcal{F}/\mathcal{O})^t$$

which is \mathcal{O} -divisible and has \mathcal{O} -corank t . Letting $\mathbf{Q}_{n,\pi}^{\text{unr},p}$ denote the unramified \mathbf{Z}_p -extension of $\mathbf{Q}_{n,\pi}$, we also have

$$D'(\mathbf{Q}_{n,\pi}^{\text{unr},p}) = D'(\mathbf{Q}_p) .$$

One sees easily that the kernels of the maps

$$H^1(\mathbf{Q}_{n,\pi}, D') \rightarrow H^1(\mathbf{Q}_{n,\pi}^{\text{unr}}, D'), \quad H^1(\mathbf{Q}_{n,\pi}, D') \rightarrow H^1(\mathbf{Q}_{n,\pi}^{\text{unr},p}, D')$$

coincide. The inflation-restriction sequence implies that this kernel is isomorphic to

$$\begin{aligned} H^1(\mathbf{Q}_{n,\pi}^{\text{unr},p}/\mathbf{Q}_{n,\pi}, D'(\mathbf{Q}_{n,\pi}^{\text{unr},p})) &= H^1(\mathbf{Q}_{n,\pi}^{\text{unr},p}/\mathbf{Q}_{n,\pi}, D'(\mathbf{Q}_p)) \\ &= \text{Hom}(\text{Gal}((\mathbf{Q}_{n,\pi}^{\text{unr},p}/\mathbf{Q}_p), D'(\mathbf{Q}_p)) . \end{aligned}$$

Since $\text{Gal}((\mathbf{Q}_{n,\pi}^{\text{unr},p}/\mathbf{Q}_p)$ has \mathbf{Z}_p -rank 1, it follows that this kernel has \mathcal{O} -corank t and therefore that $\mathcal{H}_p(\mathbf{Q}_n, D)$ indeed has \mathcal{O} -corank equal to $d^- p^n$.

Now suppose $\ell \neq p$ and let ν be a prime of \mathbf{Q}_n above ℓ . In this case, we have the formula

$$\text{corank}_{\mathcal{O}}(H^1(\mathbf{Q}_{n,\nu}, D)) = \text{corank}_{\mathcal{O}}(H^0(\mathbf{Q}_{n,\nu}, D)).$$

This \mathcal{O} -corank could be positive. But, just as above, the inflation-restriction sequence shows that the \mathcal{O} -corank of the kernel of the restriction map $H^1(\mathbf{Q}_{n,\nu}, D) \rightarrow H^1(\mathbf{Q}_{n,\nu}^{\text{unr}}, D)$ is the same as the \mathcal{O} -corank of $H^0(\mathbf{Q}_{n,\nu}, D)$ and hence the image of the restriction map is finite. The finiteness of $\mathcal{H}_\ell(\mathbf{Q}_n, D)$ follows.

Thus, both $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_n, D)$ and $\prod_{\ell \in \Sigma} \mathcal{H}_\ell(\mathbf{Q}_n, D)$ have the same \mathcal{O} -corank. Furthermore, the kernel of the global-to-local map ϕ_n is $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ which is finite by Proposition 3.1. Therefore, the cokernel

of ϕ_n has \mathcal{O} -corank 0 and must be finite. This is one of the hypotheses in Proposition 3.2.1 in [17]. Another hypothesis is called **LEO**(D) there, but that follows immediately from corollary 2.7 and Proposition 2.2. A third hypothesis is that no subquotient of $D[p] \cong T/pT$ is isomorphic to μ_p as a $G_{\mathbf{Q}_n}$ -module. However, this is satisfied because Δ has order prime to p , hence the reduction of ρ modulo p is still irreducible as a representation space over the residue field $\mathcal{O}/(p)$, and has no subquotient which is odd. Consequently, Proposition 3.2.1 in [17] implies that ϕ_n is surjective.

We can now apply the snake lemma to the commutative diagram at the beginning of this proof. The exact sequence in Proposition 4.1 is a statement about $\ker(s_n)$ and $\text{coker}(s_n)$. First of all, note that

$$\ker(h_n) \cong H^1(\Gamma_n, H^0(\mathbf{Q}_\infty, D)) .$$

Since the action of $G_{\mathbf{Q}_n}$ on D factors through a quotient of order prime to p and χ is nontrivial and irreducible, it follows that

$$H^0(\mathbf{Q}_\infty, D) = H^0(\mathbf{Q}, D) = 0 .$$

Therefore, h_n is injective, and hence so is s_n . Also, since $\Gamma_n \cong \mathbf{Z}_p$, it has p -cohomological dimension 1. This implies that $\text{coker}(h_n) = 0$. The snake lemma then implies that $\text{coker}(s_n) \cong \ker(r_n)$. For each $\ell \in \Sigma$, consider the restriction map

$$r_{n,\ell} : \mathcal{H}_\ell(\mathbf{Q}_n, D) \rightarrow \mathcal{H}_\ell(\mathbf{Q}_\infty, D) .$$

Now, if $\ell \neq p$, then ℓ is unramified in $\mathbf{Q}_\infty/\mathbf{Q}$. It follows that, for $\nu|\ell$, $\mathbf{Q}_{n,\nu}^{unr} = \mathbf{Q}_{\infty,\nu}^{unr}$. This implies that $r_{n,\ell}$ is injective.

We now determine $\ker(r_{n,p})$. We use the notation

$$\Gamma_n^{unr} = \text{Gal}(\mathbf{Q}_{n,\pi}^{unr,p}/\mathbf{Q}_{n,\pi}), \quad \Gamma_n^{cyc} = \Gamma_n = \text{Gal}(\mathbf{Q}_{\infty,\pi}/\mathbf{Q}_{n,\pi}) .$$

Just as before, the kernel of the map

$$H^1(\mathbf{Q}_{n,p}, D') \longrightarrow H^1(\mathbf{Q}_{\infty,\pi} \mathbf{Q}_{n,\pi}^{unr}, D')$$

coincides with the kernel of the map

$$H^1(\mathbf{Q}_{n,p}, D') \longrightarrow H^1(\mathbf{Q}_{\infty,\pi} \mathbf{Q}_{n,\pi}^{unr,p}, D') .$$

Furthermore, this kernel is isomorphic to the following \mathcal{O} -modules

$$\begin{aligned} H^1(\mathbf{Q}_{\infty,\pi} \mathbf{Q}_{n,\pi}^{unr,p}/\mathbf{Q}_{n,\pi}, D'(\mathbf{Q}_p)) &\cong \text{Hom}(\Gamma_n^{cyc} \times \Gamma_n^{unr}, D'(\mathbf{Q}_p)) \\ &\cong \text{Hom}(\Gamma_n^{cyc}, D'(\mathbf{Q}_p)) \times \text{Hom}(\Gamma_n^{unr}, D'(\mathbf{Q}_p)) . \end{aligned}$$

The second factor is the kernel of the restriction map

$$H^1(\mathbf{Q}_{n,\pi}, D') \rightarrow H^1(\mathbf{Q}_{n,\pi}^{unr}, D')$$

and therefore $\ker(r_{n,p})$ is isomorphic to the first factor, and hence isomorphic (non-canonically) to $D'(\mathbf{Q}_p) = H^0(\Delta_{\mathfrak{p}}, D')$. It is a cofree \mathcal{O} -module of corank t and Γ/Γ_n acts trivially. We have proved that s_n is indeed injective and has cokernel as stated in the proposition. \blacksquare

As always, we exclude in the above proposition the case where χ is the trivial character. In that case, one easily shows that $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty}) = 0$. This is so even though $S_{\chi,\varepsilon}(\mathbf{Q}_n)$ is infinite for all $n \geq 0$ according to (7). And so, $\ker(s_n)$ is infinite when χ is the trivial character.

The next result is Theorem 2 in the introduction.

Proposition 4.2. *Suppose that Hypothesis A holds. Then $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty})$ is a cofinitely generated, cotorsion $\Lambda_{\mathcal{O}}$ -module.*

Proof. By definition, $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty})$ is a $\Lambda_{\mathcal{O}}$ -submodule of $H^1(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$ and hence its Pontryagin dual $X_{\chi,\varepsilon}$ is a quotient of the Pontryagin dual of $H^1(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$ as a $\Lambda_{\mathcal{O}}$ -module. Proposition 3 in [12] asserts that $H^1(\mathbf{Q}_{\Sigma}/\mathbf{Q}_{\infty}, D)$ is cofinitely generated as a Λ -module, where $\Lambda = \mathbf{Z}_p[[\Gamma]]$. Hence its Pontryagin dual is finitely generated as a Λ -module and hence as a $\Lambda_{\mathcal{O}}$ -module. Therefore, $X_{\chi,\varepsilon}$ is also a finitely generated $\Lambda_{\mathcal{O}}$ -module.

If X is a finitely generated $\Lambda_{\mathcal{O}}$ -module of rank r , then

$$\text{rank}_{\mathcal{O}}(X_{\Gamma_n}) = rp^n + O(1)$$

as $n \rightarrow \infty$. If S is the Pontryagin dual of X , then

$$\text{corank}_{\mathcal{O}}(S^{\Gamma_n}) = \text{rank}_{\mathcal{O}}(X_{\Gamma_n}).$$

Thus, $r = 0$ if and only if $\text{corank}_{\mathcal{O}}(S^{\Gamma_n})$ is bounded. Considering $S = S_{\chi,\varepsilon}(\mathbf{Q}_{\infty})$, Propositions 3.1 and 4.1 imply the boundedness of these \mathcal{O} -coranks. Hence $r = 0$ and that means that $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty})$ is indeed $\Lambda_{\mathcal{O}}$ -cotorsion. \blacksquare

Remark 4.3. This remark concerns so-called *trivial zeros* or *exceptional zeros*. Suppose that the trivial character of $\Delta_{\mathfrak{p}}$ occurs as a constituent in $\chi|_{\Delta_{\mathfrak{p}}}$ with multiplicity t , but that $\varepsilon_{\mathfrak{p}}$ is nontrivial. Proposition 4.1 then implies that $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty})^{\Gamma}$ is infinite and that its \mathcal{O} -corank is equal to t . Thus, $(X_{\chi,\varepsilon})_{\Gamma}$ has \mathcal{O} -rank t . As in the introduction, let $\theta_{\chi,\varepsilon}$

be a generator of the characteristic ideal of the $\Lambda_{\mathcal{O}}$ -module $X_{\chi,\varepsilon}$. If $\varphi : \Gamma \rightarrow \overline{\mathbf{Q}}_p^\times$ is a continuous group homomorphism, then we can extend φ uniquely to a continuous \mathcal{O} -algebra homomorphism $\varphi : \Lambda_{\mathcal{O}} \rightarrow \overline{\mathbf{Q}}_p$. In particular, if φ_0 is the trivial character of Γ , then one obtains a continuous surjective \mathcal{O} -algebra homomorphism $\varphi_0 : \Lambda_{\mathcal{O}} \rightarrow \mathcal{O}$ whose kernel is generated by $\gamma_0 - 1$. Here γ_0 is a fixed topological generator of Γ . The fact that $(X_{\chi,\varepsilon})_{\Gamma}$ has \mathcal{O} -rank t implies that $(\gamma_0 - 1)^t$ divides $\theta_{\chi,\varepsilon}$ in $\Lambda_{\mathcal{O}}$. Thus, $\varphi_0(\theta_{\chi,\varepsilon}) = 0$. One would say that φ_0 is a zero of $\theta_{\chi,\varepsilon}$ of order $\geq t$. It seems reasonable to conjecture that the order of vanishing is exactly t .

As usual in Iwasawa theory, the \mathcal{O} -algebra $\Lambda_{\mathcal{O}}$ is isomorphic to the formal power series ring $\mathcal{O}[[T]]$. One defines such an isomorphism by sending $\gamma_0 - 1$ to T . Thus, $\theta_{\chi,\varepsilon}$ corresponds to a power series $f_{\chi,\varepsilon}(T)$ in $\mathcal{O}[[T]]$. Under the above assumption about $\chi|_{\Delta_{\varepsilon_p}}$, it follows that $f_{\chi,\varepsilon}(T) = T^t g_{\chi,\varepsilon}(T)$, where $g_{\chi,\varepsilon}(T) \in \mathcal{O}[[T]]$. The conjecture is that $g_{\chi,\varepsilon}(0) \neq 0$.

One interesting example is discussed in detail in [13], pages 227-231. The representation ρ in that example is the 2-dimensional irreducible representation of $\Delta = \text{Gal}(K/\mathbf{Q})$, where K is the splitting field over \mathbf{Q} for $x^3 - x + 1$. In fact, $\Delta \cong S_3$ and ρ is realizable over \mathbf{Q} . We can take $\mathcal{F} = \mathbf{Q}_p$ and $\mathcal{O} = \mathbf{Z}_p$. We take $p = 23$. Then $\Delta_{\mathfrak{p}}$ is a subgroup of order 2. In [13], the case where $\varepsilon_{\mathfrak{p}}$ is the nontrivial character of $\Delta_{\mathfrak{p}}$ is discussed. Then $\Delta_{\mathfrak{p}}$ acts trivially on $V/V^{(\varepsilon_{\mathfrak{p}})}$ and $t = 1$. It is shown there that $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty}) \cong \mathbf{Q}_p/\mathbf{Z}_p$. Of course, Γ acts trivially. And so, the characteristic ideal of $X_{\chi,\varepsilon}$ is generated by T . The calculation in [13] is rather subtle and depends on calculating something which one could call the \mathcal{L} -invariant for the representation ρ and $p = 23$. In particular, the \mathcal{L} -invariant turns out to be nonzero. It would be tempting to extend such a calculation to more general ρ 's satisfying Hypothesis A, but we have done nothing in that direction.

Remark 4.4. One can ask whether $H_{unr}^1(\mathbf{Q}_{\infty}, D)^{\Gamma}$ can be infinite. Let t be as in remark 4.3. If $t = 0$, then $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty})^{\Gamma}$ is finite and hence so is $H_{unr}^1(\mathbf{Q}_{\infty}, D)^{\Gamma}$. Suppose now that $t = 1$. Let ε_0 denote the trivial character of $\text{Gal}(\mathcal{K}/\mathbf{Q}_p)$. Hypothesis A is then satisfied and hence $S_{\chi,\varepsilon_0}(\mathbf{Q}_{\infty})$ is $\Lambda_{\mathcal{O}}$ -cotorsion by Proposition 4.2. Actually, Proposition 4.1 implies that $S_{\chi,\varepsilon_0}(\mathbf{Q}_{\infty})^{\Gamma}$ is finite. Since $H_{unr}^1(\mathbf{Q}_{\infty}, D) \subseteq S_{\chi,\varepsilon_0}(\mathbf{Q}_{\infty})$, it follows that $H_{unr}^1(\mathbf{Q}_{\infty}, D)^{\Gamma}$ is finite in that case too. However, if $t \geq 2$, then it turns out that $H_{unr}^1(\mathbf{Q}_{\infty}, D)^{\Gamma}$ is infinite. Its \mathcal{O} -corank will be at least $t - 1$. We will discuss this in the sequel to this paper.

Another basic result concerning $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is the following *purity* result concerning the $\Lambda_{\mathcal{O}}$ -module $X_{\chi,\varepsilon}$ and asserts equivalently that $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is an *almost divisible* $\Lambda_{\mathcal{O}}$ -module in the sense of [16].

Proposition 4.5. *Assume that Hypothesis A is satisfied. Then the Pontryagin dual $X_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ of $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ has no nontrivial, finite $\Lambda_{\mathcal{O}}$ -submodules.*

Proof. This result follows from 4.1.1 in [18]. However, that proposition is formulated in terms of a Galois representation over $\Lambda_{\mathcal{O}}$ instead of over \mathcal{O} , namely the so-called cyclotomic deformation of ρ discussed in Section 3 of [14]. Proposition 3.2 in that paper provides the $\Lambda_{\mathcal{O}}$ -module isomorphism between $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ and the Selmer group over \mathbf{Q} associated to the cyclotomic deformation of ρ . It is not difficult to verify the hypotheses (some of which were already discussed in proving the surjectivity of ϕ_n in the proof of Proposition 4.1.) One hypothesis is that $L(\mathbf{Q}_{\infty,\pi}, D)$ (which was defined above) is almost divisible as a $\Lambda_{\mathcal{O}}$ -module. This just means that $L(\mathbf{Q}_{\infty,\pi}, D)$ has no proper $\Lambda_{\mathcal{O}}$ -submodules of finite index. Now $H_{unr}^1(\mathbf{Q}_{\infty,\pi}, D')$ is isomorphic to $(\mathcal{F}/\mathcal{O})^t$ and is divisible as an \mathcal{O} -module. Furthermore, $H^1(\mathbf{Q}_{\infty,\pi}, D^{(\varepsilon_p)})$ is also divisible as an \mathcal{O} -module because $D^{(\varepsilon_p)}$ is divisible and $G_{\mathbf{Q}_{\infty,\pi}}$ has p -cohomological dimension 1. ■

Remark 4.6. The conclusion in Proposition 4.5 means that $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is an almost divisible $\Lambda_{\mathcal{O}}$ -module. One variant of the Selmer group $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is the so-called *strict* Selmer group $S_{\chi,\varepsilon}^{str}(\mathbf{Q}_\infty)$ where we continue to require elements of $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}_\infty, D)$ to be locally unramified at all $\nu|\ell$ for $\ell \neq p$, but for $\ell = p$, we require those elements to have trivial image in $H^1(\mathbf{Q}_{\infty,\pi}, D')$ (rather than in $H^1(\mathbf{Q}_{\infty,\pi}^{unr}, D')$). However, this just means that we replace

$$L(\mathbf{Q}_{\infty,\pi}, D) = H^1(\mathbf{Q}_{\infty,\pi}, D^{(\varepsilon_p)}) \oplus H_{unr}^1(\mathbf{Q}_{\infty,\pi}, D')$$

by $L^{str}(\mathbf{Q}_{\infty,\pi}, D) = H^1(\mathbf{Q}_{\infty,\pi}, D^{(\varepsilon_p)})$. Since this is an almost divisible $\Lambda_{\mathcal{O}}$ -module, it again follows that it $S_{\chi,\varepsilon}^{str}(\mathbf{Q}_\infty)$ is almost divisible as a $\Lambda_{\mathcal{O}}$ -module. In Section 5, it will be useful to know the same thing for the variant where we replace $L(\mathbf{Q}_{\infty,\pi}, D)$ by

$$L'(\mathbf{Q}_{\infty,\pi}, D) = H^1(\mathbf{Q}_{\infty,\pi}, D^{(\varepsilon_p)})_{\Lambda_{\mathcal{O}}-div} \oplus H_{unr}^1(\mathbf{Q}_{\infty,\pi}, D').$$

Here, for a discrete $\Lambda_{\mathcal{O}}$ -module H , we let $H_{\Lambda_{\mathcal{O}}-div}$ denote the maximal $\Lambda_{\mathcal{O}}$ -divisible submodule of H . As we point out in Section 5, $H^1(\mathbf{Q}_{\infty,\pi}, D^{(\varepsilon_p)})$ fails to be $\Lambda_{\mathcal{O}}$ -divisible just in the case where $\varepsilon = \omega$.

We denote the corresponding Selmer group by $S'_{\chi,\varepsilon}(\mathbf{Q}_\infty)$. Since the $\Lambda_{\mathcal{O}}$ -module $L'(\mathbf{Q}_{\infty,\pi}, D)$ is almost divisible, it will again follow that $S'_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is also an almost divisible $\Lambda_{\mathcal{O}}$ -module.

There is one case where one can prove the analogue of Proposition 4.1 even if $d^+ \neq 1$, namely the case where $d^+ = d - 1$. Then $d^- = 1$. We use the same notation as before, i.e., $\rho, V, T, D, \mathcal{F}, \mathcal{O}$, etc.. Just as in Hypothesis A, we will assume that V has a $\Delta_{\mathfrak{p}}$ -invariant subspace $W_{\mathfrak{p}}$ such that $\dim_{\mathcal{F}}(W) = d^+$ and that the $\Delta_{\mathfrak{p}}$ -representation spaces W and V/W have no irreducible constituents in common. Now $\Delta_{\mathfrak{p}}$ acts on V/W by a character $\varepsilon'_{\mathfrak{p}}$ and the assumption means that $\varepsilon'_{\mathfrak{p}}$ has multiplicity 1 in V . Let $\varepsilon_{\mathfrak{p}} = \chi - \varepsilon'_{\mathfrak{p}}$, which is the character of W as a $\Delta_{\mathfrak{p}}$ -representation space. We will denote W by $V^{(\varepsilon_{\mathfrak{p}})}$ even though $\varepsilon_{\mathfrak{p}}$ is not necessarily an irreducible character of $\Delta_{\mathfrak{p}}$. We use a similar notation for D . Thus, $D \cong D^{(\varepsilon_{\mathfrak{p}})} \oplus D^{(\varepsilon'_{\mathfrak{p}})}$, where the two summands have \mathcal{O} -coranks $d - 1$ and 1, respectively. With this notation, we can define $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ exactly as in the introduction. It turns out that we can use Proposition 4.2 (for a different Artin representation) to prove the following result.

Proposition 4.7. *Keep the above assumptions. Then $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is a cofinitely generated, cotorsion $\Lambda_{\mathcal{O}}$ -module.*

Proof. The fact that the Selmer group is cofinitely generated as a $\Lambda_{\mathcal{O}}$ -module follows from the fact that the same is true for $H^1(\mathbf{Q}_\Sigma/\mathbf{Q}, D)$. We can assume that $\mu_p \subset K$. Let $\mathcal{F}(\omega)$ be the 1-dimensional \mathcal{F} -representation space where Δ acts by character ω . Here ω gives the action of Δ on μ_p and its values are the $(p - 1)$ -st roots of unity (which are in \mathcal{F}). We let $U = \text{Hom}(V, \mathcal{F}(\omega))$ which is the underlying \mathcal{F} -representation space for an Artin representation σ factoring through Δ . Let ψ be the character of σ . Note that $d^+(\psi) = d^-(\chi) = 1$. Furthermore, the orthogonal complement of W is the $\Delta_{\mathfrak{p}}$ -invariant subspace $U^{(\delta_{\mathfrak{p}})}$ of U where $\delta_{\mathfrak{p}} = \omega\varepsilon'_{\mathfrak{p}}^{-1}$. We let δ denote the corresponding character of $\text{Gal}(K/\mathbf{Q}_p)$. Both $\dim_{\mathcal{F}}(U^{(\delta_{\mathfrak{p}})})$ and $d^+(\psi)$ are equal to 1. Furthermore, $\delta_{\mathfrak{p}}$ is not a constituent in $U/U^{(\delta_{\mathfrak{p}})}$. And so we are in the situation considered previously. Proposition 3.4 implies that $S_{\psi,\delta}(\mathbf{Q}_\infty)$ is a cotorsion $\Lambda_{\mathcal{O}}$ -module. We will use arguments in [12] to deduce that $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ is also $\Lambda_{\mathcal{O}}$ -cotorsion. In fact, there is a close relationship between the structures of $S_{\psi,\delta}(\mathbf{Q}_\infty)$ and $S_{\chi,\varepsilon}(\mathbf{Q}_\infty)$ as $\Lambda_{\mathcal{O}}$ -modules.

The action of $G_{\mathbf{Q}}$ on μ_{p^∞} is given by $\omega\kappa$, where ω and κ are as defined in the introduction. The character κ factors through the group $\Gamma = \text{Gal}(\mathbf{Q}_\infty/\mathbf{Q})$. We have $D = V/T$ as usual. We define $T^* = \text{Hom}(D, \mu_{p^\infty})$, which is a free \mathcal{O} -module of rank $d = \dim_{\mathcal{F}}(V)$.

Let $V^* = T^* \otimes_{\mathcal{O}} \mathcal{F}$, and let $D^* = V^*/T^*$. The representation space V^* of $G_{\mathbf{Q}}$ is not an Artin representation. However, if we let $\mathcal{F}(\omega\kappa)$ and $\mathcal{F}(\kappa)$ denote the 1-dimensional vector spaces over \mathcal{F} on which $G_{\mathbf{Q}}$ acts by the cyclotomic character $\omega\kappa$ and by κ , respectively, then we have

$$V^* \cong \mathrm{Hom}_{\mathcal{F}}(V, \mathcal{F}(\omega\kappa)) \cong U \otimes_{\mathcal{F}} \mathcal{F}(\kappa).$$

We will just write $V^* \cong U \otimes \kappa$ which we think of as the twist of the Artin representation U by κ . We have a filtration $0 \subset W \subset V$ and the orthogonal complements give a filtration on U and on $U \otimes \kappa$. Proposition 11 in [12] deals with an analogous situation, although the filtrations used there are defined in a different way. The Selmer groups denoted by $S_{V/T}(\mathbf{Q}_{\infty})$ and $S_{V^*/T^*}(\mathbf{Q}_{\infty})$ in that paper are also defined in terms of the filtrations, completely analogously to the definitions here. Proposition 11 implies that if one of those Selmer groups is $\Lambda_{\mathcal{O}}$ -cotorsion, then so is the other. However, κ factors through $\Gamma = \mathrm{Gal}(\mathbf{Q}_{\infty}/\mathbf{Q})$ and so $V^* \cong U$ as a representation space for $G_{\mathbf{Q}_{\infty}}$. It would then follow that $S_{D^*}(\mathbf{Q}_{\infty})$ and $S_{\psi,\delta}(\mathbf{Q}_{\infty})$ are the same as sets. More precisely, for the action of Γ , one has $S_{D^*}(\mathbf{Q}_{\infty}) \cong S_{\psi,\delta}(\mathbf{Q}_{\infty}) \otimes \kappa$. Therefore, the fact that $S_{\psi,\delta}(\mathbf{Q}_{\infty})$ is $\Lambda_{\mathcal{O}}$ -cotorsion implies the same for $S_{D^*}(\mathbf{Q}_{\infty})$ and that in turn implies that $S_D(\mathbf{Q}_{\infty}) = S_{\chi,\varepsilon}(\mathbf{Q}_{\infty})$ is $\Lambda_{\mathcal{O}}$ -cotorsion. \blacksquare

Remark 4.8. With the notation and assumptions as in Proposition 4.7 and its proof, let $X_{\chi,\varepsilon}(\mathbf{Q}_{\infty})$, $X_{D^*}(\mathbf{Q}_{\infty})$, and $X_{\psi,\delta}(\mathbf{Q}_{\infty})$ denote the Pontryagin duals of $S_{\chi,\varepsilon}(\mathbf{Q}_{\infty}) = S_D(\mathbf{Q}_{\infty})$, $S_{D^*}(\mathbf{Q}_{\infty})$, and $S_{\psi,\delta}(\mathbf{Q}_{\infty})$, respectively. They are finitely generated, torsion $\Lambda_{\mathcal{O}}$ -modules. Let $\theta_{\chi,\varepsilon}$, θ_{D^*} , and $\theta_{\psi,\delta}$ denote generators for the corresponding characteristic ideals in $\Lambda_{\mathcal{O}}$. The first two are related (up to a unit) by the involution ι of the \mathcal{O} -algebra $\Lambda_{\mathcal{O}}$ induced by the automorphism $\gamma \mapsto \gamma^{-1}$ of Γ . This is the content of Theorem 2 in [12] whose proof applies without change to the case at hand. Thus, we could simply chose $\theta_{\chi,\varepsilon}$ to be $\theta_{D^*}^{\iota}$. On the other hand, we have $X_{D^*}(\mathbf{Q}_{\infty}) \cong X_{\psi,\delta}(\mathbf{Q}_{\infty}) \otimes \kappa^{-1}$. Thus,

$$X_{\psi,\delta}(\mathbf{Q}_{\infty}) \cong X_{D^*}(\mathbf{Q}_{\infty}) \otimes \kappa$$

as $\Lambda_{\mathcal{O}}$ -modules. There is an automorphism of $\Lambda_{\mathcal{O}}$ induced by the map $\gamma \mapsto \kappa(\gamma)^{-1}\gamma$. If X is any finitely generated, torsion $\Lambda_{\mathcal{O}}$ -module, that automorphism send the characteristic ideal of X to the characteristic ideal of $X \otimes \kappa$. It follows that we can choose $\theta_{\psi,\delta}$ so that

$$\varphi(\theta_{\psi,\delta}) = \varphi\kappa^{-1}(\theta_{D^*}) = \varphi^{-1}\kappa(\theta_{\chi,\varepsilon})$$

for all $\varphi \in \mathrm{Hom}(\Gamma, \overline{\mathbf{Q}}_p^{\times})$. In particular, if taking $\varphi = \kappa^s$, we have

$$(9) \quad \kappa^{1-s}(\theta_{\chi,\varepsilon}) = \kappa^s(\theta_{\psi,\delta})$$

for all $s \in \mathbf{Z}_p$. One especially interesting case is when $d = 2$. Then $d^+(\chi)$ and $d^+(\psi)$ are both equal to 1. Assume that $\varepsilon_{\mathfrak{p}} = \omega$. Then $\Delta_{\mathfrak{p}}$ acts on $U/U^{(\delta_{\mathfrak{p}})}$ trivially. As discussed in remark 4.3, it follows that $\varphi_0(\theta_{\psi,\delta}) = 0$. Therefore, taking $s = 0$ in (9), we find that $\kappa(\theta_{\chi,\omega}) = 0$ too.

Remark 4.9. We return to the case where $d^+ = 1$. The exact sequence in Proposition 3.2 can be extended to \mathbf{Q}_{∞} by taking direct limits. One obtains the following exact sequence of discrete $\Lambda_{\mathcal{O}}$ -modules.

$$(10) \quad \begin{aligned} 0 &\longrightarrow H_{unr}^1(\mathbf{Q}_{\infty}, D) \longrightarrow S_{\chi,\varepsilon}(\mathbf{Q}_{\infty}) \\ &\longrightarrow \mathrm{Hom}_{\mathcal{O}[\Delta]} \left(\mathcal{U}_{\infty,p,\mathcal{O}}^{[\varepsilon](\chi)} / \overline{\mathcal{U}}_{\infty,p,\mathcal{O}}^{[\varepsilon](\chi)}, D \right) \longrightarrow 0. \end{aligned}$$

The intervening $\Lambda_{\mathcal{O}[\Delta]}$ -modules in the $\mathrm{Hom}_{\mathcal{O}[\Delta]}$ term are defined by

$$\mathcal{U}_{\infty,p,\mathcal{O}}^{[\varepsilon]} = \prod_{\mathfrak{p}|p} \mathcal{U}_{\infty,\mathfrak{p},\mathcal{O}}^{(\varepsilon_{\mathfrak{p}})}, \quad \overline{\mathcal{U}}_{\infty,p,\mathcal{O}}^{[\varepsilon]} = \varprojlim_n \overline{\mathcal{U}}_{n,p,\mathcal{O}}^{[\varepsilon]},$$

where the inverse limits are defined by the norm maps. They are finitely generated $\Lambda_{\mathcal{O}}$ -modules. One can then take the χ -components of those modules, although that is unnecessary since the χ -component of D is D itself. Also, $\mathcal{U}_{\infty,p,\mathcal{O}}^{[\varepsilon]}$ is a direct factor in $\mathcal{U}_{\infty,p,\mathcal{O}}$ and $\overline{\mathcal{U}}_{\infty,p,\mathcal{O}}^{[\varepsilon]}$ is the projection of $\overline{\mathcal{U}}_{\infty,p,\mathcal{O}} = \varprojlim_n \overline{\mathcal{U}}_{n,p,\mathcal{O}}$ to that factor.

The term $H_{unr}^1(\mathbf{Q}_{\infty}, D)$ in (10) can be identified with

$$\mathrm{Hom}_{\mathcal{O}[\Delta]} \left((\mathrm{Gal}(L_{\infty}/K_{\infty})_{\mathcal{O}})^{(\chi)}, D \right)$$

where L_{∞} is the maximal abelian unramified pro- p -extension of K_{∞} . Now $\mathrm{Gal}(L_{\infty}/K_{\infty})$ is a finitely generated, torsion Λ -module, a famous theorem of Iwasawa. An immediate consequence is that the Pontryagin dual of $H_{unr}^1(\mathbf{Q}_{\infty}, D)$ is a finitely generated, torsion $\Lambda_{\mathcal{O}}$ -module. Let $\theta_{unr,\chi}$ be a generator of its characteristic ideal. It is worth noting that the characteristic ideal for the $\Lambda_{\mathcal{O}}$ -module $(\mathrm{Gal}(L_{\infty}/K_{\infty})_{\mathcal{O}})^{(\chi)}$ is then generated by $\theta_{unr,\chi}^d$.

One can derive another exact sequence which is somewhat simpler than (10). It will lead to the interpolation property stated in the introduction under certain assumptions. We will discuss this carefully in the sequel to this paper and just state it here. Choose a prime \mathfrak{p} lying above p . We will use the notation $|\cdot|_{\varepsilon_{\mathfrak{p}}}$ to denote the composite map

$$\mathcal{U}_{\infty,p,\mathcal{O}} \rightarrow \mathcal{U}_{\infty,\mathfrak{p},\mathcal{O}} \rightarrow \mathcal{U}_{\infty,\mathfrak{p},\mathcal{O}}^{(\varepsilon_{\mathfrak{p}})}$$

where the two maps are the natural projection maps. The exact sequence is

$$(11) \quad \begin{aligned} 0 \rightarrow H_{unr}^1(\mathbf{Q}_\infty, D) \rightarrow S_{\chi, \varepsilon}(\mathbf{Q}_\infty) \\ \rightarrow \mathrm{Hom}_{\mathcal{O}}\left(\mathcal{U}_{\infty, p, \mathcal{O}}^{(\varepsilon_p)} / \overline{\mathcal{U}}_{\infty, p, \mathcal{O}}^{(\chi)}|_{\varepsilon_p}, D^{(\varepsilon_p)}\right) \rightarrow 0. \end{aligned}$$

Note that the $\mathrm{Hom}_{\mathcal{O}}$ term in the above exact sequence is isomorphic to the Pontryagin dual of $\mathcal{U}_{\infty, p, \mathcal{O}}^{(\varepsilon_p)} / \overline{\mathcal{U}}_{\infty, p, \mathcal{O}}^{(\chi)}|_{\varepsilon_p}$ as a $\Lambda_{\mathcal{O}}$ -module.

§5. The p -adic L -function of an Artin representation

We will discuss the construction of a p -adic L -function for the representation ρ in this section. We assume that $d = 2$. The construction relies on Hida theory and on the existence of a 2-variable p -adic L -function. As before, we choose a finite set Σ of primes containing p and ∞ and all the primes ramified in K/\mathbf{Q} . We can then regard ρ as a 2-dimensional representation of $\mathrm{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$. The basic idea is to embed ρ into a p -adic family of 2-dimensional representations ρ_k of $\mathrm{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$ for which a p -adic L -function is known to exist, and then take the limit as ρ_k tends towards ρ . Roughly speaking, for $k \geq 2$, ρ_k will be the p -adic representation corresponding to a modular form of weight k which is ordinary at p and the corresponding p -adic L -function is now classical. (See [26].)

Since $d^+ = 1$, it follows that ρ is odd. The Artin conjecture then implies that ρ is associated to a modular form f of weight 1 for $\Gamma_1(N)$, where N is the Artin conductor of ρ . Thus, f is a normalized newform of weight 1 with q -expansion $f = \sum a_n q^n$ and level N . By work of Langlands, Tunnell, Weil, and Hecke, this is known whenever the image of ρ inside $GL_2(\mathcal{O})$ is a solvable group. In the non-solvable case (i.e., when ρ is *icosahedral*), there are also partial results of Buzzard, Dickinson, Sheppard-Barron, and Taylor. For all of this, see [3] and the references there. We assume from now on that ρ is modular of level N in the above sense.

Recall that we have fixed embeddings σ_∞ and σ_p of $\overline{\mathbf{Q}}$ into \mathbf{C} and into $\overline{\mathbf{Q}}_p$, respectively. The coefficients a_n of f are algebraic integers in \mathbf{C} and can be regarded as elements of $\overline{\mathbf{Q}}_p$. In addition, if q is any prime and \mathfrak{q} is a prime of K lying above q , and $\Delta_{\mathfrak{q}}$ denotes the corresponding decomposition subgroup of Δ , then the restriction of a Frobenius automorphism for \mathfrak{q} to the maximal subspace of V on which the inertia subgroup for \mathfrak{q} acts trivially has trace a_q . Thus, the a_q 's are in \mathcal{O} for

all primes q . Furthermore, for $q \nmid N$, it is clear that $\det(\rho(\text{Frob}_q))$ is in \mathcal{O}^\times . It follows that $a_n \in \mathcal{O}$ for all $n \geq 1$.

Recall that T denotes a Δ -invariant \mathcal{O} -lattice in V . Since Hypothesis A is assumed to be satisfied, for our fixed prime \mathfrak{p} of K lying over p and our choice of $\varepsilon_{\mathfrak{p}}$, there is an exact sequence

$$(12) \quad 0 \rightarrow \mathcal{O}(\varepsilon_{\mathfrak{p}}) \rightarrow T \rightarrow \mathcal{O}(\varepsilon'_{\mathfrak{p}}) \rightarrow 0,$$

of $\mathcal{O}[\Delta_{\mathfrak{p}}]$ -modules, where $\varepsilon_{\mathfrak{p}} \neq \varepsilon'_{\mathfrak{p}}$. As in the introduction, $\varepsilon_{\mathfrak{p}}$ and $\varepsilon'_{\mathfrak{p}}$ are determined by certain characters ε and ε' of $\mathcal{G} = \text{Gal}(\mathcal{K}/\mathbf{Q}_p)$. We assume at first that ρ is ordinary in the sense that ε' is unramified. Then there exists a modular eigenform $f_1 = \sum b_n q^n$ of weight 1 with the property that $b_n = a_n$ if $(n, p) = 1$, but $b_p = \varepsilon'(p)$. The form f_1 will be a p -stabilized newform in the sense of [33], meaning that it has level N if $p \nmid N$ or level Np otherwise. If p divides N , then $f = f_1$. Since the values of ε , and hence ε' , are in \mathcal{O} , it follows that the $b_n \in \mathcal{O}$ for all $n \geq 1$.

The maximal ideal of \mathcal{O} is $p\mathcal{O}$. Since ρ is absolutely irreducible, and $p \nmid |\Delta|$, it follows that the residual representation $\bar{\rho}$ obtained by the canonical reduction map $GL_2(\mathcal{O}) \rightarrow GL_2(\mathcal{O}/p\mathcal{O})$ is also absolutely irreducible. Furthermore, $\bar{\rho}$ is also p -distinguished (i.e., the reductions $\bar{\varepsilon}$, and $\bar{\varepsilon}'$ are distinct). Using fundamental ideas due to Hida, Wiles has constructed a deformation of ρ in this setting, whose description we briefly recall. For background in Hida's theory, we refer the reader to [20], [33], and [7].

(a) The local ring $\mathbf{h}_{\mathfrak{m}}$: Let \mathbf{h}_{∞} denote Hida's universal ordinary Hecke algebra of level M , where M is the prime-to- p part of N . It is a finite flat algebra over the subring $\Lambda' = \mathbf{Z}_p[[\Gamma']]$, where $\Gamma' \cong 1 + p\mathbf{Z}_p$, considered as a subgroup of the group of diamond operators, and so Λ' is isomorphic to a formal power series ring $\mathbf{Z}_p[[T]]$. Now f_1 determines a maximal ideal \mathfrak{m} of \mathbf{h}_{∞} ; we write $\mathbf{h}_{\mathfrak{m}}$ for the completion of \mathbf{h}_{∞} at \mathfrak{m} . Since the coefficients of f_1 generate the ring \mathcal{O} and that ring is generated by roots of unity of order prime to p , one sees that $\mathbf{h}_{\mathfrak{m}}$ contains \mathcal{O} as a subring. Thus, $\mathbf{h}_{\mathfrak{m}}$ also contains the subring $\Lambda'_{\mathcal{O}} \cong \mathcal{O}[[\Gamma']]$.

(b) The Galois representation $\check{\rho}$: Since $\bar{\rho}$ is irreducible and p -distinguished, it follows from the work of Mazur, Ribet, Wiles, and others (see [35], Corollary 2 to Theorem 2.1 on page 482) that the localization $\mathbf{h}_{\mathfrak{m}}$ is a Gorenstein ring. Hida's construction then implies that there exists a representation $\check{\rho} : \text{Gal}(\mathbf{Q}_{\Sigma}/\mathbf{Q}) \rightarrow GL_2(\mathbf{h}_{\mathfrak{m}})$. Wiles has shown in [33] that there exists a weight-one prime ideal $P_1 = P_{\rho} \in \text{Spec}(\mathbf{h}_{\mathfrak{m}})$ such that $\check{\rho}$ specializes to ρ at the point P_{ρ} . If $q \nmid Np$, then the trace of a

Frobenius element $\text{Frob}(q)$ for a prime above q is given by the Hecke operator T_q regarded as an element of \mathbf{h}_m . Furthermore, we identify $G_{\mathbf{Q}_p}$ with a subgroup of $G_{\mathbf{Q}}$ by the fixed embedding σ_p (which determines \mathfrak{p}). Let \mathbf{T} denote a realization of $\check{\rho}$. Then \mathbf{T} is a free \mathbf{h}_m -module of rank 2 and there exists a $G_{\mathbf{Q}_p}$ -stable \mathbf{h}_m -submodule $\mathbf{T}_0 \subset \mathbf{T}$ such that \mathbf{T}/\mathbf{T}_0 is unramified at p . Both \mathbf{T}_0 and \mathbf{T}/\mathbf{T}_0 are free \mathbf{h}_m -modules of rank 1. The eigenvalue of $\text{Frob}(p)$ on \mathbf{T}/\mathbf{T}_0 is given by the Hecke operator U_p . Furthermore, the residual representations for \mathbf{T}_0 and \mathbf{T}/\mathbf{T}_0 are $\varepsilon_{\mathfrak{p}}$ and $\varepsilon'_{\mathfrak{p}}$, respectively. The determinant of $\check{\rho}$ is a homomorphism $\text{Gal}(\mathbf{Q}_{\Sigma}/\mathbf{Q}) \rightarrow \Lambda'_{\mathcal{O}}^{\times}$.

(c) Specialization: Recall that a point $\phi : R \rightarrow \mathbf{C}_p$ of a Λ' -algebra R is said to be arithmetic of weight $k \in \mathbf{Z}$ if the restriction $\phi|_{\Lambda'}$ is the \mathcal{O} -algebra homomorphism induced by a character $\mathbf{Z}_p^{\times} \rightarrow \overline{\mathbf{Q}}_p^{\times}$ of the form $a \mapsto \psi(a)a^k$, where ψ is a finite-order character. Note that if ϕ has values in \mathcal{O} , then ψ must have order prime to p . For each arithmetic point $\phi_k \in \text{Spec}(\mathbf{h}_m)$ of weight $k \geq 2$, the specialization of $\check{\rho}$ to ϕ_k is the representation associated by Deligne to a certain p -ordinary modular form $f_k = f_k(\phi_k)$.

(d) Λ' -adic forms: Finally, we recall that a Λ' -adic form \mathfrak{F} of tame level M with coefficients in a finite Λ' -algebra R is defined to be a formal power series

$$\mathfrak{F} = \sum a_n(\mathfrak{F})q^n \in R[[q]]$$

such that, for almost all arithmetic points $\phi_k \in \text{Spec}(R)$ with $k \geq 2$, the specialization $\mathfrak{F}(\phi_k)$ is the q -expansion of a p -stabilized newform of weight k and level Mp^r , for suitable r . We say that the form \mathfrak{F} contains the classical form f_k of weight $k \geq 1$ and level Mp^r if there exists an arithmetic point ϕ_k of weight k such that $\mathfrak{F}(\phi_k)$ is the q -expansion of f_k . The form \mathfrak{F} is said to be *ordinary* if $e\mathfrak{F} = \mathfrak{F}$, where e is Hida's ordinary projector. From now on, we only consider ordinary forms. In the ordinary case, a Λ' -adic eigenform with coefficients in R is the same thing as a Λ' -algebra homomorphism $\mathbf{h}_{\infty} \rightarrow R$. We will assume that the homomorphism factors through \mathbf{h}_m . We can assume that R is generated by the coefficients of \mathfrak{F} and hence that the homomorphism is surjective. We will also generally assume that R is a domain and so $R \cong \mathbf{h}_{\infty}/\mathfrak{a}$, where \mathfrak{a} is a prime ideal in \mathbf{h}_{∞} of height 0 (i.e., a minimal prime ideal of \mathbf{h}_{∞}). Thus, R is an irreducible component in the Λ' -algebra \mathbf{h}_{∞} . It obviously has characteristic 0 and contains Λ' as a subring.

We write \mathfrak{M}_R for the module of ordinary Λ' -adic modular forms with coefficients in the Λ' -algebra R . Then \mathfrak{M}_R is a finite free R -module, and, if $\mathfrak{M} = \mathfrak{M}_{\Lambda'}$, then there is an isomorphism $\mathfrak{M}_R = \mathfrak{M} \otimes_{\Lambda'} R$. The ring

\mathbf{h}_∞ acts on \mathfrak{M} , and the module $(\mathfrak{M}_R)_\mathfrak{m}$ is canonically isomorphic to the $\mathbf{h}_\mathfrak{m}$ -module $\mathrm{Hom}_R(\mathbf{h}_\mathfrak{m} \otimes_{\Lambda'} R, R)$.

With these notions in hand, we can state a key result.

Theorem 5.1. (Wiles) *Suppose that f_1 is the p -stabilized form associated to ρ and the unramified character ε' . Then there exists an ordinary Λ' -adic eigenform \mathfrak{F} , defined over an extension R of Λ' , which specializes to f_1 in weight 1.*

Note that in the above theorem we can assume that R is a domain and is an irreducible component in the $\Lambda'_\mathcal{O}$ -algebra $\mathbf{h}_\mathfrak{m}$. Now $\varepsilon \neq \varepsilon'$ since we are assuming that Hypothesis A holds for ρ . If ε is unramified, then one can reverse the roles of ε and ε' , obtaining a different Λ' -adic eigenform (which would correspond to a different choice of \mathfrak{m} and a different R). However, for our fixed choice of ε , there is a uniqueness result which is very important for our purpose.

Theorem 5.2. (Greenberg-Vatsal, Belläiche-Dimitrov) *Suppose that a choice of ε is fixed and that Hypothesis A holds. Then the form \mathfrak{F} is unique.*

In particular, \mathfrak{m} and R are uniquely determined by χ and ε . A version of this theorem was discovered by the authors of this paper in 1997, but never published. We needed the restrictive assumption that ε'/ε is not of order 2. It was subsequently rediscovered by Belläiche-Dimitrov [1], who actually proved the result without that assumption.

It is important to observe that the residue ring of $\mathbf{h}_\mathfrak{m}$ at the prime ideal P_ρ is exactly equal to \mathcal{O} , rather than some larger ring \mathcal{O}' . Recall that \mathcal{O} is generated over \mathbf{Z}_p by the values of χ and ε , the ring of integers in $\mathcal{F} = \mathbf{Q}_p(\chi, \varepsilon)$. We justify the claim as follows. The Hecke algebra $\mathbf{h}_\mathfrak{m}$ is generated by the Hecke operators and the diamond operators. We must show that the images of these generators are in \mathcal{O} .

For the diamond operators, this is clear since the values of $\det \circ \rho$ are in that ring. Now the image of any Hecke operator modulo P_ρ is the eigenvalue of the corresponding Hecke operator for the weight 1 form associated to ρ . We have to show that these eigenvalues lie in \mathcal{O} . For any prime q , let \mathfrak{q} be a prime of K lying over q . For the Hecke operator T_q with $q \nmid Np$, the eigenvalue of T_q is the trace of $\rho(\mathrm{Frob}(\mathfrak{q}))$, which lies in \mathcal{O} by definition. If q divides Np , then we argue as follows. As explained in the introduction, ρ can be realized in a 2-dimensional vector space V over the field \mathcal{F} . If $q \nmid p$, let $I_\mathfrak{q}$ be the inertia subgroup of Δ for the prime \mathfrak{q} . Then $V^{I_\mathfrak{q}}$ is an \mathcal{F} -subspace of V of dimension ≤ 1 . If $V^{I_\mathfrak{q}}$

is 1-dimensional, then the eigenvalue of U_q is equal to the eigenvalue of Frobenius on that subspace. That eigenvalue obviously lies in \mathcal{F} , and hence in \mathcal{O} . If that subspace has dimension 0, then the image of U_q is 0. Finally, U_p acts on f_1 with eigenvalue $\varepsilon'_p(\text{Frob}(\mathfrak{p}))$. Since $\chi|_{\Delta_p} = \varepsilon_p + \varepsilon'_p$, it follows that the values of ε'_p are in \mathcal{O} . Thus, all the eigenvalues of the Hecke operators are indeed in \mathcal{O} .

The above discussion yields a representation

$$\check{\rho} : \text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q}) \rightarrow GL_2(\mathbf{h}_m)$$

which specializes to ρ at an arithmetic specialization of weight 1. We can compose with the surjective homomorphism $\mathbf{h}_m \rightarrow R$ to obtain a representation

$$\tilde{\rho} : \text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q}) \rightarrow GL_2(R).$$

Furthermore, there is a homomorphism $\phi_\rho : R \rightarrow \mathcal{O}$ such that $\phi_\rho \circ \tilde{\rho} = \rho$. Here we are also using the notation ϕ_ρ for the induced continuous group homomorphism $GL_2(R) \rightarrow GL_2(\mathcal{O})$. Such representation $\check{\rho}$ and $\tilde{\rho}$ are called deformation of ρ . The rings \mathbf{h}_m and R are the corresponding deformation rings. Each deformation has infinitely many motivic points, namely the points corresponding to forms of higher weight. The deformation $\tilde{\rho}$ is the one we will concentrate on. These motivic points have well-defined p -adic L -functions (up to multiplying by units in the corresponding residue rings of R). We can therefore define a p -adic L -function for ρ by continuity. We now explain this more precisely.

Thus we are led to describe the p -adic L -function associated to the representation $\tilde{\rho}$. It will correspond to an element in $R[[\Gamma]]$, where $\Gamma = \text{Gal}(\mathbf{Q}_\infty/\mathbf{Q})$ is as in the previous sections. This L -function was constructed by Kitagawa [23] and by Greenberg-Stevens [19]. We want to recall the modular-symbol construction of Greenberg-Stevens, as this is the most convenient for us. Our discussion here is only a summary, and we refer the reader to the original paper of Greenberg and Stevens for a detailed exposition. For each positive integer r , let Γ_r denote the group $\Gamma_1(Np^r)$. Let \mathbf{M} denote the inverse limit of the cohomology groups $eH_{\text{par}}^1(\Gamma_r, \mathbf{Z}_p)$; then \mathbf{M} is a module over the Hecke algebra \mathbf{h}_∞ . The matrix $\iota = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ induces a Hecke-equivariant involution on \mathbf{M} , and we write \mathbf{M}^\pm for the \pm -eigenspaces under this action. Evidently we have the decomposition $\mathbf{M} = \mathbf{M}^+ \oplus \mathbf{M}^-$. Since the maximal ideal \mathfrak{m} corresponds to an irreducible representation, it follows from Thm. 4.3

and Lemma 6.9 of [19] that we have an isomorphism of \mathbf{h}_m -modules

$$\mathbf{M}_m = \text{Symb}_{\Gamma_1}(\mathbf{D})_m,$$

where \mathbf{D} is the space of \mathbf{Z}_p -valued measures on the set of primitive elements of \mathbf{Z}_p^2 , and $\text{Symb}_{\Gamma_1}(\mathbf{D})$ denotes the group of modular symbols over the group Γ_1 with values in \mathbf{D} . Furthermore, since \mathbf{h}_m is Gorenstein, a duality argument shows that, for any choice $\alpha = \pm$ of the sign, there are isomorphisms of Hecke modules $\mathfrak{M}_m = \mathbf{M}_m^\alpha = \mathbf{h}_m$.

Now let \mathfrak{F} be a Λ' -adic eigenform form, defined over the ring R , such that \mathfrak{F} specializes to $f_1 = f_\rho$ in weight 1. The choice of ε'_p made in (12) is implicit in the choice of \mathfrak{F} . Thus we have a homomorphism $\mathbf{h}_m \rightarrow R$ such that the kernel \mathfrak{P} is a minimal prime contained in the height 1 prime ideal P_ρ of \mathbf{h}_m . Extending scalars via $\Lambda' \rightarrow R$, we obtain isomorphisms $\mathfrak{M}_m \otimes_{\Lambda'} R = \mathbf{M}_m^\pm \otimes_{\Lambda'} R = \mathbf{h}_m \otimes_{\Lambda'} R$. Let $\delta^\alpha(\mathfrak{F})$ be the image of $\mathfrak{F} \in \mathfrak{M} \otimes_{\Lambda'} R$ in the module $\mathbf{M}_m^\pm \otimes_{\Lambda'} R$, and let $\delta(\mathfrak{F}) = \delta^+(\mathfrak{F}) + \delta^-(\mathfrak{F})$. Finally, let $\mathfrak{L}(\mathfrak{F}) = \delta(\mathfrak{F})(\{0\} - \{\infty\}) \in \mathbf{D} \otimes_{\Lambda'} R$ denote the special value ([19] Sec. 4.10). Then $\mathfrak{L}(\mathfrak{F})$ is an element of $\mathbf{D} \otimes_{\Lambda'} R$, and the construction of [19], equations 5.2a and 5.4, gives an element

$$(13) \quad L_p(\mathfrak{F}) \in R[[\mathbf{Z}_p^\times]]$$

known as the standard two-variable p -adic L -function associated to \mathfrak{F} . The terminology is justified as follows. Let P_k denote an arithmetic point in $\text{Spec}(R)$, of weight $k \geq 2$. Then P_k induces a specialization map $\phi_k : R \rightarrow \mathcal{O}$, as explained above. We may specialize $L_p(\mathfrak{F})$ via ϕ_k to obtain a measure $L_p(\mathfrak{F})(P_k)$ on \mathbf{Z}_p^\times ; one sees from the definitions that, if f_k denotes the specialization of \mathcal{F} to P_k , then the measure $L_p(\mathfrak{F})(P_k)$ is a p -adic L -function for f_k . Specifically, if ψ is any finite order character of \mathbf{Z}_p^\times , then $\mu_k = L_p(\mathfrak{F})(P_k)$ satisfies the property

$$(14) \quad \int_{\mathbf{Z}_p^\times} \psi d\mu_k = e_p(f_k, \psi) \tau(\psi) \frac{L(f_k \otimes \psi^{-1}, 1)}{(-2\pi i) \Omega_k^\alpha}$$

for certain periods Ω_k^\pm . Here $\tau(\psi)$ is the standard Gauss sum for ψ and

$$e_p(f_k, \psi) = \alpha_k^{-n} (1 - \alpha_k^{-1} \psi(p)),$$

where α_k is the (unit) eigenvalue of the U_p operator on f_k . The quantity $L(f_k \otimes \psi^{-1}, 1)$ denotes the standard L -function of f_k , twisted by the finite order character ψ^{-1} and evaluated at $s = 1$. A similar but more complicated formula holds for the integral of ψz^{j-1} , where j is any integer in the critical range $1 \leq j \leq k - 1$. It is essentially the formula in the proposition of Section 14 in [26].

It follows easily from Hida's control theorems [20] that the periods Ω_k^α are the *canonical periods* of f_k introduced in [31]. These periods are defined up to p -adic units. The sign $\alpha = \pm$ in the period is determined by the parity of ψ .

We can identify \mathbf{Z}_p^\times with $\text{Gal}(\mathbf{Q}(\mu_{p^\infty})/\mathbf{Q})$ in the usual way. Then $1 + p\mathbf{Z}_p$ is identified with Γ . The characters ψ of \mathbf{Z}_p^\times can be regarded as characters of that Galois group. We have $\psi = \omega^t \varphi$ where $0 \leq t \leq p-2$ and φ is a character of finite order of Γ .

The ring $R[[\mathbf{Z}_p^\times]]$ has a natural R -algebra involution ι defined by $\iota(z) = z^{-1}$ for all $z \in \mathbf{Z}_p^\times$. For our purpose, we consider the element $\iota(L_p(\mathfrak{F}))$ in $R[[\mathbf{Z}_p^\times]]$. If we specialize via ϕ_k as above, we obtain a measure μ_k' . Integrating ψ against μ_k' gives a formula like that in (14), but with ψ replaced by ψ^{-1} on the right side. In particular, the L -value occurring in the formula would now be $L(f_k \otimes \psi, 1) = L(f_k, \psi, 1)$. Now $\mathbf{Z}_p^\times \cong \mathbf{F}_p^\times \times (1 + p\mathbf{Z}_p)$ and so one has the decomposition

$$R[[\mathbf{Z}_p^\times]] \cong \bigoplus_{t=0}^{p-2} e_{\omega^t} R[[\mathbf{Z}_p^\times]]$$

where e_{ω^t} is the idempotent for ω^t . Each summand is isomorphic to $R[[1 + p\mathbf{Z}_p]]$ which we identify with $R[[\Gamma]]$. If we project $\iota(L_p(\mathfrak{F}))$ to the ω^t -component for a fixed t , we get an element $\Theta_t \in R[[\Gamma]]$. Specializing via ϕ_k , we get an element $\theta_{t,k} \in (R/P_k)[[\Gamma]]$. If we let $\psi = \omega^t \varphi$, where φ is a character of Γ , then integrating ψ against μ_k' gives the value $\varphi(\theta_{t,k})$. Note that $L(f_k \otimes \psi, 1) = L(f_k \otimes \omega^t, \varphi, 1)$.

Recall that we assumed earlier that ρ is ordinary in the sense that there is an exact sequence (12) with ε_p' unramified. However, in general, the action of $G_{\mathbf{Q}_p}$ on $T/\mathcal{O}(\varepsilon_p)$ might be ramified. One could then replace ρ by $\rho \otimes \omega^{-t}$ for some t , $0 \leq t \leq p-2$, to obtain an ordinary Artin representation. One can apply Wiles theorem to the corresponding weight 1 form f_1 and then tensor by ω^t obtaining the family $f_k \otimes \omega^t$ of modular forms which specializes in weight 1 to $f_\rho = f_1 \otimes \omega^t$. With this motivation in mind, it is natural to define $\theta_{\chi, \varepsilon}^{an}$ to be the image of Θ_t by the specialization map $\phi_\rho : R[[\Gamma]] \rightarrow \mathcal{O}[[\Gamma]] = \Lambda_{\mathcal{O}}$. This is the element alluded to in the introduction and is well-defined up to multiplication by an element of \mathcal{O}^\times .

Remark 5.3. As we have remarked several times already, the p -adic L -function $L_p(s, \chi, \varepsilon)$ depends on the choice of ε , in the case $(N, p) = 1$. Furthermore, the L -function is only defined up to a p -adic unit, owing to the indeterminacy of the complex periods in the definition of the 2-variable p -adic L -function. Finally, it is *not at all* clear that $L_p(s, \chi, \varepsilon)$

is nonzero. However, in view of Theorem 3 of the introduction, it seems reasonable to expect that $L_p(s, \chi, \varepsilon)$ should be related to the Coleman series of the norm-compatible family of units given in that theorem, and that a formula in terms of logarithms of global units similar to the one in Theorem 3 should hold. We will take up this subject in the sequel to this paper.

Remark 5.4. The modular cuspforms of weight one may be classified according to the isomorphism class of the image of the associated representation ρ in $PGL_2(\mathbf{C})$. The possible image G of Δ is isomorphic to one of the following: a dihedral group, A_4 , S_4 , or A_5 . In the event that G is dihedral, it is induced from a character of a quadratic extension of \mathbf{Q} . Then ρ is called CM (complex multiplication) or RM (real multiplication), according to whether it is induced from an imaginary quadratic field or not. An alternative approach to the p -adic L -function in this case is via the Katz 2-variable p -adic L -function; the connection with global elliptic units is then given by the p -adic Kronecker limit formula (see [5], Theorem 5.2 and the subsequent discussion). We refer the reader to Ferrara [9] for more details. When ρ has real multiplication, Ferrara has also compiled numerical evidence relating $\phi(\theta_{\chi, \varepsilon}^{an})$ to global units (in fact, Stark units) in abelian extensions of real quadratic fields when ϕ has order p . On the other hand, when ρ is an exotic form of non-dihedral type, absolutely nothing is known.

Remark 5.5. As stated in the introduction, the natural formulation of an Iwasawa main conjecture is the assertion that $\theta_{\chi, \varepsilon}^{al}$ and $\theta_{\chi, \varepsilon}^{an}$ generate the same ideal in $\Lambda_{\mathcal{O}}$. As mentioned in remark 5.3, we cannot even rule out the possibility that $\theta_{\chi, \varepsilon}^{an} = 0$. The fact that $\theta_{\chi, \varepsilon}^{al} \neq 0$ is the content of Proposition 4.2. It seems reasonable to believe that the main conjecture in weight 1 should follow from the main conjecture in higher weights. Some ideas in [28] would possibly be useful in proving a result of that kind. However, regrettably, the proof of the main conjecture in higher weight by Urban-Skinner [29] does not apply to the case at hand because those authors require the presence of a prime $q|M$ such that q is a prime of multiplicative reduction for \mathfrak{F} . Theorem 5.6 below implies that the form \mathfrak{F} admits no such prime q . On the other hand, it appears that the existence of such a q is only required to ensure that a certain anticyclotomic μ -invariant vanishes, and this can also be proven when q is a prime of supercuspidal reduction, so it may be that the result of Skinner-Urban applies when there exists a prime q such that the image of a decomposition group at q under ρ is non-abelian and irreducible. We have not pursued this idea.

We have assumed $d = 2$ until now. We close this paper with some general comments for arbitrary d . Let R be a complete Noetherian local ring with finite residue field of characteristic p . Assume that we have a continuous representation $\tilde{\rho} : \text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q}) \rightarrow GL_d(R)$, together with a continuous, surjective ring homomorphism $\phi_\rho : R \rightarrow \mathcal{O}$ such that $\phi_\rho \circ \tilde{\rho} = \rho$. That is, $\tilde{\rho}$ is a deformation of ρ . The ring R is the corresponding deformation ring. Since \mathcal{O} is generated over \mathbf{Z}_p by roots of unity of order prime to p , it is easy to see that R contains a subring which we can identify with \mathcal{O} . Therefore, R is an \mathcal{O} -algebra and ϕ_ρ is an \mathcal{O} -algebra homomorphism. Furthermore, the residue ring of R is the same as that of \mathcal{O} , namely $\mathcal{O}/p\mathcal{O}$.

We write P_ρ for the kernel of the ring homomorphism ϕ_ρ , a prime ideal in R . If P is any other prime ideal of R such that $R/P \cong \mathcal{O}'$, where \mathcal{O}' is a finite integral extension of \mathcal{O} , then composing $\tilde{\rho}$ with the group homomorphism $GL_d(R) \rightarrow GL_d(\mathcal{O}')$ defined by reducing modulo P gives a d -dimensional representation of $\text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$ over \mathcal{O}' . Thus, we get a family of representations indexed by such prime ideals P of R . Of course, $\tilde{\rho} \bmod P$ is not necessarily an Artin representation. However, if R is a domain, then the following proposition implies that it has one property in common with ρ . Namely, $\tilde{\rho} \bmod P$ is potentially unramified at all primes not dividing p . A similar result in the case $d = 2$ is contained in Theorem 2.3 in [28].

Proposition 5.6. *Let $\tilde{\rho}$ be a deformation of the Artin representation ρ . Assume that the corresponding deformation ring R is an integral domain. Suppose that q is a finite prime in Σ and that $q \neq p$. Let I_v be the inertia subgroup of $\text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$ for a prime v of \mathbf{Q}_Σ lying over q . Then the image of I_v under $\tilde{\rho}$ is finite.*

Proof. Recall that ρ factors through $\text{Gal}(K/\mathbf{Q})$. Thus, the residual representation for $\tilde{\rho}$ also factors through $\text{Gal}(K/\mathbf{Q})$. It follows that the image of $\text{Gal}(\mathbf{Q}_\Sigma/K)$ under $\tilde{\rho}$ is a pro- p group and hence that $\tilde{\rho}$ factors through $\text{Gal}(M/\mathbf{Q})$, where M is the maximal pro- p -extension of K contained in \mathbf{Q}_Σ . Therefore, since $q \neq p$, the restriction $\tilde{\rho}|_{G_{K_v}}$ factors through $\text{Gal}(K_v^{tr,p}/K_v)$, where $K_v^{tr,p}$ denotes the maximal pro- p tamely ramified extension of K_v . Enlarging K if necessary, we can assume that K , and hence K_v , contains μ_p . Now $K_v^{tr,p}$ contains the field $K_v^{unr,p}$, the maximal unramified pro- p extension of K_v . In fact, $K_v^{unr,p} = K_v(\mu_{p^\infty})$ and $K_v^{tr,p} = K_v(\mu_{p^\infty}, \sqrt[p^\infty]{q})$. Furthermore, both $\text{Gal}(K_v^{unr,p}/K_v)$ and $\text{Gal}(K_v^{tr,p}/K_v^{unr,p})$ are isomorphic to \mathbf{Z}_p . The Frobenius element of $\text{Gal}(K_v^{unr,p}/K_v)$ acts on $\text{Gal}(K_v^{tr,p}/K_v^{unr,p})$ by conjugation. This action is given by $x \mapsto x^{\mathcal{N}(v)}$ for all $x \in \text{Gal}(K_v^{tr,p}/K_v^{unr,p})$. Here $\mathcal{N}(v) = q^f$,

where f is the residue field degree for K_v/\mathbf{Q}_q . Thus, $\text{Gal}(K_v^{tr,p}/K_v)$ is an extension of \mathbf{Z}_p by $\mathbf{Z}_p(1)$.

Let τ denote a topological generator of $\text{Gal}(K_v^{tr,p}/K_v^{unr,p})$. The restriction of $\tilde{\rho}$ to the inertia subgroup $\text{Gal}(K_v^{tr,p}/K_v^{unr,p})$ is determined by the matrix $\tilde{\rho}(\tau)$. The eigenvalues of this matrix are in some extension of the fraction field of R . In addition, if α is one of those eigenvalues, then so is α^{q^f} . Since the matrix has only finitely many eigenvalues, it follows that these eigenvalues are roots of unity. Therefore, for some $t \geq 1$, the matrix $A = \tilde{\rho}(\tau^t)$ is unipotent. If we again enlarge the field K , we can assume that $t = 1$ i.e., that the eigenvalues of $\tilde{\rho}(\tau)$ are all equal to 1. Of course, f may change too.

Let \mathbf{F} denote the fraction field of R . Thus, $\text{Gal}(\mathbf{Q}_\Sigma/\mathbf{Q})$ acts on $\mathbf{V} = \mathbf{F}^d$ via $\tilde{\rho}$. Let $N = A - I_d$, a nilpotent matrix with entries in R . There is a filtration on \mathbf{V} defined by the subspaces $\mathbf{W}_k = \text{Ker}(N^k)$ for $k \geq 0$. Here $\mathbf{W}_0 = 0$ and $\mathbf{W}_t = \mathbf{V}$ if t is sufficiently large. These subspaces are G_{K_v} -invariant. The action of $\text{Gal}(K_v^{tr,p}/K_v)$ on the subquotients $\mathbf{W}_{k+1}/\mathbf{W}_k$ factors through $\text{Gal}(K_v^{unr,p}/K_v)$. We will prove that $\mathbf{W}_1 = \mathbf{V}$, i.e., that N annihilates \mathbf{V} .

Assume to the contrary that \mathbf{W}_1 is a proper subspace of \mathbf{W}_2 . Then, $N\mathbf{W}_2$ is a nontrivial subspace of \mathbf{W}_1 . We can lift the Frobenius element of $\text{Gal}(K_v^{unr,p}/K_v)$ to an element of $\text{Gal}(K_v^{tr,p}/K_v)$ which then acts on \mathbf{V} by a matrix B with entries in \mathbf{h} . Furthermore, we have $BAB^{-1} = A^{q^f}$. It follows that

$$BNB^{-1} = q^f N + CN^2$$

where C is a matrix with entries in R . One sees easily that \mathbf{W}_1 and \mathbf{W}_2 are invariant under multiplication by B and by N . Denoting the restrictions of B and N to \mathbf{W}_2 by B_2 and N_2 , respectively, we have $B_2N_2 = q^f N_2B_2$. One sees from this that if β is an eigenvalue for the action of B on $\mathbf{W}_2/\mathbf{W}_1$, then $q^f\beta$ is an eigenvalue for the action of B on \mathbf{W}_1 . Thus, B has two eigenvalues whose ratio is q^f .

Applying the homomorphism ϕ_ρ to B , we obtain a matrix with entries in \mathcal{O} which again has two eigenvalues whose ratio is q^f . However, this matrix is in the image of ρ , a finite subgroup of $GL_d(\mathcal{O})$, and its eigenvalues are roots of unity. This is a contradiction and so we have $\mathbf{W}_1 = \mathbf{V}$. It follows that τ acts trivially on \mathbf{V} . Therefore, the image of the inertia subgroup of G_K for v under $\tilde{\rho}$ is trivial. In the proof, we have possibly replaced K by a finite extension. Hence, for the original K , the image of I_v is finite. \blacksquare

In general, if we try to imitate the $d = 2$ case, there are some natural restrictions on the kinds of deformations of ρ we consider. We are

assuming that ρ satisfies Hypothesis A. The underlying \mathcal{F} -representation space V contains a $G_{\mathbf{Q}}$ -invariant \mathcal{O} -lattice T . Fix a prime \mathfrak{p} of K lying over p . Then $T^{(\varepsilon_{\mathfrak{p}})} \cong \mathcal{O}(\varepsilon_{\mathfrak{p}})$ as $\mathcal{O}[\Delta_{\mathfrak{p}}]$ -modules. Since $p \nmid |\Delta_{\mathfrak{p}}|$, $T/T^{(\varepsilon_{\mathfrak{p}})}$ is a free \mathcal{O} -module of rank $d - 1$.

It is natural to consider deformations of T which simultaneously deform $T^{(\varepsilon_{\mathfrak{p}})}$. To be precise, let \mathfrak{p} be the prime of K induced by the fixed embedding of $\overline{\mathbf{Q}}$ into $\overline{\mathbf{Q}}_p$, as in the introduction. We can then identify $G_{\mathbf{Q}_p}$ with a subgroup of $G_{\mathbf{Q}}$ and restrict $\tilde{\rho}$ to that subgroup. If \mathbf{T} is the underlying free R -module of rank d for $\tilde{\rho}$, we make the following hypothesis.

Hypothesis $\tilde{\mathbf{A}}$: *There exists a $G_{\mathbf{Q}_p}$ -invariant R -submodule \mathbf{T}_0 of \mathbf{T} with the following properties:*

- (i): \mathbf{T}_0 is a free R -module of rank 1,
- (ii): \mathbf{T}/\mathbf{T}_0 is also a free R -module,
- (iii): the image of \mathbf{T}_0 under the map ϕ_{ρ} is $T^{(\varepsilon_{\mathfrak{p}})}$.

Note that the residual representations of $G_{\mathbf{Q}_p}$ for \mathbf{T}_0 and for \mathbf{T}/\mathbf{T}_0 are the same as for $T^{(\varepsilon_{\mathfrak{p}})}$ and for $T/T^{(\varepsilon_{\mathfrak{p}})}$, respectively, and that they have no irreducible constituents in common.

Remark 5.7. Assuming that R is a domain, Hypothesis $\tilde{\mathbf{A}}$ is not as stringent as it might seem at first. As mentioned in the proof of Proposition 5.6, the image of $\text{Gal}(\mathbf{Q}_{\Sigma}/K)$ under $\tilde{\rho}$ is a pro- p group. It follows that the action of $\text{Gal}(\mathbf{Q}_{\Sigma}/\mathbf{Q})$ on \mathbf{T} factors through a group \mathcal{G} which has a normal pro- p subgroup \mathcal{N} and corresponding quotient group $\mathcal{G}/\mathcal{N} \cong \Delta$. Since $|\Delta|$ is prime to p , this group extension splits and \mathcal{G} has a subgroup \mathcal{D} which is mapped isomorphically to Δ under the map $\mathcal{G} \rightarrow \mathcal{G}/\mathcal{N}$. The subgroup \mathcal{D} is not unique, but, for simplicity of notation, we make a choice and identify \mathcal{D} with Δ by that isomorphism. With this identification, $\tilde{\rho}|_{\Delta}$ is closely related to ρ . To be precise, the reduction of $\tilde{\rho}|_{\Delta}$ modulo P_{ρ} coincides with ρ . We let \mathbf{F} be the fraction field of R (as in the proof of Proposition 5.6) and let $\mathbf{V} = \mathbf{T} \otimes_R \mathbf{F}$. Since R contains \mathcal{O} , \mathbf{F} must contain \mathcal{F} . The action of Δ on the \mathbf{F} -vector space \mathbf{V} is a d -dimensional representation over \mathbf{F} and the character of that representation must be χ . Now ρ is a representation of Δ over \mathcal{F} and this \mathbf{F} -representation of Δ must be isomorphic to the extension of scalars of ρ from \mathcal{F} to \mathbf{F} .

Hypothesis A (from the introduction) then has the following consequence for the action of $\Delta_{\mathfrak{p}}$ on \mathbf{V} : There is a 1-dimensional $\Delta_{\mathfrak{p}}$ -invariant \mathbf{F} -subspace $\mathbf{V}^{(\varepsilon_{\mathfrak{p}})}$ on which $\Delta_{\mathfrak{p}}$ acts by $\varepsilon_{\mathfrak{p}}$. Of course, that subspace is

simply $e_{\varepsilon_p} \mathbf{V}$, where $e_{\varepsilon_p} \in \mathcal{O}[\Delta_p]$ is the idempotent corresponding to ε_p . If Hypothesis \tilde{A} is satisfied, then it is clear that $\mathbf{T}_0 = e_{\varepsilon_p} \mathbf{T}$.

These observations lead to the following conclusion. It is sufficient to just assume that \mathbf{V} contains a $G_{\mathbf{Q}_p}$ -invariant \mathbf{F} -subspace \mathbf{V}_0 of dimension 1 such that the action of Δ_p on \mathbf{V}_0 is given by ε_p . That subspace must be $\mathbf{V}^{(\varepsilon_p)}$. Equivalently, it suffices to assume that $\mathbf{V}^{(\varepsilon_p)}$ is actually a $G_{\mathbf{Q}_p}$ -invariant subspace of \mathbf{V} , and not just Δ_p -invariant. Let

$$\mathbf{T}_0 = \mathbf{T} \cap \mathbf{V}_0 = \mathbf{T}^{(\varepsilon_p)},$$

the maximal subset of \mathbf{T} on which Δ_p acts by ε_p . Then \mathbf{T}_0 is a $G_{\mathbf{Q}_p}$ -invariant R -submodule of \mathbf{T} and has rank 1. Since $p \nmid |\Delta_p|$, it follows that $\mathbf{T}_0 = e_{\varepsilon_p} \mathbf{T}$ and that \mathbf{T}_0 is a direct summand in \mathbf{T} as a module over $R[\Delta_p]$. Therefore, \mathbf{T}_0 is a direct summand for \mathbf{T} as a module over R . Since \mathbf{T} is a free R -module, any direct summand for \mathbf{T} is also free. It follows that \mathbf{T}_0 is a free R -module and that the quotient \mathbf{T}/\mathbf{T}_0 is also free (and has rank $d-1$). Furthermore, the image of \mathbf{T}_0 under ϕ_ρ is just $e_{\varepsilon_p} T$ which is $T^{(\varepsilon_p)}$. Thus, Hypothesis \tilde{A} is satisfied.

Assuming that such a $G_{\mathbf{Q}_p}$ -invariant \mathbf{F} -subspace \mathbf{V}_0 does exist, note that the action of $G_{\mathbf{Q}_p}$ on \mathbf{T}_0 is given by a continuous homomorphism

$$\tilde{\varepsilon}_p : G_{\mathbf{Q}_p} \rightarrow R^\times .$$

with the property that $\phi_\rho \circ \tilde{\varepsilon}_p = \varepsilon_p$. In other words, $\tilde{\varepsilon}_p$ is a deformation of the $G_{\mathbf{Q}_p}$ -representation ε_p . It is natural to denote \mathbf{T}_0 by $\mathbf{T}^{(\tilde{\varepsilon}_p)}$. It is indeed the maximal R -submodule of \mathbf{T} on which $G_{\mathbf{Q}_p}$ acts by $\tilde{\varepsilon}_p$.

In addition to Hypothesis \tilde{A} , we would need to assume that the deformation $\tilde{\rho}$ of ρ has a rich supply of “*motivic*” specializations so that one can reasonably expect the existence of a suitable p -adic L -function. We have no way to ensure this in general. However, the remark below describes one case for $d = 3$ where we do have an ample set of motivic specializations.

Remark 5.8. Let ρ be the 2-dimensional representation associated to a modular form of weight 1. Assume that ρ factors through $\Delta = \text{Gal}(K/\mathbf{Q})$ and that $p \nmid |\Delta|$. Then $\text{Ad}^0(\rho) = \text{Sym}^2(\rho) \otimes \det(\rho)^{-1}$ is a 3-dimensional representation factoring through Δ which satisfies $d^+ = 1$. It may or may not satisfy Hypothesis A; it will do so if and only if ε/ε' has order ≥ 3 . If it does satisfy Hypothesis A, then one can take the adjoint of the Λ' -adic Galois representation associated to \mathfrak{F} to get a good deformation of $\text{Ad}^0(\rho)$. The algebraic part of our theory goes through unchanged. The analytic side also goes through, but there is

one important difference: the p -adic L -function has to be defined via continuity from the Schmidt L -function constructed in [4]. The details require more care, since in the dihedral cases the symmetric square is reducible, and in the RM case the corresponding deformation space is not smooth over the weight space at the weight one point of interest. However, it seems reasonable to expect a relationship with units to hold in the CM and exotic A_4 , S_4 and A_5 cases. We omit the details here. Regrettably, this example, and the $d = 2$ example of modular forms, are the only examples of which we are aware where $d^+ = 1$ and sufficiently many motivic deformations exist.

References

- [1] J. Bellaïche, M. Dimitrov, *On the eigencurve at classical weight 1 points*, Duke Math. Jour. **165** (2016), 245-266.
- [2] A. Brumer, *On the units of algebraic number fields*, Mathematika **14** (1967), 121-124.
- [3] K. Buzzard, M. Dickinson, N. Sheppard-Barron, R. Taylor, *On icosahedral Artin representations*, Duke Math Journal **109** (2001), 283-318.
- [4] C.-G. Schmidt, *p -adic measures attached to automorphic representations of $GL(3)$* , Invent. Math., **92**, 1988, 597-631.
- [5] E. de Shalit, *Iwasawa theory of elliptic curves with complex multiplication*, Perspectives in Mathematics, Vol 3, Academic Press, 1987.
- [6] P. Deligne, K. Ribet, *Values of abelian L -functions at negative integers over totally real fields*, Invent. Math. **59** (1980), 227-286.
- [7] M. Emerton, R. Pollack, T. Weston, *Variation of the Iwasawa invariants in a Hida family*, Invent. Math. **163** (2006), 523-580.
- [8] M. Emsalem, H. Kisilevsky, D. Wales, *Indépendance Linéaire sur $\overline{\mathbf{Q}}$ de logarithmes p -adiques de nombres algébriques et rang p -adiques du groupe des unités d'un corps de nombres*, Jour. of Number Theory **19** (1984), 384-391.
- [9] J. Ferrara, *A Stark conjecture for certain p -adic L -functions of quadratic fields*, UCSB Ph.D. thesis in preparation, 2017.
- [10] R. Greenberg, *On p -adic L -functions and cyclotomic fields II*, Nagoya Math. Jour. **67** (1977), 139-158.
- [11] R. Greenberg, *On p -adic Artin L -functions*, Nagoya Math. Jour. **89** (1983), 77-87.
- [12] R. Greenberg, *Iwasawa theory for p -adic representations*, Adv. Stud. Pure Math. **17** (1989), 97-137.
- [13] R. Greenberg, *Iwasawa theory for motives*, in L -functions and Arithmetic, LMS Lecture Note Series **153** (1991), 211-234.
- [14] R. Greenberg, *Iwasawa theory and p -adic deformations of motives*, Proceedings of Symposia in Pure Math. **55 II** (1994), 193-223.

- [15] R. Greenberg, *Introduction to Iwasawa Theory for Elliptic Curves*, LNM **1716**, 51-144.
- [16] R. Greenberg, *On the structure of certain Galois cohomology groups*, Documenta Math. · Extra Coates Volume (2006) 335-391.
- [17] R. Greenberg, *Surjectivity of the global-to-local map defining a Selmer group*, Kyoto J. of Math. **50** (2010), 853-888.
- [18] R. Greenberg, *On the structure of Selmer groups*, in Elliptic Curves, Modular Forms, and Iwasawa Theory, Springer Proceedings in Math. and Stat. **188**, 225-252.
- [19] R. Greenberg, G. Stevens, *p-adic L-functions and p-adic periods of modular forms*, Invent. Math. **111** (1993), 407-447.
- [20] H. Hida, *Galois representations into $GL_2(\mathbf{Z}_p[[X]])$ attached to ordinary cusp forms*, Invent. Math. **85** (1985), 545-613.
- [21] K. Iwasawa, *Lectures on p-adic L-functions*, Ann. Math. Studies **74**, Princeton University Press, 1972.
- [22] K. Iwasawa, *On \mathbf{Z}_ℓ -extensions of algebraic number fields*, Annals of Math. **98** (1973), 246-326.
- [23] K. Kitagawa, *On standard p-adic L-functions of families of elliptic cusp-forms*, in p-adic monodromy and the Birch-Swinnerton-Dyer Conjecture, Contemp. Math. 165, Amer. Math. Soc., 1994.
- [24] T. Kubota, W. Leopoldt, *Eine p-adische Theorie der Zetawerte (Teil I)*, J. Reine Angew. Math. **213** (1964), 328-339.
- [25] H. W. Lenstra, R. J. Schoof, *Primitive normal bases for finite fields*, Math. of Comp. **48** (1987), 217-231.
- [26] B. Mazur, J. Tate, J. Teitelbaum, *On p-adic analogs of the conjectures of Birch and Swinnerton-Dyer*, Invent. Math. **84** (1986), 1-48.
- [27] J. Neukirch, A. Schmidt, K. Wingberg, *Cohomology of Number Fields*, Grundlehren der Math. Wissenschaften **323** (2000), Springer.
- [28] T. Ochiai, *On the two-variable Iwasawa main conjecture*, Compositio Math. **142** (2006), 1157-1200.
- [29] C. Skinner, E. Urban, *The Iwasawa Main Conjectures for GL_2* , Invent. Math., **195**, (2014) 1-277.
- [30] L. Washington, *Introduction to cyclotomic fields*, Graduate Texts in Math. **83**, Springer-Verlag, 1980.
- [31] V. Vatsal, *Iwasawa theory, modular forms, and Artin representations*, Ph.D. thesis, Princeton University, 1997.
- [32] V. Vatsal *Canonical periods and congruence formulae*, Duke Math. J. **98** (1999), 397-419.
- [33] A. Wiles, *On ordinary Λ -adic representations associated to modular forms*, Inv. Math. **94** (1988), 529-573.
- [34] A. Wiles, *The Iwasawa conjecture for totally real fields*, Annals of Math. **131** (1990), 493-540.
- [35] A. Wiles, *Modular Elliptic Curves and Fermat's Last Theorem*, Annals of Math. **141** (1995), 443-551.

Department of Mathematics, University of Washington & Department of Mathematics, University of British Columbia

E-mail address: `greenber@math.washington.edu` & `vastal@math.ubc.ca`